

Homogenization of nonlinear scalar conservation laws

ANNE-LAURE DALIBARD

Abstract

We study the limit as $\varepsilon \rightarrow 0$ of the entropy solutions of the equation $\partial_t u^\varepsilon + \operatorname{div}_x [A(\frac{x}{\varepsilon}, u^\varepsilon)] = 0$. We prove that the sequence u^ε two-scale converges towards a function $u(t, x, y)$, and u is the unique solution of a limit evolution problem. The remarkable point is that the limit problem is not a scalar conservation law, but rather a kinetic equation in which the macroscopic and microscopic variables are mixed. We also prove a strong convergence result in L^1_{loc} .

1. Introduction

This article is concerned with the asymptotic behavior of the sequence $u^\varepsilon \in \mathcal{C}([0, \infty), L^1_{\text{loc}}(\mathbb{R}^N))$, as the parameter ε vanishes, where u^ε is the entropy solution of the scalar conservation law

$$\frac{\partial u^\varepsilon(t, x)}{\partial t} + \sum_{i=1}^N \frac{\partial}{\partial x_i} A_i\left(\frac{x}{\varepsilon}, u^\varepsilon(t, x)\right) = 0 \quad t \geq 0, \quad x \in \mathbb{R}^N, \quad (1)$$

$$u^\varepsilon(t=0) = u_0\left(x, \frac{x}{\varepsilon}\right). \quad (2)$$

The functions $A_i = A_i(y, v)$ ($y \in \mathbb{R}^N$, $v \in \mathbb{R}$) are assumed to be Y -periodic, where $Y = \prod_{i=1}^N (0, T_i)$ is the unit cell, and u_0 is also assumed to be periodic in its second variable.

Under regularity hypotheses on the flux, namely $A \in W^{2, \infty}_{\text{per, loc}}(\mathbb{R}^{N+1})$, and when the initial data $u^\varepsilon(t=0)$ belongs to L^∞ , it is known that there exists a unique entropy solution u^ε of the above system for all $\varepsilon > 0$ given (see [4, 16, 17, 26, 27]). The study of the homogenization of such hyperbolic scalar conservation laws has been investigated by several authors, see for instance [9, 10, 11], and in the linear case [14, 15]. In dimension one, there is also an equivalence with Hamilton-Jacobi equations which allows to use the results of [18]. In general, the results obtained by these authors can be summarized as follows: there exists a function $u^0 = u^0(t, x, y)$ such that

$$u^\varepsilon - u^0\left(t, x, \frac{x}{\varepsilon}\right) \rightarrow 0 \quad \text{in } L^1_{\text{loc}}((0, \infty) \times \mathbb{R}^N). \quad (3)$$

The function $u^0(t, x, y)$ satisfies a microscopic equation, called cell problem, and an evolution equation, which is a scalar conservation law in which the coefficients depend on the microscopic variable y . In general, there is no “decoupling” of the macroscopic variables t, x , and the microscopic variable y : the average of u^0 with respect to the variable y is not the solution of an “average” conservation law.

To our knowledge, there are no results as soon as the dimension is strictly greater than one when the flux does not satisfy a structural condition of the type $A(y, \xi) = a(y)g(\xi)$. Here, we investigate the behavior of the family u^ε for arbitrary fluxes. We prove that (3) still holds, in a sense which will be made clear later on, and the function u^0 is a solution of a microscopic cell

problem. Precisely, we prove that even though there is no simple evolution equation satisfied by the function u^0 itself, the function

$$f(t, x, y, \xi) = \mathbf{1}_{\xi < u^0}$$

is the unique solution of a linear transport equation, with a source term which is a Lagrange multiplier accounting for the constraints on f . This statement is reminiscent of the kinetic formulation for scalar conservation laws (see [19, 20, 22], the general presentation in [23], and [8] for the heterogeneous case); this is not surprising since our method of proof relies on the kinetic formulation for equation (1). However, in general, it is unclear whether u^0 is the solution of a scalar conservation law. Thus the kinetic formulation appears as the ‘‘correct’’ vision of the entropy solutions of (1), at least as far as homogenization is concerned.

The rest of this introduction is devoted to the presentation of the main results. We begin with the description of the asymptotic problem, and then we state the convergence results.

1.1. Description of the asymptotic evolution problem

We first introduce the asymptotic evolution problem, for which we state an existence and uniqueness result; then we explain how this asymptotic problem can be understood formally.

In the following, we set, for $(y, \xi) \in \mathbb{R}^{N+1}$,

$$\begin{aligned} a_i(y, \xi) &= \frac{\partial A_i}{\partial \xi}(y, \xi), \quad 1 \leq i \leq N, \\ a_{N+1}(y, \xi) &= -\operatorname{div}_y A(y, \xi). \end{aligned}$$

We set $a(y, \xi) = (a_1(y, \xi), \dots, a_{N+1}(y, \xi)) \in \mathbb{R}^{N+1}$. Notice that $\operatorname{div}_{y, \xi} a(y, \xi) = 0$. These notations were introduced in [8]. Throughout the paper, we assume that $a \in W_{\text{per,loc}}^{1,\infty}(Y \times \mathbb{R})^{N+1}$.

Before giving the definition of the limit system, let us recall the kinetic formulation for equation (1), which may shed some light on the structure of the asymptotic evolution problem. Let u^ε be an entropy solution of (1). Then there exists a non-negative measure $m^\varepsilon \in M^1((0, \infty) \times \mathbb{R}^{N+1})$ such that $f^\varepsilon = \mathbf{1}_{\xi < u^\varepsilon(t, x)}$ is a solution of the transport equation

$$\begin{aligned} \partial_t f^\varepsilon + a_i \left(\frac{x}{\varepsilon}, \xi \right) \partial_{x_i} f^\varepsilon + \frac{1}{\varepsilon} a_{N+1} \left(\frac{x}{\varepsilon}, \xi \right) \partial_\xi f^\varepsilon &= \partial_\xi m^\varepsilon, \\ f^\varepsilon(t = 0, x, \xi) &= \mathbf{1}_{\xi < u_0(x, \frac{x}{\varepsilon})}. \end{aligned} \tag{4}$$

In fact, this equation was derived in [8] for the function $g^\varepsilon(t, x, \xi) = \chi(\xi, u^\varepsilon(t, x))$, where $\chi(\xi, u) = \mathbf{1}_{0 < \xi < u} - \mathbf{1}_{u < \xi < 0}$, for $u, \xi \in \mathbb{R}$, and under the additional assumption $a_{N+1}(y, 0) = 0$ for all $y \in \mathbb{R}^N$. However, it is easily proved, using the identity $f^\varepsilon = g^\varepsilon + \mathbf{1}_{\xi < 0}$, that f^ε satisfies (4), even when $a_{N+1}(y, 0)$ does not vanish.

We now define the limit system, which is reminiscent of equation (4) :

Definition 1 *Let $f \in L^\infty([0, \infty) \times \mathbb{R}^N \times Y \times \mathbb{R})$, $u_0 \in L^\infty(\mathbb{R}^N \times Y)$. We say that f is a generalized kinetic solution of the limit problem, with initial data $\mathbf{1}_{\xi < u_0}$, if there exists a distribution $\mathcal{M} \in \mathcal{D}'_{\text{per}}([0, \infty) \times \mathbb{R}^N \times Y \times \mathbb{R})$ such that f and \mathcal{M} satisfy the following properties:*

1. *Compact support in ξ : there exists a constant $M > 0$ such that*

$$\operatorname{Supp} \mathcal{M} \subset [0, \infty) \times \mathbb{R}^N \times Y \times [-M, M], \tag{6}$$

$$f(t, x, y, \xi) = 1 \quad \text{if } \xi < -M, \tag{7}$$

$$f(t, x, y, \xi) = 0 \quad \text{if } \xi > M. \tag{8}$$

2. *Microscopic equation for f : there exists a non-negative measure $m \in M^1((0, \infty) \times \mathbb{R}^N \times Y \times \mathbb{R})$ such that f is a solution in the sense of distributions of*

$$\operatorname{div}_{y, \xi}(a(y, \xi)f(t, x, y, \xi)) = \partial_\xi m, \tag{9}$$

and $\operatorname{Supp} m \subset [0, \infty) \times \mathbb{R}^N \times Y \times [-M, M]$.

3. *Evolution equation: the couple (f, \mathcal{M}) is a solution in the sense of distributions of*

$$\begin{cases} \partial_t f + \sum_{i=1}^N a_i(y, \xi) \partial_{x_i} f = \mathcal{M}, \\ f(t=0, x, y, \xi) = \mathbf{1}_{\xi < u_0(x, y)} =: f_0(x, y, \xi); \end{cases} \quad (10)$$

In other words, for any test function $\phi \in \mathcal{D}_{per}([0, \infty) \times \mathbb{R}^N \times Y \times \mathbb{R})$,

$$\begin{aligned} \int_0^\infty \int_{\mathbb{R}^N \times Y \times \mathbb{R}} f(t, x, y, \xi) \left\{ \partial_t \phi(t, x, y, \xi) + \sum_{i=1}^N a_i(y, \xi) \partial_{x_i} \phi(t, x, y, \xi) \right\} dt dx dy d\xi = \\ = - \langle \phi, \mathcal{M} \rangle_{\mathcal{D}, \mathcal{D}'} - \int_{\mathbb{R}^N \times Y \times \mathbb{R}} \mathbf{1}_{\xi < u_0(x, y)} \phi(t=0, x, y, \xi) dx dy d\xi. \end{aligned}$$

4. *Conditions on f :*

$$\partial_\xi f \leq 0 \quad \text{in } \mathcal{D}', \quad (11)$$

$$0 \leq f(t, x, y, \xi) \leq 1 \quad \text{almost everywhere.} \quad (12)$$

And for all compact set $K \subset \mathbb{R}^N$,

$$\frac{1}{\tau} \int_0^\tau \|f(s) - f_0\|_{L^2(K \times Y \times \mathbb{R})} ds \xrightarrow{\tau \rightarrow 0} 0. \quad (13)$$

5. *Condition on \mathcal{M} : define the set*

$$\begin{aligned} \mathcal{G} := \{ \psi \in L_{loc}^\infty(Y \times \mathbb{R}), \partial_\xi \psi \geq 0, \text{ and } \exists \mu \in M_{per}^1(Y \times \mathbb{R}), \exists C > 0, \exists \alpha_- \in \mathbb{R}, \\ \text{div}_{y, \xi}(\alpha \psi) = -\partial_\xi \mu, \text{ Supp } \mu \subset Y \times [-C, C], \mu \geq 0, \\ \psi(y, \xi) = \alpha_- \text{ if } \xi < -C \}. \end{aligned}$$

For $\varphi \in \mathcal{D}(\mathbb{R}_t \times \mathbb{R}_x^N)$ such that $\varphi \geq 0$ and $\varphi(t, x) = 0$ if $t \leq 0$, consider the function

$$\begin{aligned} \mathcal{M} *_{t, x} \varphi(t, x, y, \xi) &:= \int_0^\infty \int_{\mathbb{R}^N} \mathcal{M}(s, z, y, \xi) \varphi(t-s, x-z) ds dz \\ &= \int_0^\infty \int_{\mathbb{R}^N} f(s, z, y, \xi) \{ \partial_t \varphi(t-s, x-z) + a_i(y, \xi) \partial_{x_i} \varphi(t-s, x-z) \} ds dz. \end{aligned}$$

*Then $\mathcal{M} *_{t, x} \varphi$ belongs to $\mathcal{C}([0, \infty) \times \mathbb{R}^N, L^2(Y \times \mathbb{R}))$, and*

$$\forall (t, x) \in [0, \infty) \times \mathbb{R}^N, \quad \forall \psi \in \mathcal{G}, \quad \int_{Y \times \mathbb{R}} (\mathcal{M} *_{t, x} \varphi)(t, x, \cdot) \psi \leq 0. \quad (14)$$

We now state an existence and uniqueness result for solutions of the limit problem:

Theorem 1 *Let $A \in W_{per, loc}^{2, \infty}(Y \times \mathbb{R})$.*

1. *Existence: let $u_0 \in L_{loc}^1(\mathbb{R}^N; \mathcal{C}_{per}(Y)) \cap L^\infty(\mathbb{R}^N)$, and let $f_0(x, y, \xi) = \mathbf{1}_{\xi < u_0(x, y)}$, for $(x, y, \xi) \in \mathbb{R}^N \times Y \times \mathbb{R}$. Assume that there exists a non-negative measure $m_0 = m_0(x, y, \xi)$ such that f_0 is a solution of*

$$\sum_{i=1}^N \frac{\partial}{\partial y_i} (a_i(y, \xi) f_0) + \frac{\partial}{\partial \xi} (a_{N+1}(y, \xi) f_0) = \frac{\partial m_0}{\partial \xi} \quad (15)$$

and $\text{Supp } m_0 \subset \mathbb{R}^N \times Y \times [-M, M]$, where $M = \|u_0\|_\infty$.

Assume that there exist functions $u_1, u_2 \in L^\infty(Y)$ such that $\mathbf{1}_{\xi < u_i}$ is a solution of (15) for $i = 1, 2$, for some non-negative measures m_1, m_2 , and

$$u_1(y) \leq u_0(x, y) \leq u_2(y) \quad \text{for a.e. } x \in \mathbb{R}^N, y \in Y. \quad (16)$$

Then there exists a generalized kinetic solution f of the limit problem (in the sense of Definition 1), with initial data f_0 .

2. “Rigidity”: let $u_0 \in L^\infty(\mathbb{R}^N \times Y)$, and let $f \in L^\infty([0, \infty) \times \mathbb{R}^N \times Y \times \mathbb{R})$ be a generalized kinetic solution of the limit problem, with initial data $f_0 = \mathbf{1}_{\xi < u_0}$. Then there exists a function $u \in L^\infty([0, \infty) \times \mathbb{R}^N \times Y)$ such that

$$f(t, x, y, \xi) = \mathbf{1}_{\xi < u(t, x, y)} \quad \text{almost everywhere.}$$

3. Uniqueness and contraction principle: let $u_0, v_0 \in L^\infty(\mathbb{R}^N \times Y)$, and let f, g be two generalized kinetic solutions of the limit problem with initial data $\mathbf{1}_{\xi < u_0}$ and $\mathbf{1}_{\xi < v_0}$ respectively. Then there exists a constant $C > 0$ such that for all $t > 0$, for all $R, R' > 0$,

$$\|f(t) - g(t)\|_{L^1(B_R \times Y \times \mathbb{R})} \leq e^{Ct+R} \left(\|u_0 - v_0\|_{L^1(B_{R'} \times Y)} + e^{-R'} \right). \quad (17)$$

As a consequence, for all $u_0 \in L^\infty(\mathbb{R}^N \times Y) \cap L^1_{loc}(\mathbb{R}^N, \mathcal{C}_{per}(Y))$ satisfying (15) and (16), there exists a unique generalized kinetic solution $f \in L^\infty([0, \infty) \times \mathbb{R}^N \times Y \times \mathbb{R})$ of the limit problem.

Remark 1 Notice that for any function $v \in L^\infty(Y)$, v is an entropy solution of the cell problem

$$\operatorname{div}_y A(y, v(y)) = 0 \quad (18)$$

if and only if there exists a non-negative measure $m \in M^1_{per}(Y \times \mathbb{R})$ such that equation

$$\operatorname{div}_{y, \xi}(a(y, \xi) \mathbf{1}_{\xi < v(y)}) = \partial_\xi m$$

is satisfied in the sense of distributions on $Y \times \mathbb{R}$. Hence equation (15) entails that u_0 is an entropy solution of the cell equation (18). In that case, it is said that the initial data u_0 is “well-prepared”, meaning that the microscopic profile of u_0 is adapted to the microstructure dictated by the equation.

In the case where A is divergence-free, condition (15) becomes

$$\sum_{i=1}^N \partial_{y_i}(a_i(y, \xi) \mathbf{1}_{\xi < v(y)}) = 0.$$

Indeed, in that case, v satisfies

$$\sum_{i=1}^N \partial_{y_i}(a_i(y, \xi) \mathbf{1}_{\xi < v(y)}) = \partial_\xi m$$

for some non-negative measure m such that $\operatorname{Supp} m \subset Y \times [-M, M]$. Consequently, for $\xi \geq -M$, we have

$$\sum_{i=1}^N \partial_{y_i} \left(\int_{-M-1}^{\xi} a_i(y, w) \mathbf{1}_{w < v(y)} dw \right) = m(y, \xi) \geq 0.$$

Since the left-hand side has zero mean-value on Y for all $\xi \in [-M, M]$, we deduce that $m = 0$. Thus, in the case where the flux A is divergence free, the limit system takes a slightly simpler form: conditions (9), (14) become

$$\begin{aligned} \operatorname{div}_y(a(y, \xi) f(t, x, y, \xi)) &= 0, \\ \partial_t f + \sum_{i=1}^N a_i(y, \xi) \partial_{x_i} f &= \mathcal{M}, \\ \left\{ \begin{array}{l} \int_{Y \times \mathbb{R}} (\mathcal{M} *_{t, x} \varphi)(t, x, \cdot) \psi \leq 0, \\ \forall \psi \in L^\infty_{loc}(Y \times \mathbb{R}), \operatorname{div}_y(a\psi) = 0, \text{ and } \partial_\xi \psi \geq 0. \end{array} \right. & \quad (19) \end{aligned}$$

All the other properties remain the same.

Remark 2 Assume that the flux A is divergence-free, and set

$$C_1 := \{\psi \in L^2_{loc}(Y \times \mathbb{R}), \sum_{i=1}^N \frac{\partial}{\partial y_i} (a_i(y, \xi) \psi(y, \xi)) = 0\},$$

$$C_2 := \{\psi \in L^\infty_{loc}(Y \times \mathbb{R}), \partial_\xi \psi \geq 0\}.$$

Then C_1, C_2 are convex sets of the vector space $L^2_{loc}(Y \times \mathbb{R})$. Thus condition (14) can be written as follows: for all $\varphi \in \mathcal{D}((-\infty, 0) \times \mathbb{R}^N)$ such that $\varphi \geq 0$, for all $(t, x) \in (0, \infty) \times \mathbb{R}^N$, we have

$$\mathcal{M} * \varphi(t, x) \in (C_1 \cap C_2)^\circ,$$

where C° denotes the normal cone of C . Let us recall that when the space dimension is finite (that is, if C_1, C_2 are convex cones in \mathbb{R}^d for some $d \in \mathbb{N}$), then

$$(\text{cl}(C_1) \cap \text{cl}(C_2))^\circ = \text{cl}(C_1^\circ + C_2^\circ),$$

where $\text{cl}(A)$ denotes the closure of the set A .

If we forget about the closure and the fact that we are considering convex sets in an infinite dimensional space, then we are tempted to write

$$\mathcal{M} * \varphi(t, x) \in (C_1 \cap C_2)^\circ = \mu_1 + \mu_2,$$

with $\mu_i \in C_i^\circ$, $i = 1, 2$. Moreover, very formally, we have

$$C_2^\circ = \{\partial_\xi m, m \text{ non-negative measure}\}.$$

Thus, we may think of \mathcal{M} as some distribution of the form

$$\mathcal{M} = \partial_\xi m + \mu_1,$$

with m a non-negative measure on $[0, \infty) \times \mathbb{R}^N \times Y \times \mathbb{R}$, and $\mu_1 \in C_1^\circ$.

Of course, these computations are not rigorous, but we believe they may help the reader to understand the action of the distribution \mathcal{M} (at least in the divergence-free case), even though the precise structure of \mathcal{M} shall not be needed in the proof. Inequality (14) is sufficient for all the applications in this paper.

Let us stress that uniqueness for the limit problem holds, even though the cell problem does not have a unique solution in general; indeed, in the linear divergence free case, that is, if $A(y, \xi) = b(y)\xi$, with $\text{div}_y b = 0$, then a function u is a solution of the cell problem if

$$\text{div}_y(b(y)u(y)) = 0, \quad \langle u \rangle_Y = 0.$$

The constant function equal to zero is a solution of this equation, but in general there are other entropy solutions: for instance, let us consider the case where $N = 2$, and

$$b(y_1, y_2) = (-\partial_2 \phi(y_1, y_2), \partial_1 \phi(y_1, y_2)),$$

for some function $\phi \in \mathcal{C}^2_{\text{per}}(Y)$. Then any function u of the form $g(\phi) - \langle g(\phi) \rangle$, with g a continuous function, is an entropy solution. Let us emphasize that nonlinearity assumptions on the flux are not enough to ensure uniqueness of solutions either, see for instance [18].

In Theorem 1, uniqueness of solutions of the limit system follows from a contraction principle associated with the macroscopic evolution equation, rather than the microscopic cell equation. The well-preparedness of the initial data, that is, the fact that $u_0(x, \cdot)$ is an entropy solution of the cell problem, is fundamental.

On the other hand, the lack of uniqueness of solutions of the cell problem entails that in general, there is no notion of homogenized problem. Indeed, if u is a solution of

$$\text{div}_y A(y, p + u(y)) = 0, \quad \langle u \rangle_Y = 0,$$

then in general, the quantity

$$\langle A(\cdot, p + u(\cdot)) \rangle$$

depends on u (except when $N = 1$, and in some special cases, when $N = 2$; see [13,18]). Hence the macroscopic and microscopic scales cannot be decoupled: if $\mathbf{1}_{\xi < u(t,x,y)}$ is a solution of the limit evolution problem, then $\bar{u}(t,x) = \langle u(t,x,\cdot) \rangle$ does not satisfy any remarkable equation. This is the main consequence of the absence of uniqueness for the cell problem.

Let us mention an important particular case of Theorem 1, which we call the ‘‘separate case’’. We now assume that the flux A can be written $A(y,\xi) = a_0(y)g(\xi)$, with $\operatorname{div}_y a_0 = 0$. This case has already been thoroughly investigated by Weinan E in [9] in the case where $g'(\xi) \neq 0$ for all ξ , that is, when the function g is strictly monotonous. Here, we prove that his results hold with no restriction on g .

Let us introduce the so-called ‘‘constraint space’’

$$\mathbb{K}_0 := \{f \in L^1(Y); \operatorname{div}_y(a_0 f) = 0 \text{ in } \mathcal{D}'\},$$

and the orthogonal projection P_0 on $\mathbb{K}_0 \cap L^2(Y)$ for the scalar product in $L^2(Y)$.

Then the following properties hold: for all $f, g \in L^2(Y)$, if $f \in \mathbb{K}_0$, then

$$P_0(fg) = fP_0(g).$$

And if $f, g \in \mathbb{K}_0 \cap L^2(Y)$, then the product fg belongs to \mathbb{K}_0 . Notice also that all functions which do not depend on y belong to \mathbb{K}_0 , and that $L^\infty(Y)$ is stable by P_0 .

Proposition 1 *Let $u_0 \in L^1(\mathbb{R}^N, \mathcal{C}_{per}(Y)) \cap L^\infty(\mathbb{R}^N \times Y)$ such that $u_0(x, \cdot) \in \mathbb{K}_0$ for a.e. $x \in \mathbb{R}^N$.*

For $1 \leq i \leq N$, define the vector valued function $\tilde{a}_0 \in L^\infty(Y)^N$ by $\tilde{a}_{0,i} = P_0(a_{0,i})$. Let $u = u(t, x; y)$ be the entropy solution of the scalar conservation law

$$\begin{cases} \partial_t u(t, x; y) + \operatorname{div}_x(\tilde{a}_0(y)g(u(t, x; y))) = 0, & t > 0, x \in \mathbb{R}^N, y \in Y, \\ u(t = 0, x; y) = u_0(x, y). \end{cases} \quad (20)$$

Then the function $f(t, x, y, \xi) = \mathbf{1}_{\xi < u(t,x,y)}$ is the unique generalized kinetic solution of the limit problem (10) with initial data $\mathbf{1}_{\xi < u_0(x,y)}$. In that case the distribution \mathcal{M} is given by

$$\mathcal{M} = \frac{\partial m}{\partial \xi} + g'(\xi)(\tilde{a}_0(y) - a_0(y)) \cdot \nabla_x f,$$

where m is the kinetic entropy defect measure associated with the function u , that is, f is a solution of

$$\partial_t f + g'(\xi)\tilde{a}_0(y) \cdot \nabla_x f = \partial_\xi m.$$

As a consequence, the solution $u(t, x; y)$ of (20) is an entropy solution of

$$\operatorname{div}_y A(y, u) = 0$$

for almost every $(t, x) \in (0, \infty) \times \mathbb{R}^N$.

1.2. Convergence results

Our first result is concerned with entropy solutions of (1).

Theorem 2 *Let $A \in W_{per,loc}^{2,\infty}(\mathbb{R}^{N+1})$. Assume that the initial data $u_0 \in L_{loc}^1(\mathbb{R}^N, \mathcal{C}_{per}(Y))$ satisfies (15), (16). Let $f = \mathbf{1}_{\xi < u}$ be the unique generalized kinetic solution of the limit problem, with initial data $\mathbf{1}_{\xi < u_0}$; the existence of f follows from Theorem 1. Then as ε vanishes,*

$$\mathbf{1}_{\xi < u^\varepsilon(t,x)} \stackrel{2}{\xrightarrow{sc.}} \mathbf{1}_{\xi < u(t,x,y)}. \quad (21)$$

As a consequence, for all regularization kernels φ^δ of the form

$$\varphi^\delta(x) = \frac{1}{\delta^N} \varphi\left(\frac{x}{\delta}\right), \quad x \in \mathbb{R}^N,$$

with $\varphi \in \mathcal{D}(\mathbb{R}^N)$, $\int \varphi = 1$, $0 \leq \varphi \leq 1$, we have, for all compact set $K \subset [0, \infty) \times \mathbb{R}^N$,

$$\lim_{\delta \rightarrow 0} \lim_{\varepsilon \rightarrow 0} \left\| u^\varepsilon(t, x) - u *_x \varphi^\delta\left(t, x, \frac{x}{\varepsilon}\right) \right\|_{L^1(K)} = 0. \quad (22)$$

Remark 3 Assumption (15) means that u_0 is “well-prepared”, that is, $u_0(x, \cdot)$ is an entropy solution of

$$\operatorname{div}_y (A(y, u_0(x, y))) = 0$$

for a.e. $x \in \mathbb{R}^N$. If this hypothesis is not satisfied, then it is expected that the behavior of the sequence u^ε will depend on the nature of the flux. If the flux is linear, then oscillations will propagate, and the cell equation (9) shall not be satisfied in general. If the flux satisfies some strong nonlinearity assumption, on the contrary, the conjecture is that the solution u^ε re-prepares itself in order to match the microscopic profile dictated by the equation. Few results in this direction are known in the hyperbolic case; the reader may consult for instance [2, 10, 12, 25]. In [6], the author studies the same equation as (1) in which a viscosity term of order ε is added, and proves such a result, but the method relies strongly on the parabolicity of the equation.

Remark 4 The way in which Theorem 2 is stated might seem slightly peculiar; indeed, convergence results of the type

$$u^\varepsilon(t, x) - u\left(t, x, \frac{x}{\varepsilon}\right) \rightarrow 0 \quad \text{in } L^1_{loc}$$

are expected to hold. In order to establish such a result, it seems necessary to prove that

$$\lim_{\delta \rightarrow 0} \int_K \sup_{y \in Y} |u(t, x, y) - u *_{x, \varepsilon} \varphi^\delta(t, x, y)| \, dt \, dx = 0.$$

But the evolution equation for u (or rather, for $\mathbf{1}_{\xi < u}$) is given by Definition 1; since the distribution \mathcal{M} hinders most computations, it seems difficult to derive such estimates.

The next result generalizes Theorem 1 to weaker solutions of equation (1), called kinetic solutions. In order to simplify the presentation, let us restrict the statement to the divergence-free case; it is explained in the remark following the Theorem how to derive an analogous result when the flux A is arbitrary.

For the reader’s convenience, we first recall the definition of kinetic solutions in the divergence-free case (see [8] for the heterogeneous case, and the presentation in [23] for the homogeneous case). For $(u, v) \in \mathbb{R}^2$, set

$$\chi(u, v) = \begin{cases} 1 & \text{if } 0 < u < v, \\ -1 & \text{if } v < u < 0, \\ 0 & \text{otherwise.} \end{cases}$$

Definition 2 (Kinetic solutions of (1)) Let $u^\varepsilon \in \mathcal{C}([0, \infty), L^1(\mathbb{R}^N))$. Assume that there exists a non-negative measure $m^\varepsilon \in \mathcal{C}(\mathbb{R}_\xi, M_w^1([0, \infty) \times \mathbb{R}^N))$ such that for all $T > 0$, the function

$$\xi \mapsto \int_0^T \int_{\mathbb{R}^N} m^\varepsilon(t, x, \xi) \, dt \, dx$$

is bounded on \mathbb{R} , and vanishes as $|\xi| \rightarrow \infty$.

Assume also that $f^\varepsilon(t, x, \xi) := \chi(\xi, u^\varepsilon(t, x))$ is a solution in the sense of distributions of the linear transport equation

$$\frac{\partial f^\varepsilon}{\partial t} + \sum_{i=1}^N a_i\left(\frac{x}{\varepsilon}, \xi\right) \partial_{x_i} f^\varepsilon = \frac{\partial m^\varepsilon}{\partial \xi} \quad t \geq 0, \quad x \in \mathbb{R}^N, \quad (23)$$

$$f^\varepsilon(t=0) = \chi\left(\xi, u_0\left(x, \frac{x}{\varepsilon}\right)\right), \quad (24)$$

Then it is said that u^ε is a kinetic solution of equation (1).

Remark 5 Let us recall the Definition of the space $\mathcal{C}(\mathbb{R}_\xi, M_w^1([0, \infty) \times \mathbb{R}^N))$. Let $m \in M^1([0, \infty) \times \mathbb{R}^N)$; for $\theta \in \mathcal{C}_c([0, \infty) \times \mathbb{R}^N)$, define $\mu_\theta \in M^1(\mathbb{R})$ by

$$\mu_\theta := \int_0^\infty \int_{\mathbb{R}^N} m(t, x, \cdot) \theta(t, x) \, dt \, dx.$$

Then

$$\mathcal{C}(\mathbb{R}_\xi, M_w^1([0, \infty) \times \mathbb{R}^N)) := \{m \in M^1([0, \infty) \times \mathbb{R}^N); \forall \theta \in \mathcal{C}_c([0, \infty) \times \mathbb{R}^N), \mu_\theta \in \mathcal{C}(\mathbb{R})\}.$$

The existence of kinetic solutions of (1) is only known when the flux satisfies additional regularity assumptions. Assume that $a_i \in \mathcal{C}_{\text{per}}^1(Y \times \mathbb{R})$ for $1 \leq i \leq N$, and that there exists a constant C such that

$$|a(y, \xi)| \leq C(1 + |\xi|) \quad \forall y \in Y \quad \forall \xi \in \mathbb{R}. \quad (25)$$

Under such hypotheses, it is proved in [8] that for all $u_0 \in L^1(\mathbb{R}^N, \mathcal{C}_{\text{per}}(Y))$, there exists a unique function $u^\varepsilon \in \mathcal{C}([0, \infty), L^1(\mathbb{R}^N))$ such that $\chi(\xi, u^\varepsilon)$ is a solution of (1); u^ε is called the kinetic solution of (1)-(2). And if u^ε is bounded in $L^\infty((0, T) \times \mathbb{R}^N)$ for all $T > 0$, then u^ε is the entropy solution of (1). Moreover, a contraction principle holds between kinetic solutions.

Let us now state the convergence result for kinetic solutions :

Theorem 3 *Let $A \in W_{\text{per}, \text{loc}}^{2, \infty}(Y \times \mathbb{R})$ such that $\text{div}_y A(y, \xi) = 0$ for all y, ξ . Assume that $a_i \in \mathcal{C}_{\text{per}}^1(Y \times \mathbb{R})$ for $1 \leq i \leq N$, and that (25) is satisfied. Assume that the initial data u_0 belongs to $L^1(\mathbb{R}^N, \mathcal{C}_{\text{per}}^1(Y))$ and satisfies*

$$\sum_{i=1}^N \frac{\partial}{\partial y_i} (a_i(y, \xi) \chi(\xi, u_0)) = 0.$$

Let $u^\varepsilon \in \mathcal{C}([0, \infty), L^1(\mathbb{R}^N))$ be the kinetic solution of (1) with initial data $u_0(x, x/\varepsilon)$. Then there exists a function $u \in L^\infty([0, \infty), L^1(\mathbb{R}^N \times Y))$ such that the convergence results (21) and (22) hold, and

$$\frac{\partial}{\partial y_i} (a(y, \xi) \chi(\xi, u(t, x, y))) = 0 \quad \text{in } \mathcal{D}'.$$

Moreover, if we set

$$\mathcal{M} := \frac{\partial}{\partial t} \chi(\xi, u) + \sum_{i=1}^N a_i(y, \xi) \frac{\partial}{\partial x_i} \chi(\xi, u) \in \mathcal{D}',$$

then \mathcal{M} satisfies (19).

Remark 6 *Let us explain how this result can be generalized to the general case. First, the L^1 setting is not adapted to this case, because the L^1 norm is not conserved by the equation in general. Hence another notion of kinetic solutions is needed; the correct functional space should be of the type $V + L^1(\mathbb{R}^N)$, where V is a fixed solution of the cell problem.*

Then, the crucial point in Theorem 3 is to find a sequence u_0^n such that u_0^n converges towards u_0 in $L^1(\mathbb{R}^N, \mathcal{C}_{\text{per}}(Y))$, and for all $n \in \mathbb{N}$, u_0^n satisfies (15), (16). Finding such a sequence is easy in the divergence-free case, but seems more difficult in the general case, since solutions of the cell problem are not known. This seems to be the main obstacle to the generalization of Theorem 3 to arbitrary fluxes. If this step is admitted, it is likely that the proof of Theorem 3 can be adapted to general settings.

The organization of the paper is the following: in section 2 we prove, under the hypotheses of Theorem 2, that the two-scale limit of the sequence $\mathbf{1}_{\xi < u^\varepsilon(t, x)}$ is a generalized kinetic solution of the limit system. In section 3, we study the limit problem introduced in Definition 1 and we prove the rigidity and uniqueness results in Theorem 1; hence Theorem 1 and 2 will be proved by the end of section 3. In section 4, we study a relaxation model of BGK type, approaching the limit system in the divergence free case. In section 5, we prove Proposition 1. Eventually, in section 6, we have gathered further remarks on the notion of limit evolution problem.

2. Asymptotic behavior of the sequence u^ε

In this section, we prove that the two-scale limit of the sequence $f^\varepsilon = \mathbf{1}_{\xi < u^\varepsilon(t, x)}$, say $f^0(t, x, y, \xi)$, is a generalized kinetic solution of the limit system; thus the existence result of Theorem 1 follows from this section. The organization is the following: we first derive some basic (microscopic) properties for the function f^0 . Then we explain how regularization by convolution can be used in two-scale problems. The two other subsections are devoted to the other properties of the limit system, namely condition (14) and the strong continuity at time $t = 0$.

2.1. Basic properties of f^0

We use the concept of two-scale convergence, formalized by G. Allaire after an idea of G. N'Guetseng (see [1, 21]). The fundamental result of [1] can be generalized to the present setting as follows:

Corollary 1 *Let $(g^\varepsilon)_{\varepsilon>0}$ be a bounded sequence in $L^\infty((0, \infty) \times \mathbb{R}^{N+1})$. Then there exists a function $g^0 \in L^\infty((0, \infty) \times \mathbb{R}^N \times Y \times \mathbb{R})$, and a subsequence (ε_n) such that $\varepsilon_n \rightarrow 0$ as $n \rightarrow \infty$, such that*

$$\int_0^\infty \int_{\mathbb{R}^{N+1}} g^{\varepsilon_n}(t, x, \xi) \psi\left(t, x, \frac{x}{\varepsilon_n}, \xi\right) dt dx d\xi \rightarrow \int_0^\infty \int_{\mathbb{R}^N \times Y \times \mathbb{R}} g^0(t, x, y, \xi) \psi(t, x, y, \xi) dt dx dy d\xi$$

for all functions $\psi \in L^1((0, \infty) \times \mathbb{R}^{N+1}; \mathcal{C}_{per}(Y))$.

It is said that the sequence $(g^{\varepsilon_n})_{n \in \mathbb{N}}$ two-scale converges towards g^0 .

Here, the sequence f^ε is bounded by 1 in L^∞ ; hence we can extract a subsequence, still denoted by ε , and find a function $f^0 \in L^\infty((0, \infty) \times \mathbb{R}^N \times Y \times \mathbb{R})$ such that (f^ε) two-scale converges towards f^0 . It is easily checked that f^0 inherits the following properties from the sequence f^ε

$$0 \leq f^0(t, x, y, \xi) \leq 1, \quad (26)$$

$$\partial_\xi f^0 \leq 0. \quad (27)$$

Now, let us prove (7)-(8): let

$$M := \max(\|u_1\|_\infty, \|u_2\|_\infty),$$

where u_1, u_2 are the functions appearing in assumption (16). Since $u_i(x/\varepsilon)$ is a stationary solution of (1), by a comparison principle for equation (1), we deduce that

$$u_1\left(\frac{x}{\varepsilon}\right) \leq u^\varepsilon(t, x) \leq u_2\left(\frac{x}{\varepsilon}\right) \quad \text{for almost every } t > 0, x \in \mathbb{R}^N.$$

Thus $\|u^\varepsilon\|_{L^\infty((0, \infty) \times \mathbb{R}^N)} \leq M$, and for almost every t, x, ξ , for all $\varepsilon > 0$,

$$\begin{aligned} f^\varepsilon(t, x, \xi) &= 1 & \text{if } \xi < -M, \\ f^\varepsilon(t, x, \xi) &= 0 & \text{if } \xi > M. \end{aligned}$$

Passing to the two-scale limit, we infer (7) and (8).

Now, we derive a microscopic equation for f^0 . First, multiplying (4) by $S'(\xi)$, with $S' \in \mathcal{D}(\mathbb{R})$, and integrating on $(0, T) \times B_R \times \mathbb{R}$, with $T > 0, R > 0$, yields

$$\begin{aligned} & \int_{B_R} \left(S(u^\varepsilon(T, x)) - S\left(u_0\left(x, \frac{x}{\varepsilon}\right)\right) \right) dx + \int_0^T \int_{\mathbb{R}} \int_{\partial B_R} a\left(\frac{x}{\varepsilon}, \xi\right) \cdot n_R(x) f^\varepsilon S'(\xi) d\sigma_R(x) d\xi dt - \\ & - \frac{1}{\varepsilon} \int_0^T \int_{\mathbb{R}} \int_{B_R} a_{N+1}\left(\frac{x}{\varepsilon}, \xi\right) f^\varepsilon S''(\xi) dx d\xi dt = - \int_0^T \int_{\mathbb{R}} \int_{B_R} m^\varepsilon(t, x, \xi) S''(\xi) dx d\xi dt, \end{aligned}$$

where $n_R(x)$ is the outward-pointing normal to B_R at a given point $x \in \partial B_R$, and $d\sigma_R(x)$ is the Lebesgue measure on ∂B_R .

Hence we obtain the following bound on m^ε

$$\varepsilon m^\varepsilon((0, T) \times B_R \times \mathbb{R}) \leq C_{T,R}$$

for all $\varepsilon > 0, R > 0, T > 0$, and $\text{Supp } m^\varepsilon \subset (0, \infty) \times \mathbb{R} \times [-M, M]$.

Consequently, there exists a further subsequence, still denoted by ε , and a non-negative measure $m^0 = m^0(t, x, y, \xi)$ such that $\varepsilon m^\varepsilon$ two-scale converges to m^0 (the concept of two-scale convergence can easily be generalized to measures; the arguments are the same as in [1], the only difference lies in the functional spaces). Moreover, $\text{Supp } m^0 \subset (0, \infty) \times \mathbb{R} \times Y \times [-M, M]$.

We now multiply (4) by test functions of the type $\varepsilon \varphi(t, x, x/\varepsilon, \xi)$, with $\varphi \in \mathcal{D}_{per}([0, \infty) \times \mathbb{R}^N \times Y \times \mathbb{R})$, and we pass to the two-scale limit. We obtain, in the sense of distributions on $(0, \infty) \times \mathbb{R}^N \times Y \times \mathbb{R}$

$$\frac{\partial}{\partial y_i} (a_i(y, \xi) f^0) + \frac{\partial}{\partial \xi} (a_{N+1}(y, \xi) f^0) = \frac{\partial m^0}{\partial \xi}. \quad (28)$$

Thus (9) is satisfied, which completes the derivation of the basic properties of f^0 .

Now, we define the distribution

$$\mathcal{M} := \frac{\partial f^0}{\partial t} + \sum_{i=1}^N a_i(y, \xi) \frac{\partial f^0}{\partial x_i}.$$

The distribution \mathcal{M} obviously satisfies (6). The next step is to prove that \mathcal{M} satisfies (14); since regularizations by convolution are involved in condition (14), we now describe the links between convolution and two-scale convergence.

2.2. Regularization by convolution and two-scale convergence

In this subsection, we wish to make a few remarks concerning the links between convolution and two-scale convergence. Indeed, it is a well-known fact that if a sequence (f_n) weakly converges in $L^2(\mathbb{R}^N)$ towards a function f , then for all convolution kernels $\varphi = \varphi(x)$, the sequence $(f_n * \varphi)$ two-scale converges in L^2 towards $f * \varphi$. It would be convenient to have a similar property for two-scale limits. However, in general, if a sequence $f^\varepsilon = f^\varepsilon(x)$ is bounded in $L^2(\mathbb{R}^N)$ and two-scale converges towards a function $f = f(x, y) \in L^2(\mathbb{R}^N \times Y)$, then $f^\varepsilon * \varphi$ does not two-scale converge towards $f *_{x} \varphi$. Indeed, if $\psi = \psi(x, y) \in L^2(\mathbb{R}^N, \mathcal{C}_{\text{per}}(Y))$, then

$$\begin{aligned} & \int_{\mathbb{R}^N} f^\varepsilon * \varphi(x) \psi\left(x, \frac{x}{\varepsilon}\right) dx \\ &= \int_{\mathbb{R}^{2N}} f^\varepsilon(x') \varphi(x - x') \psi\left(x, \frac{x}{\varepsilon}\right) dx dx' \\ &= \int_{\mathbb{R}^N} dx' f^\varepsilon(x') \left[\int_{\mathbb{R}^N} \varphi(x - x') \psi\left(x, \frac{x}{\varepsilon}\right) dx \right]. \end{aligned}$$

In general, the quantity between brackets in the last integral cannot be written as a function of x' and x'/ε , and it seems difficult to pass to the limit as $\varepsilon \rightarrow 0$.

In order to get rid of this difficulty, let us suggest the following construction, which is reminiscent of the doubling of variables in the papers of Kruřkov, see [16, 17]. With the same notations as above, consider the test function $(\psi *_{x} \check{\varphi})(x, x/\varepsilon)$, where $\check{\varphi}(x) := \varphi(-x) \forall x \in \mathbb{R}^N$. Then by definition of the two-scale convergence,

$$\int_{\mathbb{R}^N} f^\varepsilon(x) [\psi *_{x} \check{\varphi}]\left(x, \frac{x}{\varepsilon}\right) dx \rightarrow \int_{\mathbb{R}^N \times Y} f(x, y) [\psi *_{x} \check{\varphi}](x, y) dx dy$$

And

$$\begin{aligned} \int_{\mathbb{R}^N} f^\varepsilon(x) [\psi *_{x} \check{\varphi}]\left(x, \frac{x}{\varepsilon}\right) dx &= \int_{\mathbb{R}^{2N}} f^\varepsilon(x') \varphi(x - x') \psi\left(x, \frac{x'}{\varepsilon}\right) dx dx', \\ \int_{\mathbb{R}^N \times Y} f(x, y) [\psi *_{x} \check{\varphi}](x, y) dx dy &= \int_{\mathbb{R}^N \times Y} [f *_{x} \varphi](x, y) \psi(x, y) dx dy. \end{aligned}$$

Consequently, as $\varepsilon \rightarrow 0$,

$$\int_{\mathbb{R}^{2N}} f^\varepsilon(x') \varphi(x - x') \psi\left(x, \frac{x'}{\varepsilon}\right) dx dx' \rightarrow \int_{\mathbb{R}^N \times Y} [f *_{x} \varphi](x, y) \psi(x, y) dx dy \quad (29)$$

for all $\varphi \in \mathcal{D}(\mathbb{R}^N)$, for all $\psi \in L^2(\mathbb{R}^N, \mathcal{C}_{\text{per}}(Y))$.

In fact, different assumptions on the function ψ can be chosen; the key point is that ψ should be an admissible test function in the sense of Allaire (see [1]). In particular, if there exist functions $\psi_1 \in \mathcal{D}(\mathbb{R}^N)$, $\psi_2 \in L^\infty(Y)$ such that

$$\psi(x, y) = \psi_1(x) \psi_2(y),$$

then ψ is an admissible test function, and (29) holds.

2.3. Proof of the condition on \mathcal{M}

The goal of this subsection is to prove that with

$$\mathcal{M} = \partial_t f^0 + \sum_{i=1}^N a_i(y, \xi) \partial_i f^0,$$

condition (14) holds; hence, let $\varphi \in \mathcal{D}(\mathbb{R} \times \mathbb{R}^N)$, $\theta \in \mathcal{D}([0, \infty) \times \mathbb{R}^N)$, such that

$$\begin{aligned} \varphi &\geq 0, \quad \theta \geq 0, \\ \varphi(t, x) &= 0 \quad \forall t \geq 0, \quad \forall x \in \mathbb{R}^N. \end{aligned}$$

The function φ shall be used as a convolution kernel, which explains the hypothesis on its support. We do not assume that $\theta(t=0) = 0$.

Let $\psi \in \mathcal{G}$ arbitrary (the set \mathcal{G} was introduced in Definition 1), and let C_0 be a compact set in \mathbb{R} such that $[-M, M] \subset C_0$, where M is the constant in properties (6)-(8). We have to prove that the quantity

$$\begin{aligned} A := & \int_0^\infty \int_0^\infty \int_{\mathbb{R}^{2N} \times Y \times C_0} f^0(s, z, y, \xi) \left\{ \partial_t \varphi(t-s, x-z) + \sum_{i=1}^N a_i(y, \xi) \partial_i \varphi(t-s, x-z) \right\} \times \\ & \times \psi(y, \xi) \theta(t, x) \, d\xi \, dy \, dx \, dz \, ds \, dt \end{aligned}$$

is non-positive. Notice that properties (6)-(8) entail that the quantity A is independent of the set C_0 .

Before going into the technicalities, let us explain formally why the property is true; let us forget about the convolution and the regularity issues, and take the test function

$$\theta(t, x) \psi\left(\frac{x}{\varepsilon}, \xi\right)$$

in equation (4).

Let $R > \max(M, C+1)$; recall that M and C are such that

$$\begin{aligned} f_0(t, x, y, \xi) &= 0 \quad \text{if } \xi > M, \\ f_0(t, x, y, \xi) &= 1 \quad \text{if } \xi < -M, \end{aligned}$$

and $\psi(y, \xi) = \alpha_-$ if $\xi < -C$. Integrating on $[0, \infty) \times \mathbb{R}^N \times [-R, R]$, we obtain

$$\begin{aligned} & \int_0^\infty \int_{\mathbb{R}^N} \int_{-R}^R f^\varepsilon(t, x, \xi) \left[\partial_t \theta(t, x) + a_i\left(\frac{x}{\varepsilon}, \xi\right) \partial_{x_i} \theta(t, x) \right] \psi\left(\frac{x}{\varepsilon}, \xi\right) \, dx \, d\xi \, dt \\ & - \frac{1}{\varepsilon} \int_0^\infty \int_{\mathbb{R}^N} \int_{-R}^R f^\varepsilon(t, x, \xi) \frac{\partial \mu}{\partial \xi}\left(\frac{x}{\varepsilon}, \xi\right) \theta(t, x) \, dx \, d\xi \, dt \\ & + \alpha_- \int_0^\infty \int_{\mathbb{R}^N} \frac{1}{\varepsilon} a_{N+1}\left(\frac{x}{\varepsilon}, -R\right) \theta(t, x) \, dt \, dx \\ = & \int_0^\infty \int_{\mathbb{R}^N} \int_{-R}^R m^\varepsilon(s, z, \xi) \partial_\xi \psi\left(\frac{x}{\varepsilon}, \xi\right) \, dz \, d\xi \, ds - \int_{\mathbb{R}^N} \int_{-R}^R \mathbf{1}_{\xi < u_0(x, \frac{x}{\varepsilon})} \theta(t=0, x) \psi\left(\frac{x}{\varepsilon}, \xi\right) \, dx \, d\xi. \end{aligned}$$

Notice that

$$\frac{1}{\varepsilon} a_{N+1}\left(\frac{x}{\varepsilon}, -R\right) = -\operatorname{div}_x A\left(\frac{x}{\varepsilon}, -R\right),$$

and thus

$$\begin{aligned}
& \int_0^\infty \int_{\mathbb{R}^N} \int_{-R}^R f^\varepsilon(t, x, \xi) \left[\partial_t \theta(t, x) + a_i \left(\frac{x}{\varepsilon}, \xi \right) \partial_{x_i} \theta(t, x) \right] \psi \left(\frac{x}{\varepsilon}, \xi \right) dx d\xi dt \\
&= \int_0^\infty \int_{\mathbb{R}^N} \int_{-R}^R \left[m^\varepsilon(s, z, \xi) \partial_\xi \psi \left(\frac{x}{\varepsilon}, \xi \right) - \frac{1}{\varepsilon} \mu \left(\frac{x}{\varepsilon}, \xi \right) \partial_\xi f^\varepsilon(t, x, \xi) \right] \theta(t, x) dz d\xi ds \\
&\quad - \alpha_- \int_0^\infty \int_{\mathbb{R}^N} A_i \left(\frac{x}{\varepsilon}, -R \right) \partial_i \theta(t, x) dt dx - \int_{\mathbb{R}^N} \int_{-R}^R \mathbf{1}_{\xi < u_0(x, \frac{x}{\varepsilon})} \theta(t=0, x) \psi \left(\frac{x}{\varepsilon}, \xi \right) dx d\xi \\
&\geq -\alpha_- \int_0^\infty \int_{\mathbb{R}^N} A_i \left(\frac{x}{\varepsilon}, -R \right) \partial_i \theta(t, x) dt dx - \int_{\mathbb{R}^N} \int_{-R}^R \mathbf{1}_{\xi < u_0(x, \frac{x}{\varepsilon})} \theta(t=0, x) \psi \left(\frac{x}{\varepsilon}, \xi \right) dx d\xi.
\end{aligned}$$

Passing to the limit as $\varepsilon \rightarrow 0$, we retrieve

$$\begin{aligned}
& \int_0^\infty \int_{\mathbb{R}^N} \int_{-R}^R f^0(t, x, y, \xi) [\partial_t \theta(t, x) + a_i(y, \xi) \partial_{x_i} \theta(t, x)] \psi(y, \xi) dx dy d\xi dt \\
&\geq -\alpha_- \int_0^\infty \int_{\mathbb{R}^N \times Y} A_i(y, -R) \partial_i \theta(t, x) dt dx dy - \int_{\mathbb{R}^N} \int_{-R}^R \mathbf{1}_{\xi < u_0(x, y)} \theta(t=0, x) \psi(y, \xi) dx d\xi dy \\
&= - \int_{\mathbb{R}^N} \int_{-R}^R \mathbf{1}_{\xi < u_0(x, y)} \theta(t=0, x) \psi(y, \xi) dx d\xi.
\end{aligned}$$

This means exactly that

$$\frac{\partial}{\partial t} \int_{Y \times \mathbb{R}} f^0 \psi + \frac{\partial}{\partial x_i} \int_{Y \times \mathbb{R}} a_i f^0 \psi \leq 0,$$

or in other words, that $\int_{Y \times \mathbb{R}} \mathcal{M} \psi \leq 0$ in the sense of distributions on $[0, \infty) \times \mathbb{R}^N$.

Now, we go back to the regularizations by convolution. According to the preceding subsection,

$$\begin{aligned}
A = \lim_{\varepsilon \rightarrow 0} \int_0^\infty \int_0^\infty \int_{\mathbb{R}^{2N} \times C_0} f^\varepsilon(s, z, \xi) \left\{ \partial_t \varphi(t-s, x-z) + \sum_{i=1}^N a_i \left(\frac{z}{\varepsilon}, \xi \right) \partial_i \varphi(t-s, x-z) \right\} \times \\
\times \psi \left(\frac{z}{\varepsilon}, \xi \right) \theta(t, x) d\xi dx dz ds dt.
\end{aligned}$$

Hence, in (4), we consider the test function

$$\phi(s, z, \xi) = \left[\int_0^\infty \int_{\mathbb{R}^N} \varphi(t-s, x-z) \theta(t, x) dt dx \right] \psi_\delta \left(\frac{z}{\varepsilon}, \xi \right) K(\xi),$$

where

- K is a cut-off function such that $0 \leq K \leq 1$, $K \in \mathcal{D}(\mathbb{R})$, $K(\xi) = 1$ if $|\xi| \leq R$ (R is the same parameter as before, and satisfies $R \geq \max(M, C+1)$);
- $\psi_\delta := \psi *_y \varphi_1^\delta *_\xi \varphi_2^\delta$, with $\varphi_1 \in \mathcal{D}(\mathbb{R}^N)$, $\varphi_2 \in \mathcal{D}(\mathbb{R})$, $0 \leq \varphi_i \leq 1$, $\int \varphi_i = 1$ for $i = 1, 2$, and

$$\varphi_1^\delta(y) = \frac{1}{\delta^N} \varphi_1 \left(\frac{y}{\delta} \right), \quad \varphi_2^\delta(\xi) = \frac{1}{\delta} \varphi_2 \left(\frac{\xi}{\delta} \right), \quad 0 < \delta < 1.$$

According to (4), we have

$$\begin{aligned}
& \int_0^\infty \int_{\mathbb{R}^{N+1}} f^\varepsilon(s, z, \xi) \left[\partial_s \phi(s, z, \xi) + \sum_{i=1}^N a_i \left(\frac{z}{\varepsilon}, \xi \right) \partial_{z_i} \phi(s, z, \xi) \right] dz d\xi ds \\
&+ \frac{1}{\varepsilon} \int_0^\infty \int_{\mathbb{R}^{N+1}} f^\varepsilon(s, z, \xi) a_{N+1} \left(\frac{z}{\varepsilon}, \xi \right) \partial_\xi \phi(s, z, \xi) dz d\xi ds \tag{30} \\
&- \int_0^\infty \int_{\mathbb{R}^{N+1}} m^\varepsilon(s, z, \xi) \partial_\xi \phi(s, z, \xi) dz d\xi ds + \int_{\mathbb{R}^{N+1}} \chi \left(\xi, u_0 \left(z, \frac{z}{\varepsilon} \right) \right) \phi(s=0, z, \xi) dz d\xi \\
&= 0.
\end{aligned}$$

And

$$\begin{aligned}
\partial_s \phi(s, z, \xi) &= - \left[\int_0^\infty \int_{\mathbb{R}^N} \partial_t \varphi(t-s, x-z) \theta(t, x) dt dx \right] \psi_\delta \left(\frac{z}{\varepsilon}, \xi \right) K(\xi), \\
\nabla_z \phi(s, z, \xi) &= - \left[\int_0^\infty \int_{\mathbb{R}^N} \nabla_x \varphi(t-s, x-z) \theta(t, x) dt dx \right] \psi_\delta \left(\frac{z}{\varepsilon}, \xi \right) K(\xi) \\
&\quad + \frac{1}{\varepsilon} \left[\int_0^\infty \int_{\mathbb{R}^N} \varphi(t-s, x-z) \theta(t, x) dt dx \right] (\nabla_y \psi_\delta) \left(\frac{z}{\varepsilon}, \xi \right) K(\xi), \\
\partial_\xi \phi(s, z, \xi) &= \left[\int_0^\infty \int_{\mathbb{R}^N} \varphi(t-s, x-z) \theta(t, x) dt dx \right] K(\xi) \partial_\xi \psi_\delta \left(\frac{z}{\varepsilon}, \xi \right) \\
&\quad + \left[\int_0^\infty \int_{\mathbb{R}^N} \varphi(t-s, x-z) \theta(t, x) dt dx \right] \psi_\delta \left(\frac{z}{\varepsilon}, \xi \right) \partial_\xi K(\xi) \\
\phi(s=0, z, \xi) &= \left[\int_0^\infty \int_{\mathbb{R}^N} \varphi(t, x-z) \theta(t, x) dt dx \right] \psi_\delta \left(\frac{z}{\varepsilon}, \xi \right) K(\xi) = 0.
\end{aligned}$$

Using the assumption on the sign of θ, φ , and the fact that

$$\partial_\xi \psi_\delta = (\partial_\xi \psi) *_y \varphi_1^\delta *_\xi \varphi_2^\delta \geq 0,$$

we derive

$$\left[\int_0^\infty \int_{\mathbb{R}^N} \varphi(t-s, x-z) \theta(t, x) dt dx \right] K(\xi) \partial_\xi \psi_\delta \left(\frac{z}{\varepsilon}, \xi \right) \geq 0.$$

Moreover, due to (7), (8), and the assumptions on ψ and K , we have $\partial_\xi K = 0$ on $\text{Supp } m^\varepsilon$, and, for all $\xi \in \mathbb{R}$,

$$\begin{aligned}
&\left[\int_0^\infty \int_{\mathbb{R}^N} \varphi(t-s, x-z) \theta(t, x) dt dx \right] \psi_\delta \left(\frac{z}{\varepsilon}, \xi \right) \partial_\xi K(\xi) f^\varepsilon(s, z, \xi) \\
&= \alpha_- \left[\int_0^\infty \int_{\mathbb{R}^N} \varphi(t-s, x-z) \theta(t, x) dt dx \right] \partial_\xi K(\xi) \mathbf{1}_{\xi < M}.
\end{aligned}$$

Hence, we obtain, for all $\varepsilon, \delta > 0$,

$$\begin{aligned}
& - \int f^\varepsilon(s, z, \xi) \left\{ \partial_t \varphi(t-s, x-z) + \sum_{i=1}^N a_i \left(\frac{z}{\varepsilon}, \xi \right) \partial_i \varphi(t-s, x-z) \right\} \times \\
&\quad \times \psi_\delta \left(\frac{z}{\varepsilon}, \xi \right) \theta(t, x) d\xi dx dz ds dt \\
& + \frac{1}{\varepsilon} \int f^\varepsilon(s, z, \xi) a \left(\frac{z}{\varepsilon}, \xi \right) \cdot \nabla_{y, \xi} \psi_\delta \left(\frac{z}{\varepsilon}, \xi \right) \varphi(t-s, x-z) \theta(t, x) K(\xi) dt dx ds dz d\xi \\
& + \frac{\alpha_-}{\varepsilon} \int \varphi(t-s, x-z) \theta(t, x) \partial_\xi K(\xi) a_{N+1} \left(\frac{z}{\varepsilon}, \xi \right) \mathbf{1}_{\xi < M} dt dx ds dz d\xi \\
& \geq 0.
\end{aligned}$$

Following the formal calculations above, we have to investigate the sign of the term

$$\int f^\varepsilon(s, z, \xi) a \left(\frac{z}{\varepsilon}, \xi \right) \cdot \nabla_{y, \xi} \psi_\delta \left(\frac{z}{\varepsilon}, \xi \right) \varphi(t-s, x-z) \theta(t, x) K(\xi) dt dx ds dz d\xi.$$

Since $\text{div}_{y, \xi}(a\psi) = -\partial_\xi \mu$, we have

$$\text{div}_{y, \xi}(a\psi_\delta) = -\frac{\partial \mu_\delta}{\partial \xi} + r_\delta$$

where $\mu_\delta = \mu *_y \varphi_1^\delta *_\xi \varphi_2^\delta$, and r_δ is a remainder term. Then

$$\begin{aligned}
& - \int_0^\infty \int_{\mathbb{R}^{N+1}} f^\varepsilon(s, z, \xi) \frac{\partial \mu_\delta}{\partial \xi} \left(\frac{x}{\varepsilon}, \xi \right) \left[\int_0^\infty \int_{\mathbb{R}^N} \varphi(t-s, x-z) \theta(t, x) dt dx \right] ds dz d\xi \\
& = - \int_0^\infty \int_{\mathbb{R}^{N+1}} \delta(\xi = u^\varepsilon(s, z)) \mu_\delta \left(\frac{x}{\varepsilon}, \xi \right) \left[\int_0^\infty \int_{\mathbb{R}^N} \varphi(t-s, x-z) \theta(t, x) dt dx \right] ds dz d\xi \leq 0.
\end{aligned}$$

Hence, we have to prove that as $\delta \rightarrow 0$,

$$r_\delta \rightarrow 0 \quad \text{in } L^1_{\text{loc}}(Y \times \mathbb{R}).$$

The proof is quite classical. We have

$$\begin{aligned} r_\delta(y, \xi) &= a(y, \xi)\psi * (\nabla_{y, \xi} \varphi_1^\delta \varphi_2^\delta) - [a(y, \xi)\psi] * (\nabla_{y, \xi} \varphi_1^\delta \varphi_2^\delta) \\ &= \sum_{i=1}^N \int [a_i(y, \xi) - a_i(y_1, \xi_1)] \psi(y_1, \xi_1) \partial_{y_i} \varphi_1^\delta(y - y_1) \varphi_2^\delta(\xi - \xi_1) dy_1 d\xi_1 \\ &\quad + \int [a_{N+1}(y, \xi) - a_{N+1}(y_1, \xi_1)] \psi(y_1, \xi_1) \varphi_1^\delta(y - y_1) \partial_\xi \varphi_2^\delta(\xi - \xi_1) dy_1 d\xi_1 \end{aligned}$$

Thus, we compute, for $(y, y_1, \xi, \xi_1) \in \mathbb{R}^{2N+2}$, $1 \leq i \leq N+1$,

$$\begin{aligned} a_i(y, \xi) - a_i(y_1, \xi_1) &= (y - y_1) \cdot \int_0^1 \nabla_y a_i(\tau y + (1 - \tau)y_1, \tau \xi + (1 - \tau)\xi_1) d\tau \\ &\quad + (\xi - \xi_1) \cdot \int_0^1 \partial_\xi a_i(\tau y + (1 - \tau)y_1, \tau \xi + (1 - \tau)\xi_1) d\tau. \end{aligned}$$

Set, for $1 \leq k, i \leq N$, $y \in \mathbb{R}^N$, $\xi \in \mathbb{R}$,

$$\begin{aligned} \phi_{k,i}(y, \xi) &= y_k \frac{\partial \varphi_1}{\partial y_i}(y) \varphi_2(\xi), & \phi_{k,N+1}(y, \xi) &= y_k \frac{\partial \varphi_2}{\partial \xi}(\xi) \varphi_1(y), \\ \zeta_i(y, \xi) &= \xi \frac{\partial \varphi_1}{\partial y_i}(y) \varphi_2(\xi), & \zeta_{N+1}(y, \xi) &= \xi \frac{\partial \varphi_2}{\partial \xi}(\xi) \varphi_1(y). \end{aligned}$$

Notice that for $1 \leq k \leq N$, $1 \leq i \leq N+1$,

$$\int_{\mathbb{R}^{N+1}} \phi_{k,i} = -\delta_{k,i}, \quad \int_{\mathbb{R}^{N+1}} \zeta_i = -\delta_{N+1,i}.$$

Then

$$\begin{aligned} r_\delta(y, \xi) &= \sum_{i=1}^{N+1} \sum_{k=1}^N \int \frac{\partial a_i}{\partial y_k}(\tau y + (1 - \tau)y_1, \tau \xi + (1 - \tau)\xi_1) \psi(y_1, \xi_1) \phi_{k,i}^\delta(y - y_1, \xi - \xi_1) dy_1 d\xi_1 d\tau \\ &\quad + \sum_{i=1}^{N+1} \int \frac{\partial a_i}{\partial \xi}(\tau y + (1 - \tau)y_1, \tau \xi + (1 - \tau)\xi_1) \psi(y_1, \xi_1) \zeta_i^\delta(y - y_1, \xi - \xi_1) dy_1 d\xi_1 d\tau. \end{aligned}$$

Hence as $\delta \rightarrow 0$, r_δ converges to

$$-\text{div}_{y, \xi}(a(y, \xi)) \psi(y, \xi) = 0$$

in $L^p_{\text{loc}}(\mathbb{R}^{N+1})$ for any $p < \infty$ and for all $(t, x) \in [0, \infty) \times \mathbb{R}^N$. We now pass to the limit as $\delta \rightarrow 0$, with ε fixed, and we obtain

$$\begin{aligned} & - \int f^\varepsilon(s, z, \xi) \left\{ \partial_t \varphi(t - s, x - z) + a_i\left(\frac{z}{\varepsilon}, \xi\right) \partial_i \varphi(t - s, x - z) \right\} \psi\left(\frac{z}{\varepsilon}, \xi\right) \theta(t, x) d\xi dx dz ds dt \\ & - \alpha_- \int \theta(t, x) \partial_\xi K(\xi) A\left(\frac{z}{\varepsilon}, \xi\right) \cdot \nabla_x \varphi(t - s, x - z) dt dx ds dz d\xi \\ & \geq 0. \end{aligned}$$

Passing to the limit as ε vanishes, we are led to

$$\begin{aligned} & - \int f^0(s, z, y, \xi) \left\{ \partial_t \varphi(t - s, x - z) + a_i(y, \xi) \partial_i \varphi(t - s, x - z) \right\} \times \\ & \quad \times K(\xi) \psi(y, \xi) \theta(t, x) d\xi dx dz ds dy dt \\ & - \alpha_- \int \theta(t, x) \partial_\xi K(\xi) A(y, \xi) \cdot \nabla_x \varphi(t - s, x - z) dt dx ds dy dz d\xi \\ & \geq 0. \end{aligned}$$

Since

$$\int \theta(t, x) \nabla_x \varphi(t - s, x - z) dt dx ds dz = - \left(\int \theta(t, x) dt dx \right) \left(\int \nabla_z \varphi(s, z) ds dz \right) = 0,$$

we deduce that

$$\int f^0(s, z, y, \xi) \left\{ \partial_t \varphi(t - s, x - z) + \sum_{i=1}^N a_i(y, \xi) \partial_i \varphi(t - s, x - z) \right\} \times \\ \times \psi(y, \xi) K(\xi) \theta(t, x) d\xi dx dz ds dy dt \leq 0,$$

which means that f^0 satisfies condition (14). There only remains to check the strong continuity of f at time $t = 0$.

2.4. Strong continuity at time $t = 0$

The continuity property for f^0 is inherited from uniform continuity properties at time $t = 0$ for the sequence f^ε . This is strongly linked to the well-preparedness of the initial data (condition (15)), that is, the fact that for all $x \in \mathbb{R}^N$, $u_0(x, \cdot)$ is an entropy solution of the cell problem

$$\operatorname{div}_y A(y, u_0(x, y)) = 0.$$

First, let us consider a regularization of the initial data

$$g_n^\delta = f_0 *_{x} \rho_n *_{y} \varphi_1^\delta *_{\xi} \varphi_2^\delta.$$

with $\rho_n \in \mathcal{D}(\mathbb{R}_x^N)$ a convolution kernel ($n \in \mathbb{N}$), $\delta > 0$, and φ_i^δ defined as in the previous subsection. Then we can write

$$\begin{aligned} & \sum_{i=1}^N a_i \left(\frac{x}{\varepsilon}, \xi \right) \cdot \frac{\partial}{\partial x_i} \left[g_n^\delta \left(x, \frac{x}{\varepsilon}, \xi \right) \right] + \frac{1}{\varepsilon} a_{N+1} \left(\frac{x}{\varepsilon}, \xi \right) \frac{\partial}{\partial \xi} g_n^\delta \left(x, \frac{x}{\varepsilon}, \xi \right) \\ &= \frac{1}{\varepsilon} a \left(\frac{x}{\varepsilon}, \xi \right) \cdot (\nabla_{y, \xi} g_n^\delta) \left(x, \frac{x}{\varepsilon}, \xi \right) + \sum_{i=1}^N a_i \left(\frac{x}{\varepsilon}, \xi \right) \left(\frac{\partial}{\partial x_i} g_n^\delta \right) \left(x, \frac{x}{\varepsilon}, \xi \right) \\ &:= r_{n, \delta}^\varepsilon. \end{aligned} \quad (31)$$

Notice that

$$\|\nabla_x g_n^\delta\|_{L^\infty(\mathbb{R}^N \times Y \times \mathbb{R})} \leq \|\nabla_x \rho_n\|_{L^1(\mathbb{R}^N)},$$

and

$$a(y, \xi) \nabla_{y, \xi} g_n^\delta(x, y, \xi) = \partial_\xi m_n^\delta + r_n^\delta,$$

where

$$\begin{aligned} m_n^\delta &= m_0 *_{x} \rho_n *_{y} \varphi_1^\delta *_{\xi} \varphi_2^\delta, \\ r_n^\delta(x, y, \xi) &= a(y, \xi) \nabla_{y, \xi} g_n^\delta(x, y, \xi) - [a f_0 *_{x} \rho_n] *_{y, \xi} \nabla_{y, \xi} \varphi_1^\delta(y) \varphi_2^\delta(\xi). \end{aligned}$$

Then for all $n \in \mathbb{N}$, for all $x \in \mathbb{R}^N$, r_n^δ vanishes as $\delta \rightarrow 0$ in $L_{\text{loc}}^1(Y \times \mathbb{R})$ and almost everywhere. The proof of this fact is exactly the same as in the preceding subsection, and thus, we leave the details to the reader. As a consequence, we can write

$$r_{n, \delta}^\varepsilon(x, \xi) = \frac{1}{\varepsilon} \partial_\xi m_n^\delta \left(x, \frac{x}{\varepsilon}, \xi \right) + R_{n, \delta}^\varepsilon(x, \xi),$$

and there exists a constant C_n , independent of ε , such that for all $n \in \mathbb{N}$, for all $\varepsilon > 0$, and for almost every x, ξ

$$\limsup_{\delta \rightarrow 0} |R_{n, \delta}^\varepsilon(x, \xi)| \leq C_n.$$

Moreover, $R_{n, \delta}^\varepsilon(x, \xi) = 0$ if $\xi > M + \delta$. In the following, we take $\delta < 1$.

Now, we multiply (4) by $1 - 2g_n^\delta(x, x/\varepsilon, \xi)$, and (31) by $1 - 2f^\varepsilon(t, x, \xi)$. Setting

$$\begin{aligned} h_{n,\delta}^\varepsilon(t, x, \xi) &:= f^\varepsilon(t, x, \xi) + g_n^\delta\left(x, \frac{x}{\varepsilon}, \xi\right) - 2f^\varepsilon(t, x, \xi)g_n^\delta\left(x, \frac{x}{\varepsilon}, \xi\right) \\ &= \left|f^\varepsilon(t, x, \xi) - g_n^\delta\left(x, \frac{x}{\varepsilon}, \xi\right)\right|^2 + g_n^\delta\left(x, \frac{x}{\varepsilon}, \xi\right) - \left|g_n^\delta\left(x, \frac{x}{\varepsilon}, \xi\right)\right|^2, \end{aligned}$$

we obtain

$$\begin{aligned} \frac{\partial}{\partial t} h_{n,\delta}^\varepsilon(t, x, \xi) + \sum_{i=1}^N a_i\left(\frac{x}{\varepsilon}, \xi\right) \partial_{x_i} h_{n,\delta}^\varepsilon(t, x, \xi) + \frac{1}{\varepsilon} a_{N+1}\left(\frac{x}{\varepsilon}, \xi\right) \partial_\xi h_{n,\delta}^\varepsilon(t, x, \xi) &= \\ = \frac{\partial m^\varepsilon}{\partial \xi} \left[1 - 2g_n^\delta\left(x, \frac{x}{\varepsilon}, \xi\right)\right] + \frac{1}{\varepsilon} \partial_\xi m_n^\delta\left(x, \frac{x}{\varepsilon}, \xi\right) [1 - 2f^\varepsilon(t, x, \xi)] + R_{n,\delta}^\varepsilon(x, \xi) [1 - 2f^\varepsilon(t, x, \xi)]. \end{aligned} \quad (32)$$

Notice that

$$\begin{aligned} \partial_\xi [1 - 2f^\varepsilon(t, x, \xi)] &= 2\delta(\xi = u^\varepsilon(t, x)) \geq 0 \\ \frac{\partial}{\partial \xi} \left(1 - 2g_n^\delta\left(x, \frac{x}{\varepsilon}, \xi\right)\right) &\geq 0. \end{aligned}$$

Notice also that $f^\varepsilon(t, x, \xi) - g_n^\delta(x, x/\varepsilon, \xi) = 0$ if $|\xi|$ is large enough ($|\xi| > M + 1$), whence $h_{n,\delta}^\varepsilon$ has compact support in ξ .

Take a cut-off function $\zeta \in C^\infty(\mathbb{R}^N)$ such that $\zeta(x) = e^{-|x|}$ when $|x| \geq 1$, and $\frac{1}{\varepsilon} \leq \zeta(x) \leq 1$ for $|x| \leq 1$. Then there exists a constant C such that

$$|\nabla_x \zeta(x)| \leq C\zeta(x) \quad \forall x \in \mathbb{R}^N.$$

Hence, multiplying (32) by $\zeta(x)$ and integrating on \mathbb{R}^{N+1} , we obtain a bound of the type

$$\begin{aligned} \frac{d}{dt} \int_{\mathbb{R}^{N+1}} h_{n,\delta}^\varepsilon(t, x, \xi) \zeta(x) dx d\xi &\leq C \int_{\mathbb{R}^{N+1}} h_{n,\delta}^\varepsilon(t, x, \xi) \zeta(x) dx d\xi \\ &\quad + \int_{\mathbb{R}^{N+1}} |R_{n,\delta}^\varepsilon(x, \xi)| |1 - 2f^\varepsilon(t, x, \xi)| \zeta(x) dx d\xi. \end{aligned}$$

Using Gronwall's lemma and passing to the limit as $\delta \rightarrow 0$ with ε and $n \in \mathbb{N}$ fixed, we retrieve, for all $t \geq 0$,

$$\begin{aligned} \int_{\mathbb{R}^{N+1}} \left|f^\varepsilon(t, x, \xi) - g_n\left(x, \frac{x}{\varepsilon}, \xi\right)\right|^2 \zeta(x) dx d\xi &\leq e^{Ct} \int_{\mathbb{R}^{N+1}} \left|f_0\left(x, \frac{x}{\varepsilon}, \xi\right) - g_n\left(x, \frac{x}{\varepsilon}, \xi\right)\right|^2 \zeta(x) dx d\xi \\ &\quad + e^{Ct} \int_{\mathbb{R}^{N+1}} \left[g_n\left(x, \frac{x}{\varepsilon}, \xi\right) - \left|g_n\left(x, \frac{x}{\varepsilon}, \xi\right)\right|^2\right] \zeta(x) dx d\xi \\ &\quad + C_n(e^{Ct} - 1), \end{aligned}$$

where the constant C_n does not depend on ε , and $g_n = f_0 *_{x} \rho_n$. And for all $n \in \mathbb{N}$, $\varepsilon > 0$, we have

$$\begin{aligned} &\int_{\mathbb{R}^{N+1}} \left|f_0\left(x, \frac{x}{\varepsilon}, \xi\right) - g_n\left(x, \frac{x}{\varepsilon}, \xi\right)\right|^2 \zeta(x) dx d\xi \\ &\leq \int_{\mathbb{R}^{N+1}} \int_{\mathbb{R}^N} \left|f_0\left(x, \frac{x}{\varepsilon}, \xi\right) - f_0\left(x', \frac{x}{\varepsilon}, \xi\right)\right|^2 \rho_n(x - x') \zeta(x) dx dx' d\xi \\ &\leq \int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \left|u_0\left(x, \frac{x}{\varepsilon}, \xi\right) - u_0\left(x', \frac{x}{\varepsilon}, \xi\right)\right| \rho_n(x - x') \zeta(x) dx dx' \\ &\leq \int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \sup_{y \in Y} |u_0(x, y, \xi) - u_0(x', y, \xi)| \rho_n(x - x') \zeta(x) dx dx'. \end{aligned}$$

The right-hand side of the above inequality vanishes as $n \rightarrow \infty$ because $u_0 \in L^1_{\text{loc}}(\mathbb{R}^N, \mathcal{C}_{\text{per}}(Y))$. Similarly,

$$\begin{aligned}
& \int_{\mathbb{R}^{N+1}} \left[g_n \left(x, \frac{x}{\varepsilon}, \xi \right) - \left| g_n \left(x, \frac{x}{\varepsilon}, \xi \right) \right|^2 \right] \zeta(x) dx d\xi \\
& \leq \int_{\mathbb{R}^{N+1}} \left[g_n \left(x, \frac{x}{\varepsilon}, \xi \right) - f_0 \left(x, \frac{x}{\varepsilon}, \xi \right) \right] \zeta(x) dx d\xi \\
& \quad + \int_{\mathbb{R}^{N+1}} \left[f_0 \left(x, \frac{x}{\varepsilon}, \xi \right)^2 - g_n \left(x, \frac{x}{\varepsilon}, \xi \right)^2 \right] \zeta(x) dx d\xi \\
& \leq 3 \int_{\mathbb{R}^{N+1}} \left| g_n \left(x, \frac{x}{\varepsilon}, \xi \right) - f_0 \left(x, \frac{x}{\varepsilon}, \xi \right) \right| \zeta(x) dx d\xi \\
& \leq 3 \int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \sup_{y \in Y} |u_0(x, y, \xi) - u_0(x', y, \xi)| \rho_n(x - x') \zeta(x) dx dx'.
\end{aligned}$$

Hence, we deduce that there exists a function $\omega : [0, \infty) \rightarrow [0, \infty)$, independent of ε and satisfying $\lim_{t \rightarrow 0} \omega(t) = 0$, such that

$$\int_{\mathbb{R}^{N+1}} \left| f^\varepsilon(t, x, \xi) - f_0 \left(x, \frac{x}{\varepsilon}, \xi \right) \right| \zeta(x) dx d\xi \leq \omega(t)$$

for all $t > 0$.

Then, we prove that the same property holds for the function f^0 , that is, the two-scale limit of the sequence f^ε . Indeed, we write

$$\left| f^\varepsilon(t, x, \xi) - \mathbf{1}_{\xi < u_0(x, \frac{x}{\varepsilon})} \right| = f^\varepsilon - 2f^\varepsilon \mathbf{1}_{\xi < u_0(x, \frac{x}{\varepsilon})} + \mathbf{1}_{\xi < u_0(x, \frac{x}{\varepsilon})};$$

let $\theta \in L^\infty([0, \infty))$ with compact support and such that $\theta \geq 0$. Then for all $\varepsilon > 0$,

$$\int_0^\infty \int_{\mathbb{R}^{N+1}} \left[f^\varepsilon - 2f^\varepsilon \mathbf{1}_{\xi < u_0(x, \frac{x}{\varepsilon})} + \mathbf{1}_{\xi < u_0(x, \frac{x}{\varepsilon})} \right] \zeta(x) \theta(t) dx d\xi dt \leq \int_0^\infty \omega(t) \theta(t) dt.$$

Since $u_0 \in L^1_{\text{loc}}(\mathbb{R}^N, \mathcal{C}_{\text{per}}(Y))$, it is an admissible test function in the sense of G. Allaire (see [1]); we deduce that $\mathbf{1}_{\xi < u_0}$ is also an admissible test function. This is not entirely obvious because it is a discontinuous function of u_0 . However, this difficulty can be overcome thanks to an argument similar to the one developed below in subsection 3.3, and which we do not reproduce here. Thus, we can pass to the two-scale limit in the above inequality. We obtain

$$\begin{aligned}
\int_0^\infty \int_{\mathbb{R}^{N+1} \times Y} (f^0(t, x, y, \xi) - f^0(t, x, y, \xi))^2 + |f^0(t, x, y, \xi) - \mathbf{1}_{\xi < u_0(x, y)}|^2 \theta(t) \zeta(x) dt dx dy d\xi & \leq \\
& \leq \int_0^\infty \theta(t) \omega(t) dt
\end{aligned}$$

Notice that $f^0 - |f^0|^2 \geq 0$ almost everywhere. As a consequence, taking $\theta(t) = \mathbf{1}_{0 < t < \tau}$, with $\tau > 0$ arbitrary, we deduce that

$$\frac{1}{\tau} \int_0^\tau |f^0(t) - \chi(\xi, u_0(x, y))|^2 \zeta(x) dt dx dy \leq \frac{1}{\tau} \int_0^\tau \omega(t) dt,$$

and the left-hand side vanishes as $\tau \rightarrow 0$. Thus the continuity property is satisfied at time $t = 0$.

Hence, we have proved that any two-scale limit of the sequence f^ε is a solution of the limit system. Thus the existence result in Theorem 1 is proved, as well as the convergence result of Theorem 2. We now tackle the proof of the uniqueness and rigidity results of Theorem 1. The strong convergence result of Theorem 1 will follow from the rigidity.

3. Uniqueness of solutions of the limit evolution problem

In this section, we prove the second and the third point in Theorem 1, that is, if f is any solution of the limit evolution problem, then there exists a function $u \in L^\infty([0, \infty) \times \mathbb{R}^N \times Y)$ such that $f(t, x, y, \xi) = \mathbf{1}_{\xi < u(t, x, y)}$ almost everywhere, and if $f_1 = \mathbf{1}_{\xi < u_1}$, $f_2 = \mathbf{1}_{\xi < u_2}$ are two generalized kinetic solutions, then the contraction principle (17) holds.

3.1. The rigidity result

Let f be a generalized kinetic solution of the limit problem, with initial data $\mathbf{1}_{\xi < u_0}$. The rigidity result relies on the comparison between f and f^2 . Precisely, we prove that $f = f^2$ almost everywhere, and since $\partial_\xi f \leq 0$, there exists a function u such that $f = \mathbf{1}_{\xi < u}$. Thus, we now turn to the derivation of the equality $f = f^2$.

Let $\delta > 0$ arbitrary, and let $\theta_1 \in \mathcal{D}(\mathbb{R})$, $\theta_2 \in \mathcal{D}(\mathbb{R}^N)$ such that

$$\begin{aligned} \theta_1 &\geq 0, \quad \theta_2 \geq 0, \\ \int_{\mathbb{R}} \theta_1 &= \int_{\mathbb{R}^N} \theta_2 = 1, \\ \text{Supp } \theta_1 &\subset [-1, 0] \text{ and } \theta_1(0) = 0. \end{aligned}$$

We set, for $(t, x) \in \mathbb{R}^{N+1}$

$$\theta^\delta(t, x) = \frac{1}{\delta^{N+1}} \theta_1\left(\frac{t}{\delta}\right) \theta_2\left(\frac{x}{\delta}\right).$$

Set $f^\delta := f *_{t,x} \theta^\delta$, $\mathcal{M}^\delta := \mathcal{M} *_{t,x} \theta^\delta$. Then f^δ is a solution of

$$\frac{\partial f^\delta}{\partial t} + \sum_{i=1}^N a_i(y, \xi) \frac{\partial f^\delta}{\partial x_i} = \mathcal{M}^\delta.$$

Moreover, f^δ satisfies the following properties

$$0 \leq f^\delta \leq 1, \quad (33)$$

$$\text{div}_{y,\xi}(a(y, \xi) f^\delta) = \partial_\xi m *_{t,x} \theta^\delta, \quad (34)$$

$$\partial_\xi f^\delta \leq 0, \quad (35)$$

$$f^\delta(\cdot, \xi) = 0 \quad \text{if } \xi > M, \quad f^\delta(\cdot, \xi) = 1 \quad \text{if } \xi < -M, \quad (36)$$

whereas \mathcal{M}^δ satisfies

$$\mathcal{M}^\delta \in \mathcal{C}((0, T) \times \mathbb{R}^N, L^2(Y \times \mathbb{R})) \cap L^\infty([0, \infty) \times \mathbb{R}^N \times Y \times \mathbb{R}), \quad (37)$$

$$\mathcal{M}^\delta(\cdot, \xi) = 0 \quad \text{if } |\xi| > M, \quad (38)$$

$$\int_{Y \times \mathbb{R}} \mathcal{M}^\delta(t, x) \psi \leq 0 \quad \forall \psi \in \mathcal{G} \quad \forall (t, x) \in [0, \infty) \times \mathbb{R}^N. \quad (39)$$

In particular, notice that $(1 - 2f^\delta(t, x)) \in \mathcal{G}$ for all t, x , and $f^\delta(t, x, y, \xi) - f^\delta(t, x, y, \xi)^2 = 0$ if $|\xi| > M$.

Let $\zeta \in \mathcal{C}^\infty(\mathbb{R}^N)$ be a cut-off function as in the previous subsection. We multiply by $(1 - 2f^\delta)\zeta$ the equation satisfied by f^δ , and we integrate over $\mathbb{R}^N \times Y \times \mathbb{R}$. We obtain

$$\frac{d}{dt} \int_{\mathbb{R}^N \times Y \times \mathbb{R}} (f^\delta - |f^\delta|^2) \zeta - \int_{\mathbb{R}^N \times Y \times \mathbb{R}} a_i(y, \xi) \partial_i \zeta(x) (f^\delta - |f^\delta|^2) = \int_{\mathbb{R}^N \times Y \times \mathbb{R}} \mathcal{M}^\delta (1 - 2f^\delta) \zeta \leq 0.$$

We then deduce successively, using Gronwall's lemma,

$$\begin{aligned} \frac{d}{dt} \int_{\mathbb{R}^N \times Y \times \mathbb{R}} (f^\delta - |f^\delta|^2) \zeta &\leq C \int_{\mathbb{R}^N \times Y \times \mathbb{R}} (f^\delta - |f^\delta|^2) \zeta, \\ \int_{\mathbb{R}^N \times Y \times \mathbb{R}} (f^\delta(t) - |f^\delta(t)|^2) \zeta &\leq e^{Ct} \int_{\mathbb{R}^N \times Y \times \mathbb{R}} (f^\delta(t=0) - |f^\delta(t=0)|^2) \zeta \quad \forall t > 0, \\ \int_0^T \int_{\mathbb{R}^N \times Y \times \mathbb{R}} (f^\delta - |f^\delta|^2) \zeta &\leq \frac{e^{CT} - 1}{C} \int_{\mathbb{R}^N \times Y \times \mathbb{R}} (f^\delta(t=0) - |f^\delta(t=0)|^2) \zeta, \end{aligned} \quad (40)$$

and the constant C depends only on $\|a\|_{L^\infty(Y \times [-R, R])}$.

Now, let us check that $f^\delta(t=0)$ strongly converges towards $\mathbf{1}_{\xi < u_0} = f_0$ at time $t=0$. In fact, the main difference between the rigidity result of Theorem 1 and the one for generalized kinetic solutions of scalar conservation laws (see chapter 4 in [23]) lies in this particular point. Indeed, in the case of scalar conservation laws, the continuity property can be inferred from the equation itself; in the present case, the lack of structure of the right-hand side \mathcal{M} prevents us from deriving such a result, and hence the continuity of solutions at time $t=0$ is a necessary assumption in Definition 1.

Using hypothesis (13), we write, for almost every x, y, ξ ,

$$\begin{aligned} f^\delta(t=0, x, y, \xi) &= \int_{\mathbb{R}^{N+1}} f(s, z, y, \xi) \theta^\delta(-s, x-z) ds dz \\ f^\delta(t=0, x, y, \xi) - f_0 * \theta_2^\delta(x, y, \xi) &= \int_{\mathbb{R}^{N+1}} (f(s, z, y, \xi) - f_0(z, y, \xi)) \theta^\delta(-s, x-z) ds dz. \end{aligned}$$

As a consequence, for all $\delta > 0$

$$\begin{aligned} & \int_{\mathbb{R}^N \times Y \times \mathbb{R}} |f^\delta(t=0) - f_0 * \theta_2^\delta|^2 \zeta(x) dx dy d\xi \\ & \leq \int_{\mathbb{R}^N \times Y \times \mathbb{R}} \int_{\mathbb{R}^{N+1}} |f(s, z, y, \xi) - f_0(z, y, \xi)|^2 \zeta(x) \theta^\delta(-s, x-z) dx dy d\xi ds dz \\ & \leq \int_{\mathbb{R}} \|f(s) - f_0\|_{L^2(\mathbb{R}^N \times Y \times \mathbb{R}, \zeta(x) dx dy d\xi)}^2 \frac{1}{\delta} \theta_1\left(\frac{-s}{\delta}\right) ds dx dy d\xi + 2M|Y| \|\zeta - \zeta * \check{\theta}_2^\delta\|_{L^1(\mathbb{R}^N)} \\ & \leq \frac{C}{\delta} \int_0^\delta \|f(s) - f_0\|_{L^2(\mathbb{R}^N \times Y \times \mathbb{R}, \zeta(x) dx dy d\xi)}^2 ds + 2M|Y| \|\zeta - \zeta * \check{\theta}_2^\delta\|_{L^1(\mathbb{R}^N)}. \end{aligned}$$

The right-hand side of the last inequality vanishes as $\delta \rightarrow 0$, and thus $f^\delta(t=0)$ converges towards f_0 as $\delta \rightarrow 0$ in $L^2(\mathbb{R}^N \times Y \times \mathbb{R}, \zeta(x) dx dy d\xi)$, and hence also in $L^1(\mathbb{R}^N \times Y \times \mathbb{R}, \zeta(x) dx dy d\xi)$. Consequently,

$$\int_{\mathbb{R}^N \times Y \times \mathbb{R}} (f^\delta(t=0) - f_0)^2 \zeta \rightarrow 0 \quad \text{as } \delta \rightarrow 0.$$

Above, we have used the fact that $f_0 = \mathbf{1}_{\xi < u_0}$, and thus $f_0 = f_0^2$.

Now, we pass to the limit as $\delta \rightarrow 0$ in (40); we obtain, for all $T > 0$,

$$\int_0^T \int_{\mathbb{R}^N \times Y \times \mathbb{R}} (f - f^2) \varphi \leq 0.$$

Since the integrand in the left-hand side is non-negative, we deduce that $f = f^2$ almost everywhere. The rigidity property follows.

3.2. Contraction principle

Let f_1, f_2 be two generalized kinetic solutions of the limit problem; we denote by M_1, M_2 , and $\mathcal{M}_1, \mathcal{M}_2$, the constants and distributions associated to f_1, f_2 , respectively. Without loss of generality, we assume that $M_1 \leq M_2$. According to the rigidity result, there exist functions $u_1, u_2 \in L^\infty([0, \infty) \times \mathbb{R}^N \times Y) \cap L^1([0, \infty), L^1(\mathbb{R}^N \times Y))$ such that $f_i = \mathbf{1}_{\xi < u_i}$.

As in the previous subsection, we regularize f_i, \mathcal{M}_i by convolution in the variables t, x , and we denote by $f_i^\delta, \mathcal{M}_i^\delta$ the functions thus obtained. The strategy of the proof is the same as in [23], Theorem 4.3.1. The idea is to derive an inequality of the type

$$\frac{d}{dt} \int |f_1(t, x, y, \xi) - f_2(t, x, y, \xi)| \zeta(x) dx dy d\xi \leq C \int |f_1(t, x, y, \xi) - f_2(t, x, y, \xi)| \zeta(x) dx dy d\xi, \quad (41)$$

where ζ is a cut-off function as in the previous section.

Since $|f_1(t) - f_2(t)| = |f_1(t) - f_2(t)|^2 = f_1 + f_2 - 2f_1f_2$, let us first write the equation satisfied by $g^\delta := f_1^\delta + f_2^\delta - 2f_1^\delta f_2^\delta$. We compute

$$\begin{cases} \partial_t f_1^\delta + \sum_{i=1}^N a_i(y, \xi) \frac{\partial}{\partial x_i} f_1^\delta = \mathcal{M}_1^\delta \\ \partial_t f_2^\delta + \sum_{i=1}^N a_i(y, \xi) \frac{\partial}{\partial x_i} f_2^\delta = \mathcal{M}_2^\delta \end{cases} \times 1 - 2f_2^\delta, \\ \begin{cases} \partial_t f_1^\delta + \sum_{i=1}^N a_i(y, \xi) \frac{\partial}{\partial x_i} f_1^\delta = \mathcal{M}_1^\delta \\ \partial_t f_2^\delta + \sum_{i=1}^N a_i(y, \xi) \frac{\partial}{\partial x_i} f_2^\delta = \mathcal{M}_2^\delta \end{cases} \times 1 - 2f_1^\delta.$$

Adding the two equations thus obtained leads to

$$\partial_t g^\delta + \sum_{i=1}^N a_i(y, \xi) \frac{\partial}{\partial x_i} g^\delta = \mathcal{M}_1^\delta [1 - 2f_2^\delta] + \mathcal{M}_2^\delta [1 - 2f_1^\delta].$$

Notice that thanks to (7), (8) and the microscopic constraints (9), (11), $1 - 2f_i^\delta(t, x) \in \mathcal{G}$ for all (t, x) . Hence

$$\int_{Y \times \mathbb{R}} \mathcal{M}_2^\delta(t, x) [1 - 2f_1^\delta(t, x)] \leq 0 \quad \forall (t, x) \in [0, \infty) \times \mathbb{R}^N,$$

and the same inequality holds if the roles of f_1 and f_2 are exchanged.

Now, take a cut-off function $\zeta \in \mathcal{C}^\infty(\mathbb{R}^N)$ satisfying the same assumptions as in the previous subsection; multiply the equation on g^δ by $\zeta(x)$, and integrate over $\mathbb{R}^N \times Y \times \mathbb{R}$; this yields

$$\frac{d}{dt} \int_{\mathbb{R}^N \times Y \times \mathbb{R}} g^\delta(t, x, y, \xi) \zeta(x) dx dy d\xi \leq C \int_{\mathbb{R}^N \times Y \times \mathbb{R}} g^\delta(t, x, y, \xi) \zeta(x) dx dy d\xi \quad \forall t > 0,$$

and thus

$$\int_{\mathbb{R}^N \times Y \times \mathbb{R}} g^\delta(t, x, y, \xi) \zeta(x) dx dy d\xi \leq e^{Ct} \int_{\mathbb{R}^N \times Y \times \mathbb{R}} g^\delta(t=0, x, y, \xi) \zeta(x) dx dy d\xi.$$

According to the strong convergence results of $f_i^\delta(t=0)$ derived in the previous section, we can pass to the limit as $\delta \rightarrow 0$. We infer that for almost every $t > 0$,

$$\begin{aligned} & \int_{\mathbb{R}^N \times Y \times \mathbb{R}} |f_1(t, x, y, \xi) - f_2(t, x, y, \xi)| \zeta(x) dx dy d\xi \\ & \leq e^{Ct} \int_{\mathbb{R}^N \times Y \times \mathbb{R}} |f_1(t=0, x, y, \xi) - f_2(t=0, x, y, \xi)| \zeta(x) dx dy d\xi. \end{aligned} \quad (42)$$

This completes the derivation of the contraction principle for the limit system. Uniqueness of solutions of the limit system follows. In particular, we deduce that the whole sequence f^ε of solutions of (4) two-scale converges towards f^0 .

3.3. Strong convergence result

Here, we explain why the strong convergence result stated in Theorem 2 holds, that is, we prove (22). This fact is rather classical, and is a direct consequence of the fact that

$$\mathbf{1}_{\xi < u^\varepsilon(t, x)} \xrightarrow{2 \text{ sc.}} \mathbf{1}_{\xi < u(t, x, y)}.$$

Let us express this result in terms of Young measures: the above two-scale convergence is strictly equivalent to the fact that the two-scale Young measure $\nu_{t, x, y}$ associated with the sequence u^ε is the Dirac mass $\delta(\xi = u(t, x, y))$ (see [23], Chapter 2). And it is well-known (see [9]) that if u is a smooth function, then

$$d\nu_{t, x, y}(\xi) = \delta(\xi = u(t, x, y)) \iff u^\varepsilon - u\left(t, x, \frac{x}{\varepsilon}\right) \rightarrow 0 \quad \text{in } L^1_{\text{loc}}.$$

However, here, the function u is not smooth, but this issue is bypassed by using convolution kernels. For the reader's convenience, we now prove the result without using two-scale Young

measures. We define $u_\delta = u *_x \varphi_\delta$, with φ_δ a standard mollifier. Let $K \in \mathcal{D}(\mathbb{R})$ such that $0 \leq K \leq 1$, and $K(\xi) = 1$ if $|\xi| \leq M$. Consider also a sequence $\theta_n \in \mathcal{C}^\infty(\mathbb{R})$ such that $0 \leq \theta_n \leq 1$, and

$$\theta_n(\xi) = 1 \text{ if } \xi < -\frac{1}{n}, \quad \theta_n(\xi) = 0 \text{ if } \xi > \frac{1}{n}.$$

Then we have

$$\begin{aligned} \left| \mathbf{1}_{\xi < u^\varepsilon(t,x)} - \mathbf{1}_{\xi < u_\delta(t,x,\frac{x}{\varepsilon})} \right|^2 &= \mathbf{1}_{\xi < u^\varepsilon(t,x)} - 2\mathbf{1}_{\xi < u_\delta(t,x,\frac{x}{\varepsilon})} \mathbf{1}_{\xi < u^\varepsilon(t,x)} + \mathbf{1}_{\xi < u_\delta(t,x,\frac{x}{\varepsilon})} \\ &= \mathbf{1}_{\min(u^\varepsilon(t,x), u_\delta(t,x,\frac{x}{\varepsilon})) < \xi < \max(u^\varepsilon(t,x), u_\delta(t,x,\frac{x}{\varepsilon}))}. \end{aligned}$$

The function $\mathbf{1}_{\xi < u_\delta(t,x,\frac{x}{\varepsilon})}$ is not smooth enough to be used as an oscillating test function. Thus we replace it by

$$\theta_n \left(\xi - u_\delta \left(t, x, \frac{x}{\varepsilon} \right) \right),$$

and we evaluate the difference: for all compact set $C \subset [0, \infty) \times \mathbb{R}^N$, for all $\delta, \varepsilon > 0$,

$$\int_C \int_{\mathbb{R}} \left| \mathbf{1}_{\xi < u_\delta(t,x,\frac{x}{\varepsilon})} - \theta_n \left(\xi - u_\delta \left(t, x, \frac{x}{\varepsilon} \right) \right) \right| K(\xi) dt dx d\xi \leq \frac{2}{n} |C|.$$

According to the two-scale convergence result, for all $n \in \mathbb{N}$,

$$\begin{aligned} \int_C \int_{\mathbb{R}} \theta_n \left(\xi - u_\delta \left(t, x, \frac{x}{\varepsilon} \right) \right) \mathbf{1}_{\xi < u^\varepsilon(t,x)} K(\xi) dt dx d\xi &\rightarrow \\ &\rightarrow \int_C \int_{\mathbb{R}} \theta_n \left(\xi - u_\delta(t,x,y) \right) \mathbf{1}_{\xi < u(t,x,y)} K(\xi) dt dx dy d\xi. \end{aligned}$$

Since the sequence $\theta_n(\xi - u_\delta)$ uniformly converges towards $\mathbf{1}_{\xi < u_\delta}$ as $n \rightarrow \infty$, we can pass to the limit as $n \rightarrow \infty$, and we deduce

$$\int_C \int_{\mathbb{R}} \mathbf{1}_{\xi < u_\delta(t,x,\frac{x}{\varepsilon})} \mathbf{1}_{\xi < u^\varepsilon(t,x)} K(\xi) dt dx d\xi \rightarrow \int_C \int_{\mathbb{R} \times Y} \mathbf{1}_{\xi < u_\delta(t,x,y)} \mathbf{1}_{\xi < u(t,x,y)} K(\xi) dt dx dy d\xi.$$

Similarly, as $\varepsilon \rightarrow 0$, for all $\delta > 0$,

$$\begin{aligned} \int_C \int_{\mathbb{R}} \mathbf{1}_{\xi < u_\delta(t,x,\frac{x}{\varepsilon})} K(\xi) dt dx d\xi &\rightarrow \int_C \int_{\mathbb{R} \times Y} \mathbf{1}_{\xi < u_\delta(t,x,y)} K(\xi) dt dx dy d\xi, \\ \int_C \int_{\mathbb{R} \times Y} \mathbf{1}_{\xi < u^\varepsilon(t,x)} K(\xi) dt dx d\xi &\rightarrow \int_C \int_{\mathbb{R}} \mathbf{1}_{\xi < u(t,x,y)} K(\xi) dt dx dy d\xi. \end{aligned}$$

Thus, as $\varepsilon \rightarrow 0$, for all $\delta > 0$,

$$\int_C \int_{\mathbb{R}} \left| \mathbf{1}_{\xi < u^\varepsilon(t,x)} - \mathbf{1}_{\xi < u_\delta(t,x,\frac{x}{\varepsilon})} \right| K(\xi) dt dx d\xi \rightarrow \int_C \int_{\mathbb{R} \times Y} \left| \mathbf{1}_{\xi < u(t,x,y)} - \mathbf{1}_{\xi < u_\delta(t,x,y)} \right| K(\xi) dt dx dy d\xi$$

On the other hand,

$$\begin{aligned} \int_C \int_{\mathbb{R}} \left| \mathbf{1}_{\xi < u^\varepsilon(t,x)} - \mathbf{1}_{\xi < u_\delta(t,x,\frac{x}{\varepsilon})} \right| K(\xi) dt dx d\xi &= \left\| u^\varepsilon(t,x) - u_\delta \left(t, x, \frac{x}{\varepsilon} \right) \right\|_{L^1(C)}, \\ \int_C \int_{\mathbb{R} \times Y} \left| \mathbf{1}_{\xi < u(t,x,y)} - \mathbf{1}_{\xi < u_\delta(t,x,y)} \right| K(\xi) dt dx dy d\xi &= \|u - u_\delta\|_{L^1(C \times Y)}. \end{aligned}$$

Hence we have proved that for all $\delta > 0$, for all compact set $C \subset [0, \infty) \times \mathbb{R}^N$,

$$\lim_{\varepsilon \rightarrow 0} \left\| u^\varepsilon(t,x) - u_\delta \left(t, x, \frac{x}{\varepsilon} \right) \right\|_{L^1(C)} = \|u - u_\delta\|_{L^1(C \times Y)}.$$

Statement (22) then follows from standard convolution results.

3.4. Application: proof of the convergence result for kinetic solutions

In this subsection, we prove Theorem 3; this result is in fact an easy consequence of the convergence result stated in Theorem 2 for entropy solutions, and of the contraction principle for the limit system. Assume that $a_{N+1} \equiv 0$, and let u^ε be a kinetic solution of equation (1), with an initial data $u_0(x, x/\varepsilon)$ such that $u_0 \in L^1(\mathbb{R}^N, \mathcal{C}_{\text{per}}(Y))$ and

$$\sum_{i=1}^N \frac{\partial}{\partial y_i} (a_i(y, \xi) \chi(\xi, u_0(x, y))) = 0 \quad (43)$$

in the sense of distributions.

For $n \in \mathbb{N}$, let $u_0^n := \text{sgn}(u_0) \inf(|u_0|, n)$. Then for all $n \in \mathbb{N}$, u_0^n belongs to $L^\infty(\mathbb{R}^N \times Y)$ and

$$u_0^n \rightarrow u_0 \quad \text{as } n \rightarrow \infty \quad \text{in } L^1(\mathbb{R}^N, \mathcal{C}_{\text{per}}(Y)).$$

Moreover, $\chi(\xi, u_0^n) = \chi(\xi, u_0) \mathbf{1}_{|\xi| < n}$, and thus for all $n \in \mathbb{N}$, u_0^n satisfies (43).

For all $n, \varepsilon > 0$, let $u_n^\varepsilon \in \mathcal{C}([0, \infty), L^1(\mathbb{R}^N)) \cap L^\infty([0, \infty) \times \mathbb{R}^N)$ be the unique entropy solution of equation (1) with initial data $u_0^n(x, x/\varepsilon)$. Then by the contraction principle for kinetic solutions of scalar conservation laws, we have

$$\forall n \in \mathbb{N}, \quad \|u^\varepsilon - u_n^\varepsilon\|_{L^\infty([0, \infty), L^1(\mathbb{R}^N))} \leq \left\| u_0 \left(x, \frac{x}{\varepsilon} \right) - u_0^n \left(x, \frac{x}{\varepsilon} \right) \right\|_{L^1(\mathbb{R}^N)} \leq \|u_0 - u_0^n\|_{L^1(\mathbb{R}^N, \mathcal{C}_{\text{per}}(Y))}.$$

On the other hand, for all $n \in \mathbb{N}$, let $\mathbf{1}_{\xi < u_n}$ be the unique solution of the limit system with initial data $\mathbf{1}_{\xi < u_0^n}$. By the contraction principle for solutions of the limit system (see inequality (42)), we have, for all integers $n, m \in \mathbb{N}$, for all $t \geq 0$,

$$\begin{aligned} \int_{\mathbb{R}^N \times Y} |u_n(t, x, y) - u_m(t, x, y)| \zeta(x) dx dy &\leq e^{Ct} \int_{\mathbb{R}^N \times Y} |u_0^n(t, x, y) - u_0^m(t, x, y)| \zeta(x) dx dy \\ &\leq e^{Ct} \|u_0^n - u_0^m\|_{L^1(\mathbb{R}^N, \mathcal{C}_{\text{per}}(Y))}, \end{aligned} \quad (44)$$

where $\zeta \in \mathcal{C}^\infty(\mathbb{R}^N)$ is a cut-off function satisfying the same hypotheses as in the previous subsections.

Consequently, the sequence $(u_n)_{n \in \mathbb{N}}$ is a Cauchy sequence in $L_{\text{loc}}^\infty([0, \infty), L^1(\mathbb{R}^N \times Y, \zeta(x) dx dy))$; thus there exists a function $u \in L_{\text{loc}}^\infty([0, \infty), L^1(\mathbb{R}^N \times Y, \zeta(x) dx dy))$ such that u_n converges towards u as $n \rightarrow \infty$ in $L_{\text{loc}}^\infty([0, \infty), L^1(\mathbb{R}^N \times Y, \zeta(x) dx dy))$. Moreover, the limit u is independent of the chosen sequence u_0^n thanks to (44): indeed, let v_0^n, w_0^n be two sequences converging towards u_0 in $L^1(\mathbb{R}^N, \mathcal{C}_{\text{per}}(Y))$, and giving rise to functions v and w respectively. We construct the sequence

$$u_0^n = \begin{cases} v_0^n & \text{if } n \text{ is even,} \\ w_0^n & \text{if } n \text{ is odd.} \end{cases}$$

Then the sequence u_0^n converges towards u_0 in $L^1(\mathbb{R}^N, \mathcal{C}_{\text{per}}(Y))$, and thus the corresponding sequence u_n converges towards u , while u_{2n} converges towards v and u_{2n+1} towards w . By uniqueness of the limit, $u = v = w$.

On the other hand, since the sequence $f^\varepsilon = \chi(\xi, u^\varepsilon)$ is bounded in L^∞ , there exists a sequence $(\varepsilon_k)_{k \in \mathbb{N}}$ of positive numbers, $\varepsilon_k \rightarrow 0$, and a function $f \in L^\infty([0, \infty) \times \mathbb{R}^N \times Y \times \mathbb{R})$, such that

$$\chi(\xi, u^{\varepsilon_k}(t, x)) \xrightarrow{2\text{-sc.}} f(t, x, y, \xi).$$

Now, for all $k, n \in \mathbb{N}$,

$$\|\chi(\xi, u^{\varepsilon_k}) - \chi(\xi, u_n^{\varepsilon_k})\|_{L^\infty([0, \infty), L^1(\mathbb{R}^{N+1}))} \leq \|u_0 - u_0^n\|_{L^1(\mathbb{R}^N, \mathcal{C}_{\text{per}}(Y))},$$

and for all $n \in \mathbb{N}$, since $\chi(\xi, u) = \mathbf{1}_{\xi < u} - \mathbf{1}_{\xi < 0}$, we have, as $k \rightarrow \infty$,

$$\chi(\xi, u_n^{\varepsilon_k}) \xrightarrow{2\text{-sc.}} \chi(\xi, u_n).$$

Let $\varphi \in \mathcal{D}_{\text{per}}([0, \infty) \times \mathbb{R}^N \times Y \times \mathbb{R})$. According to the definition of two-scale convergence,

$$\begin{aligned} \int_0^\infty \int_{\mathbb{R}^{N+1}} [\chi(\xi, u^{\varepsilon_k}(t, x)) - \chi(\xi, u_n^{\varepsilon_k}(t, x))] \varphi \left(t, x, \frac{x}{\varepsilon_k}, \xi \right) dt dx d\xi &\rightarrow \\ &\rightarrow \int_0^\infty \int_{\mathbb{R}^N \times Y \times \mathbb{R}} [f(t, x, y, \xi) - \chi(\xi, u_n(t, x, y))] \varphi(t, x, y, \xi) dt dx dy d\xi. \end{aligned}$$

And for all $k \in \mathbb{N}$, the following inequality holds:

$$\begin{aligned} \left| \int_0^\infty \int_{\mathbb{R}^{N+1}} [\chi(\xi, u^{\varepsilon_k}(t, x)) - \chi(\xi, u_n^{\varepsilon_k}(t, x))] \varphi \left(t, x, \frac{x}{\varepsilon_k}, \xi \right) dt dx d\xi \right| &\leq \\ &\leq \|\varphi\|_{L^1([0, \infty), L^\infty(\mathbb{R}^N \times Y \times \mathbb{R}))} \|u_0 - u_0^n\|_{L^1(\mathbb{R}^N, \mathcal{C}_{\text{per}}(Y))}. \end{aligned}$$

Passing to the limit as $k \rightarrow \infty$, we deduce that for all $n \in \mathbb{N}$, $\varphi \in \mathcal{D}_{\text{per}}([0, \infty) \times \mathbb{R}^N \times Y \times \mathbb{R})$,

$$\begin{aligned} \left| \int_0^\infty \int_{\mathbb{R}^N \times Y \times \mathbb{R}} [f(t, x, y, \xi) - \chi(\xi, u_n(t, x, y))] \varphi(t, x, y, \xi) dt dx d\xi \right| &\leq \\ &\leq \|u_0 - u_0^n\|_{L^1(\mathbb{R}^N, \mathcal{C}_{\text{per}}(Y))} \|\varphi\|_{L^1([0, \infty), L^\infty(\mathbb{R}^N \times Y \times \mathbb{R}))}. \end{aligned}$$

Thus, we pass to the limit as $n \rightarrow \infty$ and we infer that $f = \chi(\xi, u(t, x, y))$ almost everywhere. Hence the limit is unique, and the whole sequence $\chi(\xi, u^\varepsilon)$ converges (in the sense of two-scale convergence).

Eventually, let us pass to the limit as $n \rightarrow \infty$ in the limit evolution problem for $\chi(\xi, u_n)$. We set $f = \chi(\xi, u)$, and define the distribution

$$\mathcal{M} := \partial_t f + a(y, \xi) \cdot \nabla_x f.$$

Then $\mathcal{M}_n \rightharpoonup \mathcal{M}$ in the sense of distributions, and it is easily checked that inequality (19) is preserved when passing to the (weak) limit. Thus \mathcal{M} satisfies (19).

In the divergence-free case, the main difference between the L^∞ and the L^1 setting, that is, Theorem 2 and Theorem 3, lies in the fact that uniqueness for the limit system in the L^1 setting seems difficult to derive; indeed, the proof of uniqueness in the L^∞ case uses several times the fact that the distribution \mathcal{M} has compact support. In a L^1 setting, this assumption ought to be replaced by a hypothesis expressing that \mathcal{M} vanishes as $|\xi| \rightarrow \infty$, in some sense. But it is unclear how to retrieve such a property from the hydrodynamic limit (see section 4), for instance. The above argument only proves that uniqueness holds among L^1 solutions which are obtained as the limit of a sequence of L^∞ solutions. Thus we have left open the correct notion of limit system in a weak L^1 setting, and the derivation of uniqueness therein.

Nonetheless, we wish to stress that the contraction principle in the L^∞ setting is sufficient to ensure that the whole sequence $\chi(\xi, u^\varepsilon)$ converges, even if uniqueness for the limit system fails.

4. A relaxation model for the limit evolution problem

In this section, we exhibit another way of finding solutions of the limit system in the divergence-free case. Indeed, the existence result of Theorem 1 was proved by passing to the two-scale limit in (4), and it may be interesting to have another way of constructing solutions, which does not involve a homogenization process.

Hence, we introduce a relaxation model of BGK type, in which we pass to the limit as the relaxation parameter goes to infinity. The drawback of this method lies in the fact that the existence of solutions of the limit system is not a consequence of the construction. Indeed, we shall prove that if a solution of the limit system exists, then the family of solutions of the relaxation model strongly converges towards it in the hydrodynamic limit. Hence the proof is not self-contained, because the existence of a solution of the limit system is required in order to pass to the limit. Nevertheless, the final result may be useful in other applications.

In the whole section, the words “limit system” refer to the modified equations introduced in Remark 1. In the divergence-free case, it is also slightly more convenient to work with the function $\chi(\xi, u)$, rather than $\mathbf{1}_{\xi < u}$. Hence a solution of the limit problem is a function g satisfying

$$\sum_{i=1}^N \frac{\partial}{\partial y_i} (a_i(y, \xi)g) = 0, \quad (45)$$

$$\frac{\partial g}{\partial t} + \sum_{i=1}^N a_i(y, \xi) \frac{\partial g}{\partial x_i} = \mathcal{M}, \quad (46)$$

$$\partial_\xi g = \delta(\xi) - \nu(t, x, y, \xi), \quad \nu \geq 0, \quad (47)$$

and \mathcal{M} is such that for all $\varphi \in \mathcal{D}([0, \infty) \times \mathbb{R}^N)$ such that $\varphi \geq 0$, the function $\mathcal{M} *_{t,x} \varphi$ belongs to $\mathcal{C}([0, \infty) \times \mathbb{R}^N, L^2(Y \times \mathbb{R}))$, and

$$\begin{cases} \int_{Y \times \mathbb{R}} (\mathcal{M} *_{t,x} \varphi)(t, x, \cdot) \psi \leq 0, \\ \forall \psi \in L_{\text{loc}}^\infty(Y \times \mathbb{R}), \operatorname{div}_y(a\psi) = 0, \text{ and } \partial_\xi \psi \geq 0. \end{cases} \quad (48)$$

4.1. A relaxation model

The goal of this subsection is to introduce a system approaching (45)-(48). With this aim in view, a relaxation model of BGK type is introduced, which takes into account the constraints of the limit system, that is, equations (45)-(48). Let

$$\begin{aligned} M &:= \|u_0\|_{L^\infty(Y \times \mathbb{R})}, \\ E &:= \{f \in L^2(Y \times \mathbb{R}), \operatorname{Supp} f \subset Y \times [-M, M]\}, \\ \mathbb{K} &:= \{\varphi \in E, \operatorname{div}_y(a(y, \xi)\varphi(y, \xi)) = 0 \text{ in } \mathcal{D}'\}, \\ \mathcal{K} &:= \mathbb{K} \cap \{\varphi \in E, \exists \nu \in M_{\text{per}}^1(Y \times \mathbb{R}), \nu \geq 0, \partial_\xi \varphi = \delta(\xi) - \nu\}. \end{aligned}$$

Then E endowed with the usual scalar product on L^2 is a Hilbert space, and \mathcal{K} is a nonempty closed convex set in E . Thus the projection \mathcal{P} on \mathcal{K} is well-defined.

The main result of this subsection is the following :

Proposition 2 *Let $\lambda, T > 0$ be arbitrary. Set*

$$X_T := \mathcal{C}([0, T], L^2(\mathbb{R}_x^N \times Y \times \mathbb{R}_\xi))$$

and

$$f_0(x, y, \xi) = \chi(\xi, u_0(x, y)), \quad (x, y, \xi) \in \mathbb{R}^N \times Y \times \mathbb{R}.$$

Then there exists a unique solution $f_\lambda \in X_T$ of the equation

$$\begin{cases} \partial_t f_\lambda + a(y, \xi) \cdot \nabla_x f_\lambda + \lambda f_\lambda = \lambda \mathcal{P}(f_\lambda), & t > 0, (x, y, \xi) \in \mathbb{R}^N \times Y \times \mathbb{R}, \\ f_\lambda(t = 0, x, y, \xi) = f_0(x, y, \xi). \end{cases} \quad (49)$$

The function f_λ has the following properties :

1. For almost every t, x, y, ξ ,

$$\begin{aligned} f_\lambda(t, x, y, \xi) &= 0 \text{ if } \xi \geq M, \\ \operatorname{sgn}(\xi) f_\lambda(t, x, y, \xi) &= |f_\lambda(t, x, y, \xi)| \leq 1. \end{aligned}$$

2. L^2 estimate: for all $\lambda > 0$,

$$\|f_\lambda\|_{X_T} \leq \|u_0\|_{L^1(\mathbb{R}^N \times Y)}. \quad (50)$$

3. Strong continuity at time $t = 0$: there exists a function $\omega : [0, \infty) \rightarrow [0, \infty)$, such that $\lim_{0+} \omega = 0$, and such that for all $\lambda > 0, t \geq 0$,

$$\|f_\lambda(t) - f_0\|_{L^1(\mathbb{R}^N \times Y \times \mathbb{R})} \leq \omega(t). \quad (51)$$

4. *Fundamental inequality for $\mathcal{M}_\lambda := \lambda(\mathcal{P}(f_\lambda) - f_\lambda)$: for all $g \in \mathcal{K}$, for almost every (t, x) ,*

$$\int_{Y \times \mathbb{R}} \mathcal{M}_\lambda (\mathcal{P}(f_\lambda) - g) \leq 0. \quad (52)$$

In equation (49), the projection \mathcal{P} acts on the variables y, ξ only; since f is a function of t, x, y, ξ , $\mathcal{P}(f)$ should be understood as

$$\mathcal{P}(f)(t, x, \cdot) = \mathcal{P}(f(t, x, \cdot)),$$

and the above equality holds between functions in $L^2(Y \times \mathbb{R})$, almost everywhere in t, x .

Proof. *First step. Construction of f_λ .* The existence and uniqueness of f_λ follows from a fixed point theorem in X_T . We define the application $\phi_T : X_Y \rightarrow X_T$ by $\phi_T(f) = g$, where g is the solution of the linear equation

$$\begin{cases} \partial_t g + a(y, \xi) \cdot \nabla_x g + \lambda g = \lambda \mathcal{P}(f), \\ g(t = 0, x, y, \xi) = \chi(\xi, u_0(x, y)) \end{cases}$$

The existence and uniqueness of g follow from well-known results on the theory of linear transport equations (recall that $a \in W_{\text{per,loc}}^{1,\infty}(Y \times \mathbb{R})$). Moreover, if $f_1, f_2 \in X_T$ and $g_i = \phi_T(f_i)$, $i = 1, 2$, then $g = g_1 - g_2$ is a solution of

$$\begin{cases} \partial_t g + a(y, \xi) \cdot \nabla_x g + \lambda g = \lambda [\mathcal{P}(f_1) - \mathcal{P}(f_2)], \\ g(t = 0, x, y, \xi) = 0. \end{cases}$$

Multiplying the above equation by g , and integrating on $\mathbb{R}_x^N \times Y \times \mathbb{R}_\xi$, we obtain the estimate

$$\frac{1}{2} \frac{d}{dt} \|g(t)\|_{L^2(\mathbb{R}^N \times Y \times \mathbb{R})}^2 + \lambda \|g(t)\|_{L^2(\mathbb{R}^N \times Y \times \mathbb{R})}^2 \leq \lambda \int_{\mathbb{R}^N \times Y \times \mathbb{R}} [\mathcal{P}(f_1) - \mathcal{P}(f_2)] g.$$

Recall that the projection \mathcal{P} is Lipschitz continuous with Lipschitz constant 1. Thus

$$\begin{aligned} \int_{\mathbb{R}^N \times Y \times \mathbb{R}} [\mathcal{P}(f_1) - \mathcal{P}(f_2)] g &\leq \frac{1}{2} \|\mathcal{P}(f_1(t)) - \mathcal{P}(f_2(t))\|_{L^2(\mathbb{R}^N \times Y \times \mathbb{R})}^2 + \frac{1}{2} \|g(t)\|_{L^2(\mathbb{R}^N \times Y \times \mathbb{R})}^2 \\ &\leq \frac{1}{2} \|(f_1 - f_2)(t)\|_{L^2(\mathbb{R}^N \times Y \times \mathbb{R})}^2 + \frac{1}{2} \|g(t)\|_{L^2(\mathbb{R}^N \times Y \times \mathbb{R})}^2. \end{aligned}$$

Eventually, we obtain

$$\frac{d}{dt} \|g(t)\|_{L^2(\mathbb{R}^N \times Y \times \mathbb{R})}^2 + \lambda \|g(t)\|_{L^2(\mathbb{R}^N \times Y \times \mathbb{R})}^2 \leq \lambda \|(f_1 - f_2)(t)\|_{L^2(\mathbb{R}^N \times Y \times \mathbb{R})}^2 \leq \lambda \|f_1 - f_2\|_{X_T}^2.$$

A straightforward application of Gronwall's lemma yields

$$\|g_1 - g_2\|_{X_T} \leq \sqrt{1 - e^{-\lambda T}} \|f_1 - f_2\|_{X_T}.$$

Thus ϕ_T is a contractant application and has a unique fixed point in X_T , which we call f_λ .

Second step. L^2 estimate. Multiplying (49) by f_λ and integrating on $\mathbb{R}^N \times Y \times \mathbb{R}$, we infer

$$\frac{1}{2} \frac{d}{dt} \|f_\lambda(t)\|_{L^2(\mathbb{R}^N \times Y \times \mathbb{R})}^2 + \lambda \|f_\lambda(t)\|_{L^2(\mathbb{R}^N \times Y \times \mathbb{R})}^2 \leq \lambda \int_{\mathbb{R}^N \times Y \times \mathbb{R}} \mathcal{P}(f_\lambda) f_\lambda.$$

Notice that $0 \in \mathcal{K}$; thus the Lipschitz continuity of \mathcal{P} entails that for almost every t, x

$$\|\mathcal{P}(f_\lambda)(t, x)\|_E \leq \|f_\lambda(t, x)\|_E.$$

Hence, using the Cauchy-Schwartz inequality, we deduce that $t \mapsto \|f_\lambda(t)\|_{L^2(\mathbb{R}^N \times Y \times \mathbb{R})}$ is non-increasing on $[0, T]$. The equality

$$\int_{\mathbb{R}^N \times Y \times \mathbb{R}} |\chi(\xi, u_0(x, y))|^2 dx dy d\xi = \int_{\mathbb{R}^N \times Y \times \mathbb{R}} |\chi(\xi, u_0(x, y))| dx dy d\xi = \int_{\mathbb{R}^N \times Y} |u_0(x, y)| dx dy$$

then yields the desired result.

Third step. Compact support in ξ . Let us prove now that $f_\lambda(\cdot, \xi) = 0$ if $|\xi| > M$: let $\varphi \in \mathcal{D}(\mathbb{R})$ be an arbitrary test function such that $\varphi(\xi) = 0$ when $|\xi| \leq M$. Then $\mathcal{P}(f_\lambda)\varphi = 0$ since $\mathcal{P}(f_\lambda) \in \mathcal{K}$, and thus $f_\lambda\varphi$ is a solution of

$$\begin{aligned} \frac{\partial}{\partial t} (f_\lambda\varphi) + a \cdot \nabla_x (f_\lambda\varphi) + \lambda (f_\lambda\varphi) &= 0, \\ (f_\lambda\varphi)(t = 0, x, y, \xi) &= 0. \end{aligned}$$

Hence $(f_\lambda\varphi)(t, x, y, \xi) = 0$ for almost every t, x, y, ξ , and $f_\lambda(\cdot, \xi) = 0$ if $|\xi| > M$.

Fourth step. Sign property. We now prove the sign property, namely

$$\operatorname{sgn}(\xi)f_\lambda = |f_\lambda| \leq 1 \quad \text{a.e.}$$

This relies on the following fact: if $g \in \mathcal{K}$, then $\operatorname{sgn}(\xi)g(y, \xi) \in [0, 1]$ for almost every y, ξ . Indeed, $g(\cdot, \xi) = 0$ if $\xi < -M$, and thus if $-M < \xi < 0$,

$$g(y, \xi) = - \int_{-M}^{\xi} \nu(y, \xi') d\xi' \leq 0.$$

Hence $g(y, \cdot)$ is non-positive and nonincreasing on $(-\infty, 0)$. Similarly, $g(y, \cdot)$ is non-negative and nondecreasing on $(0, \infty)$. And if $\xi < 0 < \xi'$, then

$$g(y, \xi') - g(y, \xi) = 1 - \int_{\xi}^{\xi'} \nu(y, w) dw \leq 1.$$

Hence the sign property is true for all functions in \mathcal{K} .

Multiplying (49) by $\operatorname{sgn}(\xi)$, we are led to

$$\frac{\partial}{\partial t} (\operatorname{sgn}(\xi)f_\lambda) + a(y, \xi) \cdot \nabla_x (\operatorname{sgn}(\xi)f_\lambda) + \lambda (\operatorname{sgn}(\xi)f_\lambda) = \lambda \mathcal{P}(f_\lambda) \in [0, \lambda].$$

At time $t = 0$, $\operatorname{sgn}(\xi)f_\lambda(t = 0) = |\chi(\xi, u_0)| \in [0, 1]$. Thus, using a maximum principle for the linear transport equation above, we deduce that the sign property is satisfied for f_λ .

Fifth step. Uniform continuity at time $t = 0$. Let $\delta > 0$ be arbitrary, and let $f_0^\delta := f_0 *_{x} \theta^\delta$, with θ^δ a standard mollifier. Then $f_0^\delta(x) \in \mathcal{K}$ for all $x \in \mathbb{R}^N$, and thus $f_\lambda - f_0^\delta$ is a solution of the equation

$$\frac{\partial}{\partial t} (f_\lambda - f_0^\delta) + a(y, \xi) \cdot \nabla_x (f_\lambda - f_0^\delta) + \lambda (f_\lambda - f_0^\delta) = \lambda (\mathcal{P}(f_\lambda) - \mathcal{P}(f_0^\delta)) - a(y, \xi) \cdot (f_0 *_{x} \nabla \theta^\delta).$$

Multiply the above equation by $(f_\lambda - f_0^\delta)$ and integrate on $\mathbb{R}^N \times Y \times \mathbb{R}$. Using once more the Lipschitz continuity of the projection \mathcal{P} , we obtain

$$\begin{aligned} \frac{1}{2} \frac{d}{dt} \|f_\lambda - f_0^\delta\|_{L^2(\mathbb{R}^N \times Y \times \mathbb{R})}^2 &\leq \|a\|_{L^\infty(Y \times (-M, M))} \|f_\lambda - f_0^\delta\|_{L^2(\mathbb{R}^N \times Y \times \mathbb{R})} \|f_0\|_{L^2(\mathbb{R}^N \times Y \times \mathbb{R})} \|\nabla \theta^\delta\|_{L^1} \\ \frac{d}{dt} \|f_\lambda - f_0^\delta\|_{L^2(\mathbb{R}^N \times Y \times \mathbb{R})} &\leq \frac{C}{\delta}. \end{aligned}$$

As a consequence, we obtain the following estimate, which holds for all $t > 0$, $\lambda > 0$ and $\delta > 0$

$$\|f_\lambda(t) - f_0^\delta\|_{L^2(\mathbb{R}^N \times Y \times \mathbb{R})} \leq \frac{Ct}{\delta} + \|f_0 - f_0^\delta\|_{L^2(\mathbb{R}^N \times Y \times \mathbb{R})}.$$

Hence the uniform continuity property is true, with

$$\omega(t) := \inf_{\delta > 0} \left(\frac{Ct}{\delta} + 2 \|f_0 - f_0^\delta\|_{L^2(\mathbb{R}^N \times Y \times \mathbb{R})} \right).$$

Sixth step. Inequality for \mathcal{M}_λ . Inequality (52) is merely a particular case of the inequality

$$\langle \mathcal{P}(f) - f, \mathcal{P}(f) - g \rangle_E \leq 0$$

which holds for all $f \in E$, for all $g \in \mathcal{K}$.

4.2. The hydrodynamic limit

In this subsection, we prove the following result :

Proposition 3 *Let $(f_\lambda)_{\lambda>0}$ be the family of solutions of the relaxation model (49), and let $f(t) = \chi(\xi, u)$ be the unique solution of the limit system (45)-(48) with initial data $\chi(\xi, u_0(x, y))$. Then as $\lambda \rightarrow \infty$,*

$$f_\lambda \rightarrow f \quad \text{in } L^2((0, T) \times \mathbb{R}^N \times Y \times \mathbb{R}).$$

The above Proposition relies on an inequality of the type

$$\frac{d}{dt} \int_{\mathbb{R}^N \times Y \times \mathbb{R}} |f_\lambda - f|^2 \leq r_\lambda(t),$$

with $r_\lambda(t) \rightarrow 0$ as $\lambda \rightarrow \infty$. The calculations are very similar to those of the contraction principle in the previous section; the only difference lies in the fact that f_λ and f are not solutions of the same equation.

Before tackling the proof itself, let us derive a few properties on the weak limit of the sequence f_λ . Since the sequence f_λ is bounded in $X_T \subset L^2((0, T) \times \mathbb{R}^N \times Y \times \mathbb{R})$, there exists a subsequence, which we relabel f_λ , and a function $g \in L^2((0, T) \times \mathbb{R}^N \times Y \times \mathbb{R})$ such that f_λ weakly converges to g in L^2 . Moreover, the sequence $\mathcal{P}(f_\lambda)$ is bounded in $L^2((0, T) \times \mathbb{R}^N \times Y \times \mathbb{R})$, for all $T > 0$. Hence, extracting a further subsequence if necessary, there exists a function $h \in L^2((0, T) \times \mathbb{R}^N \times Y \times \mathbb{R})$ such that $\mathcal{P}(f_\lambda)$ weakly converges towards h as $\lambda \rightarrow \infty$. Notice that the convex set \mathcal{K} is closed for the weak topology in L^2 . Consequently, $h(t, x) \in \mathcal{K}$ for almost every t, x . And we have

$$\mathcal{P}(f_\lambda) - f_\lambda = \mathcal{O}\left(\frac{1}{\lambda}\right),$$

where the \mathcal{O} is meant in the sense of distributions. Hence, $g = h$, and in particular, we deduce that $g(t, x) \in \mathcal{K}$ for almost every (t, x) .

We are now ready to prove the contraction inequality; consider a mollifying sequence θ^δ as in the previous section, and set $f^\delta = f *_{t,x} \theta^\delta$, $f_\lambda^{\delta'} = f_\lambda *_{t,x} \theta^{\delta'}$. Then

$$\begin{aligned} \partial_t f^\delta + a(y, \xi) \cdot \nabla_x f^\delta &= \mathcal{M}^\delta, \\ \partial_t f_\lambda^{\delta'} + a(y, \xi) \cdot \nabla_x f_\lambda^{\delta'} &= \mathcal{M}_\lambda^{\delta'}. \end{aligned}$$

Let us multiply the first equation by $\text{sgn}(\xi) - 2f_\lambda^{\delta'}$, the second by $2(f_\lambda^{\delta'} - f^\delta)$, and add the two identities thus obtained; setting $F_\lambda^{\delta, \delta'} = \text{sgn}(\xi) f^\delta + |f_\lambda^{\delta'}|^2 - 2f^\delta f_\lambda^{\delta'}$, we have

$$\partial_t F_\lambda^{\delta, \delta'} + a(y, \xi) \cdot \nabla_x F_\lambda^{\delta, \delta'} = \mathcal{M}^\delta \left(\text{sgn}(\xi) - 2f_\lambda^{\delta'} \right) + 2\mathcal{M}_\lambda^{\delta'} (f_\lambda^{\delta'} - f^\delta).$$

We integrate over $(0, t) \times \mathbb{R}^N \times Y \times \mathbb{R}$ and obtain

$$\begin{aligned} \int_{\mathbb{R}^N \times Y \times \mathbb{R}} F_\lambda^{\delta, \delta'}(t, x, y, \xi) dx dy d\xi &= \int_0^t \int_{\mathbb{R}^N \times Y \times \mathbb{R}} \mathcal{M}^\delta \left(\text{sgn}(\xi) - 2f_\lambda^{\delta'} \right) \\ &\quad + 2 \int_0^t \int_{\mathbb{R}^N \times Y \times \mathbb{R}} \mathcal{M}_\lambda^{\delta'} (f_\lambda^{\delta'} - f^\delta) \\ &\quad + \int_{\mathbb{R}^N \times Y \times \mathbb{R}} F_\lambda^{\delta, \delta'}(t=0, x, y, \xi) dx dy d\xi. \end{aligned}$$

We now pass to the limit as $\delta' \rightarrow 0$, with all the other parameters fixed. Notice that

$$\begin{aligned} \lim_{\delta' \rightarrow 0} \int_0^t \int_{\mathbb{R}^N \times Y \times \mathbb{R}} \mathcal{M}_\lambda^{\delta'} (f_\lambda^{\delta'} - f^\delta) &= \int_0^t \int_{\mathbb{R}^N \times Y \times \mathbb{R}} \mathcal{M}_\lambda (f_\lambda - f^\delta) \\ &= -\lambda \int_0^t \int_{\mathbb{R}^N \times Y \times \mathbb{R}} (f_\lambda - \mathcal{P}(f_\lambda))^2 \\ &\quad + \int_0^t \int_{\mathbb{R}^N \times Y \times \mathbb{R}} \mathcal{M}_\lambda (\mathcal{P}(f_\lambda) - f^\delta) \\ &\leq 0, \end{aligned}$$

since $f^\delta(t, x) \in \mathcal{K}$ for all t, x . The passage to the limit in $F_\lambda^{\delta, \delta'}(t = 0)$ does not rise any difficulty because of the strong continuity of the functions f_λ at time $t = 0$. Hence, we retrieve

$$\begin{aligned} & \int_{\mathbb{R}^N \times Y \times \mathbb{R}} \left\{ \left(|f^\delta(t)| - |f^\delta(t)|^2 \right) + |f^\delta(t) - f_\lambda(t)|^2 \right\} \\ & \leq \int_0^t \int_{\mathbb{R}^N \times Y \times \mathbb{R}} \mathcal{M}^\delta(\operatorname{sgn}(\xi) - 2f_\lambda) \\ & \quad + \int_{\mathbb{R}^N \times Y \times \mathbb{R}} \left\{ \left(|f^\delta(t=0)| - |f^\delta(t=0)|^2 \right) + |f^\delta(t=0) - \chi(\xi, u_0)|^2 \right\}, \end{aligned}$$

and thus, integrating once again this inequality for $t \in [0, T]$,

$$\begin{aligned} & \int_0^T \int_{\mathbb{R}^N \times Y \times \mathbb{R}} \left\{ \left(|f^\delta| - |f^\delta|^2 \right) + |f^\delta(t) - f_\lambda|^2 \right\} \\ & \leq \int_0^T dt \left[\int_0^t \int_{\mathbb{R}^N \times Y \times \mathbb{R}} \mathcal{M}^\delta(s) (\operatorname{sgn}(\xi) - 2f_\lambda(s)) ds \right] \\ & \quad + T \int_{\mathbb{R}^N \times Y \times \mathbb{R}} \left\{ \left(|f^\delta(t=0)| - |f^\delta(t=0)|^2 \right) + |f^\delta(t=0) - \chi(\xi, u_0)|^2 \right\}. \end{aligned}$$

We now pass to the limit as $\lambda \rightarrow \infty$, with $\delta > 0$ fixed. Then

$$\liminf_{\lambda \rightarrow \infty} \|f_\lambda - f^\delta\|_{L^2((0, T) \times \mathbb{R}^N \times Y \times \mathbb{R})}^2 \geq \|g - f^\delta\|_{L^2((0, T) \times \mathbb{R}^N \times Y \times \mathbb{R})}^2,$$

and

$$\begin{aligned} & \lim_{\lambda \rightarrow \infty} \int_0^T dt \left[\int_0^t \int_{\mathbb{R}^N \times Y \times \mathbb{R}} \mathcal{M}^\delta(s) (\operatorname{sgn}(\xi) - 2f_\lambda(s)) ds \right] \\ & = \int_0^T dt \left[\int_0^t \int_{\mathbb{R}^N \times Y \times \mathbb{R}} \mathcal{M}^\delta(s) (\operatorname{sgn}(\xi) - 2g(s)) ds \right] \leq 0. \end{aligned}$$

Thus, we obtain, for all $\delta > 0$

$$\|g - f^\delta\|_{L^2((0, T) \times \mathbb{R}^N \times Y \times \mathbb{R})}^2 \leq T \int_{\mathbb{R}^N \times Y \times \mathbb{R}} \left\{ |f^\delta(t=0)| - |f^\delta(t=0)|^2 + |f^\delta(t=0) - \chi(\xi, u_0)|^2 \right\}.$$

We have already proved in the previous section that the family $f^\delta(t = 0)$ strongly converges towards $\chi(\xi, u_0)$ as δ vanishes, due to the continuity assumption at time $t = 0$. Hence, we obtain in the limit

$$\|g - f\|_{L^2((0, T) \times \mathbb{R}^N \times Y \times \mathbb{R})}^2 \leq 0,$$

and consequently, $g = f$. Hence the result is proved.

5. The separate case : identification of the limit problem

This section is devoted to the proof of Proposition 1. Thus we focus on the limit system in the case where the flux A can be written as

$$A(y, \xi) = a_0(y)g(\xi), \quad \text{with } \operatorname{div}_y a_0 = 0.$$

The interest of this case lies in the special structure of the limit system; indeed, we shall prove that the function u , which is the two-scale limit of the sequence u^ε , is the solution of the scalar conservation law (20). In view of Theorem 1, we wish to emphasize that Proposition 1 implies in particular that the entropy solution of (20) satisfies the constraint equation

$$\operatorname{div}_y (a_0(y)g(u(t, x; y))) = 0$$

for almost every $t > 0, x \in \mathbb{R}^N$; this fact is not completely obvious when $g \neq \operatorname{Id}$. We will prove in the sequel that $u(t, x)$ actually belongs to the constraint space \mathbb{K}_0 for almost every t, x .

Before addressing the proof of Proposition 1, let us mention that the limit problem (20) is not the one which is expected from a vanishing viscosity approach. Precisely, for any given $\delta > 0$, let u_δ^ε be the solution of

$$\partial_t u_\delta^\varepsilon + \operatorname{div}_x A\left(\frac{x}{\varepsilon}, u_\delta^\varepsilon\right) - \varepsilon \delta \Delta_x u_\delta^\varepsilon = 0,$$

with the initial data $u_\delta^\varepsilon(t=0, x) = u_0(x, x/\varepsilon)$. Then for all $\varepsilon > 0$, $u_\delta^\varepsilon \rightarrow u^\varepsilon$ as $\delta \rightarrow 0$; moreover, the behavior of u_δ^ε as $\varepsilon \rightarrow 0$ is known for each $\delta > 0$ (see [5, 6]). In the divergence-free case, for all $\delta > 0$,

$$\lim_{\varepsilon \rightarrow 0} u_\delta^\varepsilon = \bar{u}(t, x) \quad \text{in } L^1_{\text{loc}},$$

where \bar{u} is the entropy solution of

$$\begin{cases} \partial_t \bar{u} + \operatorname{div}_x(\langle a \rangle g(\bar{u})) = 0, & t > 0, x \in \mathbb{R}^N, \\ \bar{u}(t=0, x) = \langle u_0(x, \cdot) \rangle, & x \in \mathbb{R}^N. \end{cases}$$

Hence, it could be expected that the limits $\varepsilon \rightarrow 0$ and $\delta \rightarrow 0$ can be commuted, that is

$$\lim_{\varepsilon \rightarrow 0} \lim_{\delta \rightarrow 0} u_\delta^\varepsilon = \lim_{\delta \rightarrow 0} \lim_{\varepsilon \rightarrow 0} u_\delta^\varepsilon,$$

which would entail

$$\lim_{\varepsilon \rightarrow 0} u^\varepsilon = \bar{u}.$$

In general, this equality is false, even in a weak sense: a generic counter-example is the one of shear flows (see for instance the calculations in [9]). Indeed, if $N = 2$ and $A(y, \xi) = (a_1(y_2)\xi, 0)$, then equation (20) becomes

$$\begin{cases} \partial_t u + a_1(y_2) \partial_{x_1} u = 0, & t > 0, x \in \mathbb{R}^2, y \in [0, 1]^2 \\ u(t=0, x, y) = u_0(x_1, x_2, y_2), & x \in \mathbb{R}^2, y \in [0, 1]^2. \end{cases}$$

It is then easily checked that in general, the average of u over $Y = [0, 1]^2$ is not the solution of the transport equation

$$\partial_t \bar{u} + \langle a_1 \rangle \partial_{x_1} \bar{u} = 0.$$

We now turn to the proof of Proposition 1. In view of Theorem 1, it is sufficient to prove that the entropy solution of (20) belongs to \mathbb{K}_0 for a.e. t, x , or in other words, that \mathbb{K}_0 is invariant by the semi-group associated to equation (20). We prove this result in the slightly more general context of kinetic solutions. The core of the proof lies in the following Proposition:

Proposition 4 *Let $u_0 \in L^1(\mathbb{R}^N, L^\infty(Y))$ such that $u_0(x, \cdot) \in \mathbb{K}_0$ for almost every $x \in \mathbb{R}^N$.*

Let $v = v(t, x; y) \in \mathcal{C}([0, \infty); L^1(\mathbb{R}^N \times Y))$ be the kinetic solution of

$$\begin{cases} \partial_t v(t, x; y) + \operatorname{div}_x(\tilde{a}_0(y)g(v(t, x; y))) = 0, & t > 0, x \in \mathbb{R}^N, y \in Y, \\ v(t=0, x; y) = u_0(x, y). \end{cases} \quad (53)$$

Then for almost every $t > 0, x \in \mathbb{R}^N$, $v(t, x) \in \mathbb{K}_0$.

Remark 7 *Let us recall that the function v is the kinetic solution of (53) if there exists a non-negative measure $m \in \mathcal{C}(\mathbb{R}_\xi, M_w^1([0, \infty) \times \mathbb{R}^N \times Y))$ such that the following properties hold:*

– *The function $f^1(t, x, y, \xi) := \chi(\xi, v(t, x; y))$ is a solution in the sense of distributions of*

$$\begin{cases} \partial_t f^1 + \tilde{a}_0(y) \cdot \nabla_x f^1 g'(\xi) = \partial_\xi m, & t > 0, x \in \mathbb{R}^N, y \in Y, \xi \in \mathbb{R}, \\ f^1(t=0, x, y, \xi) = \chi(\xi, u_0(x, y)); \end{cases} \quad (54)$$

– *There exists a function $\mu \in L^\infty(\mathbb{R})$ such that*

$$\begin{aligned} \int_0^\infty \int_{\mathbb{R} \times Y} m(t, x, y, \xi) dx dy dt &\leq \mu(\xi) \quad \forall \xi \in \mathbb{R}, \\ \lim_{|\xi| \rightarrow \infty} \mu(\xi) &= 0. \end{aligned}$$

Proof. First, let us recall (see [23,24]) that for all $T > 0$,

$$f^1 = \lim_{\lambda \rightarrow \infty} f_\lambda \quad \text{in } \mathcal{C}([0, T]; L^1(\mathbb{R}^N \times Y \times \mathbb{R})),$$

where $f_\lambda = f_\lambda(t, x, y, \xi)$ ($\lambda > 0$) is the unique solution of the system

$$\begin{cases} \partial_t f_\lambda + \tilde{a}_0(y) \cdot \nabla_x f_\lambda g'(\xi) + \lambda f_\lambda = \lambda \chi(\xi, u_\lambda), \\ u_\lambda(t, x, y) = \int_{\mathbb{R}} f_\lambda(t, x, y, \xi) d\xi, \\ f_\lambda(t=0) = \chi(\xi, u_0). \end{cases} \quad (55)$$

Moreover, for every $\lambda > 0$, u_λ is the unique fixed point of the contractant application

$$\phi_\lambda : \mathcal{C}((0, T); L^1(\mathbb{R}^N \times Y)) \rightarrow \mathcal{C}((0, T); L^1(\mathbb{R}^N \times Y)) \\ u_1 \mapsto u_2$$

where $u_2 = \int_\xi f$ and f is the solution of

$$\begin{cases} \partial_t f + \tilde{a}_0(y) \cdot \nabla_x f g'(\xi) + \lambda f = \lambda \chi(\xi, u_1), \\ f(t=0) = \chi(\xi, u_0). \end{cases} \quad (56)$$

Thus, it is sufficient to prove that the space

$$\mathcal{I}_0 := \{u \in \mathcal{C}([0, T]; L^1(\mathbb{R}^N \times Y)); u(t, x) \in \mathbb{K}_0 \text{ a.e}\}$$

is invariant by the application ϕ_λ .

First, let us stress that for all $u \in L^1(Y)$,

$$u \in \mathbb{K}_0 \iff \operatorname{div}_y(a(y)\chi(\xi, u)) = 0 \text{ in } \mathcal{D}'(Y \times \mathbb{R}). \quad (57)$$

Indeed, if $u \in \mathbb{K}_0$, then for all $\delta > 0$, set $u_\delta = u * \theta^\delta$, with θ^δ a standard mollifier. The function u_δ is a solution of

$$\operatorname{div}_y(a_0 u_\delta) = r_\delta,$$

and the remainder r_δ vanishes strongly in $L^1(Y)$ (see the calculations in the previous sections). Since the function u_δ is smooth, if $G \in \mathcal{C}^1(\mathbb{R}^N)$, we have

$$\operatorname{div}_y(a_0 G(u_\delta)) = G'(u_\delta) r_\delta.$$

Passing to the limit as δ vanishes, we infer $\operatorname{div}_y(a_0 G(u)) = 0$ for all $G \in \mathcal{C}^1(\mathbb{R}^N)$. At last, taking a sequence of smooth functions approaching $\chi(\xi, u)$, we deduce that $\operatorname{div}_y(a_0 \chi(\xi, u)) = 0$ in $\mathcal{D}'_{\text{per}}(Y \times \mathbb{R})$. Conversely, assume that $\operatorname{div}_y(a_0 \chi(\xi, u)) = 0$; integrating this equation with respect to ξ yields $u \in \mathbb{K}_0$. Hence (57) is proved.

Now, let us prove that the space \mathcal{I}_0 is invariant by the application ϕ_λ . Let $u_1 \in \mathcal{C}([0, T]; L^1(\mathbb{R}^N \times Y))$ such that $u_1(t, x) \in \mathbb{K}_0$ almost everywhere. Then

$$\operatorname{div}_y(a_0(y)\chi(\xi, u_1(t, x, y))) = 0.$$

Let f be the solution of (56); since $\tilde{a}_0 \in \mathbb{K}_0^N$, the distribution $\operatorname{div}_y(a_0 f)$ satisfies the transport equation

$$\partial_t (\operatorname{div}(a_0 f)) + g'(\xi) \tilde{a}_0(y) \cdot \nabla_x (\operatorname{div}(a_0 f)) + \lambda \operatorname{div}(a_0 f) = 0, \quad t > 0, \quad x \in \mathbb{R}^N, \quad y \in Y, \quad \xi \in \mathbb{R},$$

and $\operatorname{div}(a_0 f)(t=0) = 0$ because $u_0(x) \in \mathbb{K}_0$ almost everywhere. Hence

$$\operatorname{div}_y(a_0(y)f(t, x, y, \xi)) = 0 \quad \text{in } \mathcal{D}'.$$

Integrating this equation with respect to ξ yields $u_2 \in \mathbb{K}_0$ almost everywhere.

Consequently, for all $\lambda > 0$, $u_\lambda(t, x; \cdot) \in \mathbb{K}_0$ for almost every t, x . Passing to the limit, we deduce that $v(t, x; \cdot) \in \mathbb{K}_0$ almost everywhere. Hence Proposition 4 is proved.

We now turn to the proof of Proposition 1: setting $b(y) = a_0(y) - \tilde{a}_0(y)$, we have

$$\partial_t f^1 + a_0(y) \nabla_x f^1 g'(\xi) = \partial_\xi m - b(y) \nabla_x f^1 g'(\xi) =: \mathcal{M}_1.$$

If $u_0 \in L^\infty(\mathbb{R}^N)$, then $v \in L^\infty([0, \infty) \times \mathbb{R}^N \times Y)$, and it is easily checked that f^1 and \mathcal{M}_1 satisfy the compact support assumptions. According to Proposition 4, f^1 also satisfies (45), and thanks to the structure of the right-hand side, the distribution \mathcal{M}_1 satisfies (48). Thus f^1 is the unique solution of the limit system, and Proposition 1 is proved.

6. Further remarks on the notion of limit system

Here, we have gathered, by way of conclusion, a few remarks on the limit evolution system introduced in Definition 1. The main idea behind this section is that the limit system is not unique (although its solution always is), and thus several other relevant equations can be written instead of (10). Unfortunately, there does not seem to be any rule which would allow to decide between two limit systems.

Let us illustrate these words by a first series of examples: assume that the flux is divergence free, and let

$$\begin{aligned} a'(y, \xi) &= (a_1(y, \xi), \dots, a_N(y, \xi)) \in \mathbb{R}^N, \quad (y, \xi) \in Y \times \mathbb{R}, \\ \mathbb{K} &:= \{f \in L^1_{\text{loc}}(Y \times \mathbb{R}), \operatorname{div}_y(a'f) = 0 \text{ in } \mathcal{D}'\}. \end{aligned}$$

We denote by P the projection on \mathbb{K} in $L^1_{\text{loc}}(Y \times \mathbb{R})$. Precisely, consider the dynamical system $X(t, y; \xi)$ defined by

$$\begin{cases} \dot{X}(t, y; \xi) = a'(X(t, y; \xi), \xi), & t > 0, \\ X(0, y; \xi) = y \in Y. \end{cases}$$

Then for all $\xi \in \mathbb{R}$, the Lebesgue measure on Y is invariant by the semi-group $X(t; \xi)$ because $a'(\cdot, \xi)$ is divergence free for all ξ . Hence, by the ergodic theorem, for all $f \in L^1_{\text{loc}}(Y \times \mathbb{R})$ there exists a function $P(f)$ in $L^1_{\text{loc}}(Y \times \mathbb{R})$, such that

$$P(f)(y, \xi) = \lim_{T \rightarrow \infty} \frac{1}{T} \int_0^T f(X(t, y; \xi), \xi) dt,$$

and the limit holds a.e. in y, ξ and in $Y \times (-R, R)$ for all $R > 0$.

For $1 \leq i \leq N$, set $\tilde{a}_i := P(a_i)$. Then if f is a solution of the limit system, f also satisfies

$$\partial_t f + \tilde{a}(y, \xi) \cdot \nabla_x f = \tilde{\mathcal{M}}$$

and $f, \tilde{\mathcal{M}}$ satisfy (9) and (11) - (14). Indeed,

$$\tilde{\mathcal{M}} = \mathcal{M} + [\tilde{a}(y, \xi) - a'(y, \xi)] \cdot \nabla_x f$$

and the term $[\tilde{a}(y, \xi) - a'(y, \xi)] \cdot \nabla_x (f *_x \varphi)(t, x, y, \xi)$ belongs to \mathbb{K}^\perp for all t, x and for all $\varphi \in \mathcal{D}([0, \infty) \times \mathbb{R}^N)$. Of course, as long as \tilde{a} is sufficiently smooth, say $\tilde{a} \in W^{1, \infty}_{\text{loc}}(\mathbb{R}^{N+1})$, uniqueness holds for this limit system (the proof is exactly the same as the one in section 3). Thus this constitutes as legitimate a limit system as the one in Definition 1. In fact, in the separate case, Proposition 1 indicates that the above system seems to be the relevant one, rather than the one in Definition 1. Notice that the distribution $\tilde{\mathcal{M}}$ satisfies the additional property

$$\tilde{\mathcal{M}} *_t *_x \phi(t, x) \in \mathbb{K}^\perp \quad \forall t, x.$$

Let us now go a little further: let $\theta \in \mathcal{C}^1(\mathbb{R})$ such that $0 \leq \theta \leq 1$, and let

$$a_\theta(y, \xi) = \theta(\xi)a'(y, \xi) + (1 - \theta(\xi))\tilde{a}(y, \xi).$$

Then f is a solution of

$$\partial_t f + a_\theta(y, \xi) \cdot \nabla_x f = \mathcal{M}_\theta,$$

for some distribution \mathcal{M}_θ satisfying (14). Thus this constitutes yet another limit system which has the same structure as the one of Definition 1. Hence the limit system is highly non unique, and it must be seen as a way of identifying the two-scale limit of the sequence f^ε , rather than as a kinetic formulation of a given conservation law, for instance. Let us also emphasize the following fact: consider the solution $v = v(t, x, y)$ of the scalar conservation law

$$\begin{cases} \partial_t v + \operatorname{div}_x \tilde{A}(y, v) = 0, \\ v(t = 0, x, y) = u_0(x, y), \end{cases}$$

where the flux \tilde{A} is such that $\partial_\xi \tilde{A}_i(y, \xi) = \tilde{a}_i(y, \xi)$. Then, in general, the function $\mathbf{1}_{\xi < v}$ is not a solution of the limit system, except in the so-called ‘‘separate case’’ described in Proposition 1.

Indeed, the function v is not a solution of the cell problem in general, even if u_0 is. In other words, the set \mathbb{K} is not invariant by the evolution equation

$$\partial_t g + \sum_{i=1}^N \tilde{a}_i(y, \xi) \partial_{x_i} g = \partial_\xi m,$$

where m is a nonnegative measure and $g = \mathbf{1}_{\xi < v}$.

Let us now assume that the flux A is not divergence free. Then there are cases where a different notion of limit problem can be given: assume that there exists real numbers $p_1 < p_2$, and a family $\{v(\cdot, p)\}_{p_1 \leq p \leq p_2}$, which satisfies the following properties:

1. The function $(y, p) \mapsto v(y, p)$ belongs to $L^\infty(Y \times [p_1, p_2])$;
2. For all $p \in [p_1, p_2]$, $v(\cdot, p)$ is an entropy solution of the cell problem; in other words, there exists a nonnegative measure $m(y, \xi; p)$ such that $f(y, \xi; p) = \mathbf{1}_{\xi < v(y, p)}$ is a solution of

$$\sum_{i=1}^N \frac{\partial}{\partial y_i} (a_i(y, \xi) f) + \frac{\partial}{\partial \xi} (a_{N+1}(y, \xi) f) = \frac{\partial}{\partial \xi} m;$$

3. For all $p \in [p_1, p_2]$, $\langle v(\cdot, p) \rangle_Y = 0$;
4. The distribution $\partial_p v$ is a nonnegative function in $L^1(Y \times [p_1, p_2])$; this implies in particular that for all couples $(p, p') \in [p_1, p_2]^2$ such that $p \geq p'$, for almost every $y \in Y$,

$$v(y, p) \geq v(y, p').$$

Under these conditions, a kinetic formulation for equation (1) can be derived, based on the family $v(x/\varepsilon, p)$ of stationary solutions of (1), rather than on the family of Kruzkov's inequalities. This kind of construction was achieved in [7] in a parabolic setting, following an idea developed by Emmanuel Audusse and Benoît Perthame in [3]; these authors define a new notion of entropy solutions for a heterogeneous conservation law in dimension one, based on the comparison with a family of stationary solutions. Let us explain briefly how the kinetic formulation for entropy solutions of (1) is derived: let u^ε be an entropy solution of (1). Define the distribution $m^\varepsilon \in \mathcal{D}'((0, \infty) \times \mathbb{R}^N \times (p_1, p_2))$ by

$$m^\varepsilon(t, x, p) := - \left\{ \frac{\partial}{\partial t} \left(u^\varepsilon - v \left(\frac{x}{\varepsilon}, p \right) \right)_+ + \frac{\partial}{\partial y_i} \left[\mathbf{1}_{v(\frac{x}{\varepsilon}, p) < u^\varepsilon} \left(A_i \left(\frac{x}{\varepsilon}, u^\varepsilon \right) - A_i \left(\frac{x}{\varepsilon}, v \left(\frac{x}{\varepsilon}, p \right) \right) \right) \right] \right\}. \quad (58)$$

Then according to the comparison principle (which was known by Kruzkhov, see [16,17]), m^ε is a nonnegative measure on $(0, \infty) \times \mathbb{R}^N \times [p_1, p_2]$. Now, set

$$f^\varepsilon(t, x, p) := \mathbf{1}_{v(\frac{x}{\varepsilon}, p) < u^\varepsilon(t, x)} \in L^\infty([0, \infty) \times \mathbb{R}^N \times [p_1, p_2]).$$

Thanks to the regularity assumptions on the family $v(\cdot, p)$, we can differentiate equality (58) (which is meant in the sense of distributions) with respect to p , and we are led to

$$\frac{\partial}{\partial t} \left(f^\varepsilon v_p \left(\frac{x}{\varepsilon}, p \right) \right) + \frac{\partial}{\partial x_i} \left(f^\varepsilon v_p \left(\frac{x}{\varepsilon}, p \right) a_i \left(\frac{x}{\varepsilon}, v \left(\frac{x}{\varepsilon}, p \right) \right) \right) = \frac{\partial m^\varepsilon}{\partial p}. \quad (59)$$

This equation is in fact the appropriate kinetic formulation in the heterogeneous case; its main advantage over equation (4) is the absence of the highly oscillating term

$$\frac{1}{\varepsilon} \partial_\xi \left[a_{N+1} \left(\frac{x}{\varepsilon}, \xi \right) \mathbf{1}_{\xi < u^\varepsilon} \right].$$

Notice that for all $p \in [p_1, p_2]$,

$$\operatorname{div}_y \left(\frac{\partial v(y, p)}{\partial p} a(y, v(y, p)) \right) = 0 \quad \text{in } \mathcal{D}'_{\text{per}}(Y). \quad (60)$$

This equation is derived by differentiating equation

$$\operatorname{div}_y A(y, v(y, p)) = 0$$

with respect to p . Thus, if we set

$$\check{a}(y, p) := \frac{\partial v(y, p)}{\partial p} a(y, v(y, p)),$$

the vector field $\check{a} \in L^1(Y \times [p_1, p_2])$ is divergence-free, and the same kind of limit system as in the divergence free case can be built. Of course, the interest of such a construction lies in the simplicity of the structure of the limit system in the divergence free case.

Definition 3 Let $f \in L^\infty([0, \infty), L^1(\mathbb{R}^N \times Y \times \mathbb{R}))$, $u_0 \in L^1 \cap L^\infty(\mathbb{R}^N \times Y)$. It is said that f is a generalized kinetic solution of the limit problem associated with the family $v(\cdot, p)$ if there exists a distribution $\mathcal{M} \in \mathcal{D}'_{per}([0, \infty) \times \mathbb{R}^N \times Y \times \mathbb{R})$ such that f and \mathcal{M} satisfy the following properties:

1. Compact support in p : there exists $(p'_1, p'_2) \in [p_1, p_2]^2$ such that $p_1 < p'_1 \leq p'_2 < p_2$, and

$$\begin{aligned} \text{Supp} \mathcal{M} &\subset [0, \infty) \times \mathbb{R}^N \times Y \times [p'_1, p'_2]; \\ f(t, x, y, p) &= 1 \text{ if } p_1 < p < p'_1, \quad f(t, x, y, p) = 0 \text{ if } p'_2 < p < p_2. \end{aligned}$$

2. Microscopic equation for f : f is a solution in the sense of distributions on $Y \times (p_1, p_2)$ of

$$\text{div}_y(\check{a}(y, p)f(t, x, y, p)) = 0. \quad (61)$$

3. Evolution equation: the couple (f, \mathcal{M}) is a solution in the sense of distributions on $[0, \infty) \times \mathbb{R}^N \times Y \times (p_1, p_2)$ of

$$\begin{cases} \partial_t(v_p(y, p)f) + \check{a}(y, p) \cdot \nabla_x f = \mathcal{M}, \\ f(t=0, x, y, p) = \mathbf{1}_{v(y, p) < u_0(x, y)} =: f_0(x, y, p); \end{cases} \quad (62)$$

In other words, for any test function $\phi \in \mathcal{D}_{per}([0, \infty) \times \mathbb{R}^N \times Y \times (p_1, p_2))$,

$$\begin{aligned} \int_0^\infty \int_{\mathbb{R}^N \times Y \times \mathbb{R}} f(t, x, y, p) v_p(y, p) \{ \partial_t \phi(t, x, y, p) + a(y, v(y, p)) \cdot \nabla_x \phi(t, x, y, p) \} dt dx dy d\xi = \\ = - \langle \phi, \mathcal{M} \rangle_{\mathcal{D}, \mathcal{D}'} - \int_{\mathbb{R}^N \times Y \times \mathbb{R}} \mathbf{1}_{v(y, p) < u_0(x, y)} v_p(y, p) \phi(t=0, x, y, p) dx dy d\xi. \end{aligned}$$

4. Conditions on f :

$$\partial_p f \leq 0, \quad (63)$$

$$0 \leq f(t, x, y, \xi) \leq 1 \quad a.e., \quad (64)$$

$$\frac{1}{\tau} \int_0^\tau \|f(s) - f_0\|_{L^2(\mathbb{R}^N \times Y \times (p_1, p_2))} ds \xrightarrow{\tau \rightarrow 0} 0. \quad (65)$$

5. Condition on \mathcal{M} : for all $\varphi \in \mathcal{D}([0, \infty) \times \mathbb{R}^N)$ such that $\varphi \leq 0$, the function $\mathcal{M} *_{t,x} \varphi$ belongs to $\mathcal{C}([0, \infty) \times \mathbb{R}^N, L^1(Y \times \mathbb{R}))$, and

$$\begin{cases} \int_{Y \times \mathbb{R}} (\mathcal{M} *_{t,x} \varphi)(t, x, \cdot) \psi \leq 0, \\ \forall \psi \in L^\infty_{loc}(Y \times \mathbb{R}), \text{div}_y(\check{a}\psi) = 0, \text{ and } \partial_\xi \psi \geq 0. \end{cases} \quad (66)$$

We now state without proof a result analogue to Theorems 1, 2 :

Proposition 5 Let $A \in W^{2, \infty}_{per, loc}(Y \times \mathbb{R})$. Assume that $a \in \mathcal{C}^1_{per}(Y \times \mathbb{R})$ and that $\check{a} \in W^{1, 1}(Y \times (p_1, p_2))$. Let $u_0 \in L^\infty(\mathbb{R}^N \times Y) \cap L^1_{loc}(\mathbb{R}^N, \mathcal{C}_{per}(Y))$ such that $u_0(x, \cdot)$ is an entropy solution of the cell problem for almost every $x \in \mathbb{R}^N$. Assume furthermore that there exists $p'_1 < p'_2$ in $(p_1, p_2)^2$ such that

$$v(y, p'_1) \leq u_0(x, y) \leq v(y, p'_2),$$

and let

$$f_0(x, y, p) := \mathbf{1}_{v(y, p) < u_0(x, y)}$$

Then the following results hold :

1. There exists a unique generalized kinetic solution f of the limit problem associated with the family $(v(\cdot, p))_{p_1 \leq p \leq p_2}$ with initial data f_0 . Moreover, there exists a function $u \in L^\infty([0, \infty) \times \mathbb{R}^N \times Y)$ such that

$$f(t, x, y, p) = \mathbf{1}_{v(y, p) < u(t, x, y)} \quad a.e.$$

2. Let $u^\varepsilon \in L^\infty([0, \infty) \times \mathbb{R}^N)$ be the entropy solution of (1) with initial data $u_0(x, x/\varepsilon)$. Let $f(t, x, y, p) = \mathbf{1}_{v(y, p) < u(t, x, y)}$ be the unique solution of the limit problem. Then for all regularization kernels φ^δ of the form

$$\varphi^\delta(x) = \frac{1}{\delta^N} \varphi\left(\frac{x}{\delta}\right), \quad x \in \mathbb{R}^N,$$

with $\varphi \in \mathcal{D}(\mathbb{R}^N)$, $\int \varphi = 1$, $0 \leq \varphi \leq 1$, we have, for all compact $K \subset [0, \infty) \times \mathbb{R}^N$,

$$\lim_{\delta \rightarrow 0} \lim_{\varepsilon \rightarrow 0} \left\| u^\varepsilon(t, x) - u *_x \varphi^\delta\left(t, x, \frac{x}{\varepsilon}\right) \right\|_{L^1(K)} = 0. \quad (67)$$

Hence a whole variety of limit systems can be given, depending on the choice of the family of solutions of the cell problem. However, it is not obvious that any given system is “better” than another one. But the important result, as far as homogenization is concerned, is that all systems have a unique solution.

References

1. GRÉGOIRE ALLAIRE: Homogenization and two-scale convergence. *SIAM J. Math. Anal.* **23**, 1482–1518 (1992).
2. DEBORA AMADORI AND DENIS SERRE: Asymptotic behavior of solutions to conservation laws with periodic forcing. *J. Hyperbolic Differential Equations* **3**, 387–401 (2006).
3. EMMANUEL AUDUSSE AND BENOÎT PERTHAME: Uniqueness for scalar conservation laws with discontinuous flux via adapted entropies. *Proc. of the Royal Soc. of Edinburgh* **135A**, 253–266 (2005).
4. CONSTANTINE M. DAFERMOS: *Hyperbolic conservation laws in Continuum Physics*. Grundlehren Mathematischen Wissenschaften, no. GM 325, Springer-Verlag, Berlin, Heidelberg, New York, 205, second edition.
5. A.-L. DALIBARD: Homogenization of a quasilinear parabolic equation with vanishing viscosity. *J. Math. Pures et Appl.* **86**, 133–154 (2006).
6. A.-L. DALIBARD: Initial layer for the homogenization of a quasilinear parabolic equation with vanishing viscosity. *Arch. Rat. Mech. Anal.* **185**, 515–543 (2007).
7. A.-L. DALIBARD: Kinetic formulation for a parabolic conservation law. Application to homogenization. *SIAM J. Math. Anal.* **39**, 891–915 (2007).
8. A.-L. DALIBARD: Kinetic formulation for heterogeneous scalar conservation laws. *Annales de l’IHP (C) : Analyse non linéaire* **23**, 475–498 (2006).
9. WEINAN E: Homogenization of linear and nonlinear transport equations. *Comm. Pure Appl. Math.* **45**, 301–326 (1992).
10. WEINAN E: Homogenization of scalar conservation laws with oscillatory forcing terms. *SIAM J. Appl. Math.* **52**, 959–972 (1992).
11. WEINAN E AND DENIS SERRE: Correctors for the homogenization of conservation laws with oscillatory forcing terms. *Asymptotic Analysis* **5**, 311–316 (1992).
12. BJORN ENGQUIST AND WEINAN E: Large time behavior and homogenization of solutions of two-dimensional conservation laws. *Comm. Pure Appl. Math.* **46**, 1–26 (1993).
13. L. C. EVANS AND D. GOMES: Effective Hamiltonians and averaging for Hamiltonian dynamics. II. *Arch. Ration. Mech. Anal.* **161**, 271–305 (2002).
14. THOMAS Y. HOU AND XUE XIN: Homogenization of linear transport equations with oscillatory vector fields. *SIAM J. Appl. Math.* **52**, 34–45 (1992).
15. P. E. JABIN AND A. TZAVARAS: Kinetic decomposition of homogenization problems. Preprint (2006).
16. S. N. N. KRUIZKOV: Generalized solutions of the Cauchy problem in the large for nonlinear equations of first order. *Soviet Math. Dokl.* **10**, 785–788 (1969).
17. S. N. N. KRUIZKOV: First order quasilinear equations in several independent variables. *Math. USSR Sb.* **10**, 217–243 (1970).
18. P.-L. LIONS, G. PAPANICOLAOU, AND S. R. S. VARADHAN: Homogenization of Hamilton-Jacobi equations. Unpublished (1987).
19. PIERRE-LOUIS LIONS, BENOÎT PERTHAME, AND EITAN TADMOR: Formulation cinétique des lois de conservation scalaires multidimensionnelles. *C.R. Acad. Sci. Paris, Série I*, **312**, 97–102 (1991).
20. PIERRE-LOUIS LIONS, BENOÎT PERTHAME, AND EITAN TADMOR: A kinetic formulation of multidimensional conservation laws and related equations. *J. Amer. Math. Soc.* **7**, 169–191 (1994).

21. GABRIEL N'GUETSENG: A general convergence result for a functional related to the theory of homogenization. *SIAM J. Math. Anal.* **20**, 608–623 (1989).
22. BENOÎT PERTHAME: Uniqueness and error estimates in first order quasilinear conservation laws via the kinetic entropy defect measure. *J. Math. Pures et Appl.* **77**, 1055–1064 (1998).
23. BENOÎT PERTHAME: *Kinetic formulation of conservation laws*. Oxford Lecture Series in Mathematics and its Applications, no. 21, Oxford University Press, 2002.
24. BENOÎT PERTHAME AND EITAN TADMOR: A kinetic equation with kinetic entropy functions for scalar conservation laws. *Comm. Math. Phys.* **136**, 501–517 (1991).
25. D. SERRE: *L^1 -stability of nonlinear waves in scalar conservation laws*. Evolutionary equations (C.M.Dafermos and E.Feireisl, eds.), Handb. Differ. Equ., vol. 1, North-Holland, 2004, pp. 473–553.
26. DENIS SERRE: *Systems of conservation laws. 1. Hyperbolicity, entropies, shock waves*. Cambridge University Press, Cambridge, 1999.
27. DENIS SERRE: *Systems of conservation laws. 2. Geometric structures, oscillations, and initial-boundary value problems*. Cambridge University Press, Cambridge, 2000.

CEREMADE-UMR 7534
Université Paris-Dauphine
Place du maréchal de Lattre de Tassigny
75775 Paris Cedex 16, FRANCE
email: dalibard@ceremade.dauphine.fr