

Separation for the stationary Prandtl equation

Anne-Laure Dalibard* Nader Masmoudi†

February 28, 2018

Abstract

In this paper, we prove that separation occurs for the stationary Prandtl equation, in the case of adverse pressure gradient, for a large class of boundary data at $x = 0$. We justify the Goldstein singularity: more precisely, we prove that under suitable assumptions on the boundary data at $x = 0$, there exists $x^* > 0$ such that $\partial_y u_{y=0}(x) \sim C\sqrt{x^* - x}$ as $x \rightarrow x^*$ for some positive constant C , where u is the solution of the stationary Prandtl equation in the domain $\{0 < x < x^*, y > 0\}$. Our proof relies on three main ingredients: the computation of a “stable” approximate solution, using modulation theory arguments; a new formulation of the Prandtl equation, for which we derive energy estimates, relying heavily on the structure of the equation; and maximum principle and comparison principle techniques to handle some of the nonlinear terms.

Contents

1	Introduction	2
1.1	Setting of the problem and state of the art	3
1.2	Main result	5
2	Strategy of proof	7
2.1	Self-similar change of variables	7
2.2	Construction of an approximate solution	9
2.3	Error estimates	13
2.3.1	Commutator of \mathcal{L}_U with $\partial_s + \frac{b}{2}Y\partial_Y$	16
2.3.2	Action of \mathcal{L}_U on the remainder term \mathcal{R}	16
2.3.3	Energy estimates on transport-diffusion equations of the type (2.18)	17
2.4	Construction of sub- and super-solutions	21
2.5	Bootstrap argument	22
2.6	Organization of the rest of the paper	25
3	Derivation of L^∞ estimates and construction of sub and super solutions	26
3.1	Uniform bounds on $\partial_{YY}U$	27
3.2	Construction of sub and super solutions for W	30
3.3	Refined lower bound on $\partial_{YY}U$	33

*Sorbonne Université, Université Paris-Diderot SPC, CNRS, Laboratoire Jacques-Louis Lions, LJLL, F-75005 Paris

†NYUAD campus, Saadiyat Island Abu Dhabi, UAE and Courant Institute, New York University, 251 Mercer Street, New York, NY 10012.

4	Main tools for the energy estimates	33
4.1	Proof of Lemma 2.6	33
4.2	Proof of the commutator result (Lemma 2.8)	38
4.3	Structure of the diffusion term	40
4.4	Structure of the commutator	46
4.5	Useful inequalities	47
4.5.1	Evaluation of some remainder terms	47
4.5.2	Control of integral tails	48
4.5.3	A special case of Hardy inequalities	50
5	Sequence of estimates on V	51
5.1	Preliminary step: estimates on (E_0, D_0)	54
5.2	Estimate on $\partial_Y^2 \mathcal{L}_U V$: proof of Proposition 2.9	56
5.3	Estimate on $\partial_Y^2 \mathcal{L}_U^2 V$: proof of Proposition 2.13	66
A	Useful formulas	76
B	Estimate on the modulation rate	77
C	Proof of Lemma 3.6	79
D	Proof of Lemma 3.7	82
E	Proof of Lemma 4.11	85
F	Proofs of Lemmas 2.14 and 2.15	87

1 Introduction

One of the main open problems in the mathematical analysis of fluid flows is the understanding of the inviscid limit in the presence of boundaries. In the case of a fixed bounded domain, it is an open problem to know whether solutions to the Navier-Stokes system with no slip boundary condition (zero Dirichlet boundary condition) do converge to a solution to the Euler system when the viscosity goes to zero. The main problem here comes from the fact that we cannot impose a no slip boundary condition for the Euler system. To recover a zero Dirichlet condition, Prandtl proposed to introduce a boundary layer [30] in a small neighborhood of the boundary in which viscous effects are still present. It turns out that the system that governs the flow in this small neighborhood, namely the Prandtl system has many mathematical difficulties. One of the outcome is that the justification of the approximation of the Navier-Stokes system by the Euler system in the interior and the Prandtl system in a boundary layer is still mainly open. We refer to Sammartino and Caffisch [31, 32] for this justification in the analytic case. There is also a well known convergence criterion due to Kato [14] that states that the convergence from Navier-Stokes to Euler holds as long as there is no viscous dissipation in a small layer around the boundary (see also [22]).

Let us also mention that when the no slip boundary condition is replaced by a Navier type condition or an inflow condition, the situation gets much better: Bardos [1] proved that the convergence holds for some special type of boundary condition (vorticity equal to zero on the boundary) which does not require the construction of any boundary layer. For Navier

boundary conditions, a boundary layer can be constructed and controlled (see for instance [3, 36, 15, 10, 2, 16, 24, 25]).

We are interested in the present paper in the stationary version of the Prandtl equation, namely

$$\begin{aligned} uu_x + vu_y - u_{yy} &= -\frac{dp_E(x)}{dx}, & x > 0, y > 0, \\ u_x + v_y &= 0, & x > 0, y > 0, \\ u|_{x=0} &= u_0, \quad u|_{y=0} = 0, \quad \lim_{y \rightarrow \infty} u(x, y) = u_E(x), \end{aligned} \tag{1.1}$$

where $y = 0$ stands for the rigid wall, x (resp. y) is the tangential (resp. normal) variable to the wall. The functions u_E, p_E are given by the outer flow: more precisely u_E (resp. p_E) is the trace at the boundary of the tangential velocity (resp. of the pressure) of a flow satisfying the Euler equations. The functions u_E, p_E are linked by the relation

$$u_E u_E' = -\frac{dp_E(x)}{dx}.$$

Existence results for (1.1) were first obtained by Oleinik (see [29, Theorem 2.1.1]). Indeed, as long as u is positive (i.e. when there is no recirculation within the boundary layer), (1.1) can be considered as a non-local transport-diffusion equation in which the tangential variable x plays the role of “time”. The function u_0 , which is the input flow, is then considered as an “initial data”. However, this point of view breaks down as soon as u takes negative values. Physical experiments and numerical simulations show that such behavior may occur; in this case, the boundary layer seems to detach itself from the boundary. This phenomenon is therefore referred to as “boundary layer separation” (see figure 1).

The goal of this paper is to prove that separation does occur for the stationary Prandtl model (1.1), and to give a quantitative description of the solution close to (but on the left of) the separation point. In particular, we will justify rigorously the “Goldstein singularity” (see [7]).

1.1 Setting of the problem and state of the art

The first mathematical study of the stationary Prandtl equation was performed by Oleinik (see [29]):

Proposition 1.1 (Oleinik). *Let $\alpha > 0$, $X \in]0, +\infty[$. Let $u_0 \in C_b^{2,\alpha}(\mathbb{R})$ such that $u_0(0) = 0$, $u_0'(0) > 0$, $\lim_{y \rightarrow \infty} u_0(y) = u_E(0) > 0$, and such that $u_0(y) > 0$ for $y > 0$. Assume that $dp_E/dx \in C^1([0, X])$, and that for $y \ll 1$ the following compatibility condition is satisfied*

$$u_0''(y) - \frac{dp_E(0)}{dx} = O(y^2). \tag{1.2}$$

Then there exists $x^ \leq X$ such that equation (1.1) admits a solution $u \in C^1([0, x^*[\times \mathbb{R}_+)$ enjoying the following properties:*

- *Regularity: u is bounded and continuous in $[0, x^*] \times \mathbb{R}_+$, $\partial_y u, \partial_y^2 u$ are bounded and continuous in $[0, x^*[\times \mathbb{R}_+$, and $\partial_x u, v$ and $\partial_y v$ are locally bounded and continuous in $[0, x^*[\times \mathbb{R}_+$;*
- *Non-degeneracy: $u(x, y) > 0$ for all $y > 0$ $x \in [0, x^*[$, and for all $\bar{x} < x^*$ there exists $y_0 > 0$, $m > 0$ such that $\partial_y u(x, y) \geq m$ for all $(x, y) \in [0, \bar{x}] \times [0, y_0]$.*




FIG. 1. Velocity profiles in the boundary layer near separation

Figure 1: From Stewartson [33].

- *Sufficient condition for global solutions: if $\frac{dp_E(x)}{dx} \leq 0$, then the solution is global, i.e. $x^* = X$.*

In this paper, we are interested in the case where the solution of (1.1) is not global: more precisely, we consider the equation (1.1) with $dp_E/dx = 1$, i.e.

$$\begin{aligned}
 uu_x + vu_y - u_{yy} &= -1, & x \in (0, x_0), & y > 0, \\
 u_x + v_y &= 0, & x \in (0, x_0), & y > 0, \\
 u|_{x=0} = u_0, & u|_{y=0} = 0, & \lim_{y \rightarrow \infty} u(x, y) &= u_E(x),
 \end{aligned} \tag{P}$$

with $u_E(x) = \sqrt{2(x_0 - x)}$, for some $x_0 > 0$, and u_0 satisfies the assumptions of Proposition 1.1. Hence it is known that local solutions (in x) of (P) do exist. However, heuristically, it can be expected that the negative source term will diminish the value of the tangential velocity u , and that there might exist a point x^* beyond which the result of Proposition 1.1 cannot be used to extend the solution. More precisely, it can be checked easily that the compatibility condition (1.2) is propagated by equation (P). As a consequence, we have $x^* < x_0$ if and only if one of the following two conditions is satisfied:

- (i) $u_y(x^*, 0) = 0$;
- (ii) There exists $y^* > 0$ such that $u(x^*, y^*) = 0$.

In order to simplify the mathematical analysis, we will work with solutions of (P) that are increasing in y . This property is propagated by the equation, and ensures that situation (ii)

above never occurs. Consequently, for solutions which are increasing in y , we have $x^* < x_0$ if and only if

$$\frac{\partial u}{\partial y}(x^*, 0) = 0. \quad (1.3)$$

In the Physics literature (see for instance the seminal work of Goldstein [7], followed by the one of Stewartson [33]), this condition is used as a characterization of the “separation point”.

The first computational works on this subject go back to Goldstein [7] and Landau [20, Chapter 4, §40]. In particular, Goldstein uses an asymptotic expansion in self-similar variables to compute the profile of the singularity close to the separation point. These computations are later extended by Stewartson [33]. However, these calculations are formal; furthermore, some of the coefficients of the asymptotic expansion cannot be computed by either method. Independently, Landau proposes another characterization of the separation point, and gives an argument suggesting that $\partial_y u|_{y=0} \sim \sqrt{x^* - x}$ close to the separation point.

There are very few rigorous results about separation in the math literature. In the paper [4], Weinan E announces a result obtained in collaboration with Luis Caffarelli. This result states, under some structural assumption on the initial data, that the existence time x^* of the solutions of (P) in the sense of Oleinik is finite, and that the family $u_\mu(x, y) := \frac{1}{\sqrt{\mu}} u(\mu(x^* - x), \mu^{1/4} y)$ is compact in $\mathcal{C}(\mathbb{R}_+^2)$. Moreover, the author states two technical Lemmas playing a key role in the proof. However, to the best of our knowledge, the complete proof of this result was never published.

Let us also mention recent works by Guo and Nguyen [9] and by Iyer [12, 11, 13], in which the authors justify the Prandtl expansion either over a moving plate or over a rotating disk. Note that in these two cases, the velocity of the boundary layer on the boundary is non zero, which somehow prevents recirculation and separation.

In the time-dependent framework, boundary layer separation has also been tackled recently by Kukavica, Vicol and Wang [19], extending computations by Engquist and E [5]: starting from an analytic initial data, for a specific Euler flow, the authors prove that some Sobolev norm blows up in finite time. This is known as the van Dommelen and Shen singularity. Note that in this time-dependent context, separation is defined as the apparition of a singular behaviour, which is a somewhat different notion from the one we are describing in the present paper. This is related to the bad mathematical properties of the time-dependent Prandtl equation, which is known to be locally well-posed in analytic or Gevrey spaces [31, 21, 18, 17, 6] and in the monotone case [26], but ill-posed in Sobolev spaces [8, 6]. We also refer to the recent preprint [35] for a proof of separation in a time-dependent setting with the same criterion as the one used in the present paper, using virial-type arguments.

1.2 Main result

Our main result states that for a suitable class of initial data u_0 , the maximal existence “time” $x^* > 0$ of the solution given by Oleinik’s Theorem is finite: in other words, setting

$$\lambda(x) := \partial_y u|_{y=0},$$

there exists $x^* \in]0, +\infty[$ such that $\lim_{x \rightarrow x^*} \lambda(x) = 0$. Furthermore, for this class of initial data, we are able to quantify the rate of cancellation of $\lambda(x)$.

Let us now explicit our assumptions on the initial data u_0 :

(H1) $u_0 \in \mathcal{C}^7(\mathbb{R}_+)$, u_0 is increasing in y and $\lambda_0 := u_0'(0) > 0$;

(H2) There exists a constant $C_0 > 0$ such that

$$\begin{aligned} \forall y \geq 0, \quad -C_0 \inf(y^2, 1) \leq u_0''(y) - 1 \leq 0, \\ C_0^{-1} \leq -u_0^{(4)}(0) \leq C_0, \\ \|u_0\|_{W^{7,\infty}} \leq C_0. \end{aligned}$$

(H3) $u_0 = u_0^{\text{app}} + v_0$, where

$$\begin{aligned} u_0^{\text{app}} &= \lambda_0 y + \frac{y^2}{2} + u_0^{(4)}(0) \frac{y^4}{4!} \\ &\quad - c_7 (u_0^{(4)}(0))^2 \frac{y^7}{\lambda_0} + c_{10} (u_0^{(4)}(0))^3 \frac{y^{10}}{\lambda_0^2} + c_{11} (u_0^{(4)}(0))^3 \frac{y^{11}}{\lambda_0^3} \quad \text{for } y \leq \lambda_0^{3/7}, \\ |u_0^{\text{app}}| &\leq C_0 \quad \text{for } y \geq \lambda_0^{3/7}, \end{aligned}$$

and

$$|v_0| \leq C_0 \left(\lambda_0^{-3/2} (\lambda_0 y^7 + c_8 y^8) + \lambda_0^{-2} y^{10} + \lambda_0^{-3} y^{11} \right) \quad \text{for } y \leq \lambda_0^{3/7}.$$

In the expressions above, the constants c_i are universal and can be computed explicitly.

Remark 1.2. • *These assumptions are actually not optimal: in fact, condition (H3) merely ensures that some energy-like quantities are small enough. However, the actual condition we need is complicated to state at this stage: we refer to the statement of Theorem 3, in rescaled variables, for a less stringent condition.*

- *Notice that $|v_0| \ll u_0^{\text{app}}$ if $\lambda_0 \ll 1$: the term v_0 is the initial data for the corrector term $v = u - u^{\text{app}}$. The main issue of the paper is to have a good control of v close to $y = 0$.*
- *The monotony assumption on u_0 ensures that separation occurs at $y = 0$. The monotony is preserved by the Prandtl equation for $x > 0$.*
- *Notice that we prescribe the Taylor expansion of u_0 up to order 7. In other words, we impose a high order compatibility condition on the initial data, because we need to derive estimates on derivatives of u .*

Theorem 1. *Consider the Prandtl equation with adverse pressure gradient (P) and with an initial data $u_0 \in C^7(\mathbb{R}_+)$ satisfying (H1)-(H3). Then for any $\eta > 0$, $C_0 > 0$, there exists $\epsilon_0 > 0$ such that if $\lambda_0 < \epsilon_0$, the “existence time” x^* is finite, and $x^* = O(\lambda_0^2)$. Furthermore, setting $\lambda(x) := \partial_y u|_{y=0}(x)$, there exists a constant $C > 0$, depending on u_0 , such that*

$$\lambda(x) \sim C \sqrt{x^* - x} \quad \text{as } x \rightarrow x^*.$$

The proof of Theorem 1 relies on several ingredients: the first step is to perform a self-similar change of variables, using $\lambda(x)$ as a scaling factor. Then the issue is to control the variations of λ , or more precisely, of $b := -2\lambda_x \lambda^3$. The method thanks to which we construct an approximate solution and find the ideal ODE on b is inspired from the theory of modulation of variables, which was initiated formally by Zakharov and Shabat (see [37] and the presentation in the book by Sulem and Sulem [34]) and rigorously applied by Merle and Raphaël to blow-up phenomena in the nonlinear Schrödinger equation [28, 27].

Once the approximate solution is constructed, the whole problem amounts to controlling the remainder v . To that end, we exhibit a transport-diffusion structure of equation (P) (or of its rescaled version, see equation (2.15)). Let us emphasize that this structure, to our knowledge, is entirely new. We perform energy estimates that rely strongly on the structure of the equation. In order to handle nonlinearities, we will also need to control u in L^∞ . Therefore we derive pointwise estimates on u and its derivatives by constructing sub and super-solutions and using the maximum principle.

Let us point out that in order to carry these estimates, we will use three different versions of the equation. The first one is merely a rescaling of equation (P) (see (2.3)). It will be used to compute explicitly the approximate solution and find the ODE on b . The second one is a transport equation with a non local diffusion term (see (2.16)). Its purpose is to perform energy estimates, and the major difficulty will be to find good coercivity inequalities on the diffusion. Eventually, we will use a change of variables to transform (2.3) into a nonlinear transport diffusion equation of porous medium type (see (2.24)). This last form was already used by Oleinik in [29] and will be suitable for the maximum principle and will help us prove the L^∞ estimates.

In the next section, we present our scheme of proof and state our main intermediate results. The reader who is not interested in the technical details of the proof can focus on section 2, that gives an overall idea of the main arguments involved. The third section is devoted to the construction of sub and super solutions. In section 4, we introduce several tools that play an important role in the energy estimates: coercivity of the diffusion term, commutator Lemma, computation of the remainder... Eventually, we prove the energy estimates in section 5.

Remark 1.3. *Our result actually gives much more information on u : in fact, we construct an approximate solution u^{app} , which contains the main order terms in the Taylor expansion of u , and we control $v = u - u^{app}$. As a corollary, we find that the sequence of functions $(u^\mu)_{\mu>0}$ from the statement of Luis Caffarelli and Weinan E converges towards $z^2/2$ in the zone $z \leq \mu^{-1/12}\xi^{1/6}$, $\xi \lesssim 1$ (see Remark 2.17 for more details). Hence our result holds under more stringent assumptions on the initial data, but it gives a much more quantitative and precise description of the asymptotic behaviour.*

2 Strategy of proof

2.1 Self-similar change of variables

Let us first recall that equation (P) has a scaling invariance: indeed, if (u, v) is a solution of (P), then for any $\mu > 0$, the couple (u_μ, v_μ) defined by

$$u_\mu = \frac{1}{\sqrt{\mu}}u(\mu x, \mu^{1/4}y), \quad v_\mu = \mu^{1/4}v(\mu x, \mu^{1/4}y),$$

is still a solution of (P). This scaling invariance has been used by Goldstein [7] and Stewartson [33] to compute exact solutions of (P) close to the separation point. These special solutions were sought as formal infinite series in some rescaled variables.

In the present article, the idea is to perform a change of variables which relies on this scaling invariance and which depends on the solution itself. It incorporates information on the “separation rate”, i.e. on the speed of cancellation of $\partial_y u|_{y=0}$. This type of idea was used

by F. Merle and P. Raphaël in the context of singularity analysis for the nonlinear Schrödinger equation [28, 27]. More precisely, define

$$\lambda(x) := \partial_y u|_{y=0} \quad \text{and} \quad Y = \frac{y}{\lambda(x)}.$$

We also change the tangential variable and define the variable s by

$$\frac{ds}{dx} = \frac{1}{\lambda^4(x)}. \quad (2.1)$$

Then the new unknown function is

$$U(s, Y) := \lambda^{-2}(x(s))u(x(s), \lambda(x(s))Y). \quad (2.2)$$

It can be easily checked that U is a solution of the equation

$$UU_s - U_Y \int_0^Y U_s - bU^2 + \frac{3b}{2}U_Y \int_0^Y U - U_{YY} = -1, \quad (2.3)$$

where

$$b = -2\lambda_x \lambda^3 = -2\frac{\lambda_s}{\lambda}. \quad (2.4)$$

The boundary conditions become

$$U|_{Y=0} = 0, \quad \lim_{Y \rightarrow \infty} U(s, Y) = U_\infty(s), \quad (2.5)$$

where U_∞ satisfies $U_\infty U'_\infty - bU_\infty^2 = -1$. Moreover, thanks to the definition of λ , we have

$$\partial_Y U|_{Y=0} = 1. \quad (2.6)$$

From now on, we will work with equation (2.3) only. The goal is to construct an approximate solution of (2.3), together with $b(s)$ and $\lambda(s)$, having nice stability properties as $s \rightarrow \infty$. Note that the limit $s \rightarrow \infty$ corresponds to the limit $x \rightarrow x^*$ in the original variables. As we will see in the next paragraph, the stability properties of the approximate solution are intimately connected to the asymptotic law of b as $s \rightarrow \infty$. Eventually, the asymptotic behavior of b will dictate the rate of cancellation of λ close to $x = x^*$. We prove that the behavior $b(s) \sim s^{-1}$ is stable. This asymptotic law corresponds to the separation rate announced in Theorem 1, namely $\lambda(x) \sim C\sqrt{x^* - x}$.

In the next paragraphs, we explain how we construct the approximate solution, and which energy estimates are used to prove its stability. We deal with nonlinearities in the equation by using the maximum principle, together with Sobolev embeddings. Let us recall that we will in fact use three different forms of equation (2.3):

- Due to its polynomial form, equation (2.3) itself is very useful to construct the approximate solution and find the correct asymptotic law for b ;
- In order to perform energy estimates, we will transform (2.3) into a transport-diffusion equation (with a non-local diffusion term), see (2.15) and (2.16);
- Eventually, in order to use the maximum principle, we rely on a third version of (2.3), that uses von Mises variables. The equation then becomes a nonlinear local transport-diffusion equation.

2.2 Construction of an approximate solution

The heuristic idea behind the construction of stable approximate solutions is the following: we look for an approximate solution U^{app} of (2.3) with a remainder as small as possible. In particular, the remainder for U^{app} should have the lowest possible growth at infinity. This implies that the function U^{app} itself should have the lowest possible growth as $Y \rightarrow \infty$, as we shall see in a moment. As in the work of Merle and Raphaël in the context of the nonlinear Schrödinger equation, this low growth condition has an immediate impact on the asymptotic behavior of the function b .

We decompose the definition of approximate solutions into three zones: the main zone goes from 0 to s^α , for some $\alpha > 0$ to be defined later on. In this zone, we compute a Taylor expansion of $U(s, Y)$ for Y close to zero, and we try to push the expansion as far as possible, which amounts to the “low growth condition” explained above. In the second zone, we only keep the largest term in the Taylor expansion, namely $Y^2/2$. It can be checked that $Y^2/2$ is a stationary solution of (2.3). This stationary solution corresponds to a solution of (P) which is independent of x and scaling invariant, namely $(x, y) \mapsto y^2/2$. In the third zone, we connect $Y^2/2$ to an asymptotic profile $U_\infty^{\text{app}}(s)$. Notice that if $b(s) = s^{-1} + O(s^{-\eta-1})$ for some $\eta > 0$, then $U_\infty(s) = s + 1 + o(1)$, and therefore we also take $U_\infty^{\text{app}}(s) \sim s$.

Throughout this paragraph, we will rely on the polynomial form of the rescaled Prandtl equation, namely (2.3).

- *Taylor expansion of U for Y close to zero:*

Let us first recall that thanks to the change of variables (2.2), we have

$$U(s, 0) = 0, \quad \partial_Y U(s, 0) = 1.$$

It then follows from (2.3) that

$$\partial_{YY} U(s, 0) = 1.$$

The first terms of the Taylor expansion of U for Y close to zero are therefore $Y + \frac{Y^2}{2}$.

The first natural idea is to define a sequence of polynomials in Y with coefficients depending on s thanks to the induction relation

$$\begin{aligned} U_1(s, Y) &:= Y + \frac{Y^2}{2}, \\ \partial_{YY}(U_{N+1} - U_N) &:= 1 + U_N \partial_s U_N - \partial_Y U_N \int_0^Y \partial_s U_N - b U_N^2 + \frac{3b}{2} \partial_Y U_N \int_0^Y U_N - \partial_{YY} U_N. \end{aligned} \quad (2.7)$$

We obtain easily that

$$U_2(s, Y) := Y + \frac{Y^2}{2} - a_4 b Y^4, \quad \text{with } a_4 = \frac{1}{48}. \quad (2.8)$$

Let us now compute the error terms generated by U_2 . We have

$$\begin{aligned} &U_2 \partial_s U_2 - \partial_Y U_2 \int_0^Y \partial_s U_2 - b U_2^2 + \frac{3b}{2} \partial_Y U_2 \int_0^Y U_2 - \partial_{YY} U_2 + 1 \\ &= -a_4 \left(\frac{4}{5} b_s + \frac{13}{10} b^2 \right) Y^5 - \frac{3}{10} a_4 (b_s + b^2) Y^6 + a_4^2 \frac{b}{5} (b_s + b^2) Y^8. \end{aligned}$$

Let us recall that we expect that $b(s) = O(s^{-1})$ as $s \rightarrow \infty$. Therefore the coefficient of the last term in the right-hand side is one order of magnitude smaller than the first two terms. We thus focus on the comparison between the first two terms in the right-hand side. As explained above, the goal is to choose the approximate solution with the smallest growth at infinity. Note that the remainder term $(b_s + b^2)Y^6$ would yield in U_3 a term proportional to $(b_s + b^2)Y^8$, whereas the remainder term $(\frac{4}{5}b_s + \frac{13}{10}b^2)Y^5$ would yield a term proportional to $(\frac{4}{5}b_s + \frac{13}{10}b^2)Y^7$. Consequently, we choose to “cancel out” the term $(b_s + b^2)Y^8$ in U_3 . In other words, in the induction formula (2.7) defining the sequence $(U_N)_{N \geq 1}$, we replace every occurrence of b_s by $-b^2$. The polynomial U_3 is therefore defined by

$$U_3(s, Y) := Y + \frac{Y^2}{2} - a_4 b Y^4 - a_7 b^2 Y^7, \quad \text{with } a_7 = \frac{1}{84} a_4.$$

It follows that for $Y \ll 1$,

$$U(s, Y) \simeq U_3(s, Y) + V_3(s, Y),$$

where $V_3(s, Y) = -a_7 \frac{8}{5} (b_s + b^2) Y^7 - a_4 \frac{3}{10 \times 7 \times 8} (b_s + b^2) Y^8 + O(Y^{10})$. (2.9)

Remark 2.1. *In the work of Merle and Raphaël, the choice of the parameters λ and b stems from orthogonality properties of the quantity $U - U^{app}$ on some well chosen functions. In the present case, these orthogonality properties can be seen as a cancellation at high enough order of $U - U^{app}$ at $Y = 0$.*

For technical reasons, it is necessary to push further the expansion of U . We thus compute U_4 . We find that

$$\begin{aligned} & U_3 \partial_s U_3 - \partial_Y U_3 \int_0^Y \partial_s U_3 - b U_3^2 + \frac{3b}{2} \partial_Y U_3 \int_0^Y U_3 - \partial_{YY} U_3 + 1 \\ = & (b_s + b^2) \left[-\frac{4}{5} a_4 Y^5 - \frac{3}{10} a_4 Y^6 - \frac{a_4}{60} b Y^8 - \frac{3}{4} a_7 b Y^9 + \frac{3}{5} a_4 a_7 b^2 Y^{11} + \frac{1}{4} a_7^2 b^3 Y^{14} \right] \\ & - \frac{27}{16} a_7 b^3 Y^8 - \frac{3}{16} a_7 b^3 Y^9 + \frac{11}{16} a_4 a_7 b^4 Y^{11} + \frac{3}{8} a_7^2 b^5 Y^{14}. \end{aligned}$$

It follows that

$$U_4 = U_3 - a_{10} b^3 Y^{10} - a_{11} b^3 Y^{11} + a_{13} b^4 Y^{13} + a_{16} b^5 Y^{16},$$

where

$$a_{10} = \frac{27}{16 \times 90} a_7, \quad a_{11} = \frac{3}{16 \times 110} a_7, \quad a_{13} = \frac{11 a_4 a_7}{13 \times 12 \times 16}, \quad a_{16} = \frac{a_7^2}{16 \times 5 \times 8}. \quad (2.10)$$

• *Definition of the approximate solution:*

We now define the approximate solution U^{app} in the following way: let $\Theta \in \mathcal{C}^2(\mathbb{R}_+)$ be such that $\Theta(\xi) = \frac{\xi^2}{2}$ for $\xi \leq c_0$ for some $c_0 > 0$, Θ strictly increasing, and $\Theta(\xi) \rightarrow 1$ as $\xi \rightarrow \infty$. Let $\chi \in \mathcal{C}_0^\infty(\mathbb{R}_+)$ be such that $\chi \equiv 1$ in a neighbourhood of zero. We take

$$U^{app}(s, Y) := \chi \left(\frac{Y}{s^{2/7}} \right) \left[Y - a_4 b Y^4 - a_7 b^2 Y^7 - a_{10} b^3 Y^{10} - a_{11} b^3 Y^{11} \right] + \frac{1}{b} \Theta(\sqrt{b} Y). \quad (2.11)$$

Notice that $U^{\text{app}} \simeq U_4$ as long as $Y \lesssim s^{2/7}$ (the highest order terms have been removed, mainly because they do not lower the size of the remainder while making the computations heavier), and that $U^{\text{app}} \rightarrow \frac{1}{b}$ as $Y \rightarrow \infty$. Therefore we do not require that $U^{\text{app}} - U(s, Y) \rightarrow 0$ as $Y \rightarrow \infty$. But this is not an issue, since we will measure the distance between U and U^{app} in weighted Sobolev spaces, with weights decreasing polynomially (with a large power) after s^β , for some $\beta < 2/7$.

Remark 2.2. *The zone after which we cut-off the first part of the approximate solution is irrelevant: we could have used any cut-off $\chi(\cdot/s^\alpha)$ as long as $\alpha \in]1/4, 1/3[$. The choice $\alpha = 2/7$ simplifies some of the statements on U^{app} since it ensures that Y and $-a_4 b Y^4$ are the largest terms in $Y - a_4 b Y^4 - a_7 b^2 Y^7 - a_{10} b^3 Y^{10} - a_{11} b^3 Y^{11}$.*

We also set, in the rest of the paper, $V := U - U^{\text{app}}$. The computations above and in particular (2.9) show that

$$V(s, Y) = -a_7 \frac{8}{5} (b_s + b^2) Y^7 - a_4 \frac{3}{10 \times 7 \times 8} (b_s + b^2) Y^8 + O(Y^{10}) \quad \text{for } 0 < Y \ll 1.$$

In particular, let \mathcal{N} be a (semi-)norm on functions $W \in \mathcal{C}^8(\mathbb{R}_+)$ such that $W = O(Y^7)$ for Y close to zero. Assume that there exists a constant $C_{\mathcal{N}}$ such that

$$\mathcal{N}(W) \geq C_{\mathcal{N}} |\partial_Y^7 W|_{Y=0}.$$

Then $\mathcal{N}(U - U^{\text{app}}) \geq C_{\mathcal{N}} |b_s + b^2|$. Therefore the goal of the paper is to use the structure of the equation (2.3) in order to find a semi-norm \mathcal{N} which satisfies the assumptions above, and to prove that

$$\mathcal{N}(U - U^{\text{app}}) \leq C s^{-2-\eta},$$

for some positive constant η and for s sufficiently large, or alternatively, that

$$\int_{s_0}^{\infty} s^{3+2\eta} \mathcal{N}(U(s) - U^{\text{app}}(s)) ds < +\infty.$$

Indeed, we have the following result:

Lemma 2.3. *Let $s_0 := b_0^{-1}$. Assume that the variables x, s and the parameters λ, b are related by the formulas (2.1), (2.4) with the initial conditions $\lambda|_{s=s_0} = \lambda_0$, $b|_{s=s_0} = b_0 = s_0^{-1}$, and that there exists a constant c_0 such that*

$$c_0^{-1} \leq \frac{b_0}{\lambda_0^2} \leq c_0.$$

Assume furthermore that there exist constants $\eta > 0$ and $\epsilon \in (0, 1)$ such that for all $s \geq s_0$,

$$\begin{aligned} \int_{s_0}^{\infty} s^{3+2\eta} |b_s + b^2|^2 ds &< \infty, \\ \frac{1-\epsilon}{s} &\leq b(s) \leq \frac{1+\epsilon}{s}. \end{aligned} \tag{2.12}$$

Then there exists $x^ > 0$ such that $\lambda(x) \rightarrow 0$ as $x \rightarrow x^*$. Furthermore, if $\lambda_0 \ll 1$, then $x^* = O(\lambda_0^2)$ and there exists a constant C such that*

$$\lambda(x) \sim C(x^* - x)^{1/2} \quad \text{as } x \rightarrow x^*.$$

Proof. First, setting

$$J := \int_{s_0}^{\infty} s^{3+2\eta} |b_s + b^2|^2 ds$$

and using Lemma B.1 in the Appendix, we know that

$$b(s) = \frac{1}{s} + r(s),$$

where

$$\forall s \geq s_0, \quad |r(s)| \leq \epsilon \frac{1 + \epsilon s_0}{1 - \epsilon s^2} + J^{1/2} \frac{1 + \epsilon}{(1 - \epsilon)^2} \frac{1}{s^{1+\eta}}.$$

As a consequence,

$$\int_{s_0}^{\infty} |r(s)| ds \leq \epsilon \frac{1 + \epsilon}{1 - \epsilon} + J^{1/2} \frac{1 + \epsilon}{(1 - \epsilon)^2 \eta s_0^\eta} < \infty.$$

From (2.1) and (2.4), it follows that

$$\lambda(s) = \lambda(s_0) \exp\left(-\frac{1}{2} \int_{s_0}^s b(s') ds'\right) = \lambda(s_0) \left(\frac{s_0}{s}\right)^{1/2} \exp\left(-\frac{1}{2} \int_{s_0}^s r(s') ds'\right).$$

We have $\lambda(s_0) = \lambda_0 = O(s_0^{-1/2})$ by assumption. Moreover, according to the estimate of r above, the function $\psi(s) := \exp\left(-\frac{1}{2} \int_{s_0}^s r(s') ds'\right)$ has a finite, strictly positive limit ψ_∞ as $s \rightarrow \infty$. As a consequence $\lambda(s) = \left(\lambda_0 s_0^{1/2}\right) \psi(s) s^{-1/2}$ for all $s \geq s_0$. According to (2.1), we have

$$x^* := \int_{s_0}^{\infty} \lambda(s)^4 ds = \left(\lambda_0 s_0^{1/2}\right)^4 \int_{s_0}^{\infty} \psi(s)^4 s^{-2} ds < \infty.$$

Thus separation occurs at a finite x^* . Moreover,

$$x^* - x(s) = \left(\lambda_0 s_0^{1/2}\right)^4 \int_s^{\infty} \psi(s')^4 s'^{-2} ds' \sim \left(\lambda_0 s_0^{1/2}\right)^4 \psi_\infty^4 s^{-1} \quad \text{as } s \rightarrow \infty.$$

Going back to the original variables, we deduce that

$$\lambda(x) \sim \frac{1}{\left(\lambda_0 s_0^{1/2}\right) \psi_\infty} (x^* - x)^{1/2} \quad \text{as } x \rightarrow x^*.$$

Using the above formulas, we also infer that if $s_0 \gg 1$, $x^* = O(s_0^{-1}) = O(\lambda_0^2)$. □

Remark 2.4. Notice that the precise value of the separation point x^* depends on the whole function $u_0(y)$ (and not only on its derivatives at $y = 0$) through the function $\psi(s)$. This intricate dependance might explain why the coefficients in Goldstein's expansion were undetermined.

Remark 2.5. Let us now give some examples of norms \mathcal{N} such that

$$\mathcal{N}(W) \geq C_{\mathcal{N}} |\partial_Y^7 W|_{Y=0} \tag{2.13}$$

for $W \in \mathcal{C}^8(\mathbb{R}_+)$ with $W = O(Y^7)$ for Y close to zero. We can take for instance

$$\mathcal{N}(W)^2 := \int_0^\infty (\partial_Y^7 W)^2 + (\partial_Y^8 W)^2,$$

or

$$\mathcal{N}(W)^2 := \int_0^\infty \left(\frac{W}{Y^7} \right)^2 + \left(\partial_Y \frac{W}{Y^7} \right)^2.$$

More generally, we can use any norm \mathcal{N} such that

$$\mathcal{N}(W) \gtrsim \left\| Y^{k-7} \partial_Y^k W \right\|_{H^1(0, Y_0)},$$

for some fixed $Y_0 > 0$ and for any $k \in \{0, \dots, 7\}$. The norm \mathcal{N} we will use eventually will be equivalent to a linear combination of such norms in a neighbourhood of $Y = 0$.

2.3 Error estimates

In this paragraph, we explain roughly how estimates on $V := U - U^{\text{app}}$ are derived. More details will be given in sections 4 and 5. We emphasize that all energy estimates written in this paper are new. The first step is to compute an evolution equation of transport-diffusion type on V . To that end, let us consider equation (2.3), and set, for $W_1, W_2 \in \mathcal{C}(\mathbb{R}_+)$,

$$L_{W_1} W_2 := W_1 W_2 - \partial_Y W_1 \int_0^Y W_2,$$

so that equation (2.3) can be written as

$$L_U U_s - bU^2 + \frac{3b}{2} U_Y \int_0^Y U - \partial_{YY} U = -1.$$

Notice that since $U(s, Y) > 0$ for all $s, Y > 0$,

$$\frac{L_U W}{U^2} = \partial_Y \left(\frac{\int_0^Y W}{U} \right).$$

Hence we can define the inverse of the operator L_U , for functions f such that $f(Y)/Y^2$ is integrable in a neighbourhood of zero: we have

$$L_U^{-1} f = \left(U \int_0^Y \frac{f}{U^2} \right)_Y = U_Y \int_0^Y \frac{f}{U^2} + \frac{f}{U}. \quad (2.14)$$

As a consequence, the equation on U can be written as

$$\partial_s U + bL_U^{-1} \left(\frac{3}{2} U_Y \int_0^Y U - U^2 \right) - L_U^{-1} (\partial_{YY} U - 1) = 0.$$

It follows immediately from the definition that

$$L_U^{-1}(U^2) = \partial_Y \left(U \int_0^Y 1 \right) = (YU)_Y.$$

On the other hand,

$$\begin{aligned} L_U^{-1} \left(U_Y \int_0^Y U \right) &= L_U^{-1} \left(U_Y \int_0^Y U - U^2 \right) + (YU)_Y \\ &= -L_U^{-1} L_U U + (YU)_Y \\ &= -U + (YU)_Y = YU_Y. \end{aligned}$$

We infer that the equation on U becomes

$$\partial_s U - bU + \frac{b}{2}Y\partial_Y U - L_U^{-1}(\partial_{YY}U - 1) = 0. \quad (2.15)$$

The whole non-linearity of the equation is now encoded in the diffusion term $L_U^{-1}(\partial_{YY}U - 1)$.

Setting $\mathcal{L}_U := L_U^{-1}\partial_{YY}$, the equation on $V = U - U^{\text{app}}$ becomes

$$\partial_s V - bV + \frac{b}{2}Y\partial_Y V - \mathcal{L}_U V = \mathcal{R}, \quad (2.16)$$

where the remainder \mathcal{R} is defined by

$$\mathcal{R} := - \left(\partial_s U^{\text{app}} - bU^{\text{app}} + \frac{b}{2}Y\partial_Y U^{\text{app}} \right) + L_U^{-1}(\partial_{YY}U^{\text{app}} - 1).$$

Equation (2.16) is the second form we will be using for the rescaled Prandtl equation. It will be handy for the derivation of energy estimates.

We have the following result, which is proved in section 4:

Lemma 2.6. *The remainder term \mathcal{R} can be decomposed as*

$$\begin{aligned} \mathcal{R} &= (b_s + b^2) \chi \left(\frac{Y}{s^{2/7}} \right) [a_4 Y^4 + 2a_7 b Y^7 + 3a_{10} b^2 Y^{10} + 3a_{11} b^2 Y^{11}] \\ &+ \chi \left(\frac{Y}{s^{2/7}} \right) [a_{10} b^4 Y^{10} + a_{11} 3b^4 Y^{11}/2] \\ &+ \frac{b}{2} L_U^{-1}(L_V Y) + \frac{a_7 b^3}{2} L_U^{-1} \left(\chi \left(\frac{Y}{s^{2/7}} \right) (L_V Y^7 + L_{-a_4 b Y^4 - a_7 b^2 Y^7 + a_{10} b^3 Y^{10} + a_{11} b^3 Y^{11} Y^7}) \right) \\ &+ P_1(s, Y) + L_U^{-1}(P_2(s, Y)) \end{aligned}$$

where $P_1, P_2 \in \mathcal{C}^0([s_0, \infty), \mathcal{C}^\infty(\mathbb{R}_+))$ are such that P_i has at most polynomial growth in s and Y and $P_i(s, Y) = 0$ for $Y \leq cs^{2/7}$ for some $c > 0$.

Remark 2.7. *Following the decomposition of Lemma 2.6, we write $\mathcal{R} = \sum_{i=1}^4 \mathcal{R}_i$. Each of the remainder terms \mathcal{R}_i will play a different role and will be treated separately. More precisely:*

1. *Cancellations will occur in the remainder term \mathcal{R}_1 ;*
2. *The size of the term \mathcal{R}_2 dictates the final rate of convergence of the energy. This is where the choice of the approximate solution plays an important role;*
3. *The term \mathcal{R}_3 can be treated as perturbation of the zero order term bV and of the transport term $bY\partial_Y V$ as soon as $Y \gg 1$, and as a perturbation of the diffusion $\mathcal{L}_U V$ if $Y \ll s^{1/4}$. Indeed, think of L_U^{-1} as a division by U , and of a derivation with respect to Y as a division by Y . Then*

$$|bL_U^{-1}(L_V Y)| \lesssim b \frac{Y^2}{U} |V_Y| \lesssim b \frac{1}{1+Y} |Y\partial_Y V|.$$

Thus if $Y \gg 1$, this term is small compared to $bY\partial_Y V$. On the other hand, heuristically, $|\mathcal{L}_U V| \gtrsim U^{-1} |\partial_Y^2 V| \gtrsim (YU)^{-1} |\partial_Y V|$ (think for instance of a Hardy inequality). Thus as long as $Y^2 U \ll b^{-1}$, i.e. $Y \ll s^{1/4}$, the diffusion term $\mathcal{L}_U V$ dominates $bY\partial_Y V$, and therefore \mathcal{R}_3 .

4. The last term \mathcal{R}_4 will not play any role in the energy estimates: indeed, we will choose weights with a strong polynomial decay for $Y \geq s^\beta$ for some $\beta < 2/7$, so that the error stemming from \mathcal{R}_4 can be made $O(s^{-P})$ for any $P > 0$ by an appropriate choice of the weight.

The idea is now to perform weighted energy estimates on equation (2.16), with the help of a norm \mathcal{N} satisfying assumption (2.13). These estimates rely on the following ideas:

1. Let \mathcal{N} be a norm satisfying (2.13), and define an energy $E(s)$ by

$$E(s) = \mathcal{N}(V(s))^2.$$

In order to prove that $b_s + b^2 = O(s^{-2-\eta})$ (or that $\int_{s_0}^{+\infty} s^{3+2\eta}(b_s + b^2)^2 ds < +\infty$) for some $\eta > 0$, it is enough to show that

$$\frac{dE}{ds} + \frac{\alpha}{s}E(s) \leq \rho(s) \quad \forall s \geq s_0 \quad (2.17)$$

with $4 + \eta \leq \alpha$, and with a right-hand side $\rho(s)$ such that $\int_{s_0}^{\infty} s^\alpha \rho(s) ds < +\infty$. Indeed, integrating (2.17) between s_0 and s yields

$$E(s) \leq s^{-\alpha} \left(E(s_0)s_0^\alpha + \int_{s_0}^{\infty} s^\alpha \rho(s) ds \right).$$

Assuming additionally that $E(s_0) \leq C_0 s_0^{-\alpha}$ for some constant C_0 independent of s_0 , we are led to

$$c_{\mathcal{N}} |b_s + b^2|^2 \leq E(s) \leq C s^{-4-\eta} \quad \forall s \geq s_0,$$

and using Lemma 2.3, we obtain the desired result.

2. The property $\alpha > 4$ in (2.17) is derived thanks to algebraic manipulations on (2.16). Schematically, if we only keep the transport part in (2.16) and if we consider the model equation

$$\partial_s f - \frac{1}{s}f + \frac{1}{2s}Y \partial_Y f = r,$$

we see that the k -th derivative of f satisfies

$$\partial_s \partial_Y^k f + \left(\frac{k}{2} - 1 \right) \frac{1}{s} \partial_Y^k f + \frac{1}{2s} Y \partial_Y^{k+1} f = \partial_Y^k r.$$

Hence

$$\frac{d}{ds} \|\partial_Y^k f\|_{L^2}^2 + \left(k - \frac{5}{2} \right) \frac{1}{s} \|\partial_Y^k f\|_{L^2}^2 = \int \partial_Y^k r \partial_Y^k f$$

Taking $k = 7$ and $k = 8$ and summing the two estimates, we obtain the desired result with $\alpha = \frac{9}{2}$ and $\mathcal{N}(f) := \|\partial_Y^7 f\|_{H^1}$.

3. The fact that the remainder is integrable with a weight s^α in (2.17) stems from our choice of approximate solution. In particular, if we modify the algorithm of construction of $(U_N)_{N \geq 1}$ described in the previous paragraph and replace every occurrence of b_s by $-cb^2$ for some constant $c \neq 1$, the estimate is no longer true.

Let us now explain the main steps in the derivation of estimate (2.17). The difficulties lie in the complex structure of the diffusion operator \mathcal{L}_U , and in the estimation of some commutator terms. The idea is to apply several times the operator \mathcal{L}_U to equation (2.16). This requires:

1. computing the commutator of \mathcal{L}_U with $\partial_s + \frac{b}{2}Y\partial_Y$;
2. understanding the action of \mathcal{L}_U on the remainder term \mathcal{R} ;
3. obtaining energy estimates on transport-diffusion equations of the type

$$\partial_s f + Cbf + \frac{b}{2}Y\partial_Y f - \mathcal{L}_U f = r. \quad (2.18)$$

Let us now explain how we deal with each of the points above.

2.3.1 Commutator of \mathcal{L}_U with $\partial_s + \frac{b}{2}Y\partial_Y$

The commutator result is stated in the following

Lemma 2.8 (Computation of the commutator). *For any function $W \in W_{loc}^{1,1}((s_0, \infty) \times \mathbb{R}_+)$ such that $W = O(Y^2)$ for Y close to zero,*

$$\left[L_U^{-1}, \partial_s + \frac{b}{2}Y\partial_Y \right] W = bL_U^{-1}W - \left(\mathcal{D} \int_0^Y \frac{W}{U^2} \right)_Y + 2 \left(U \int_0^Y \frac{W}{U^3} \mathcal{D} \right)_Y,$$

where $\mathcal{D} := L_U^{-1}(\partial_{Y^2}U - 1)$.

In the rest of the article, we define the commutator operator

$$\mathcal{C}[W] := - \left(\mathcal{D} \int_0^Y \frac{W}{U^2} \right)_Y + 2 \left(U \int_0^Y \frac{W}{U^3} \mathcal{D} \right)_Y.$$

The quantity \mathcal{D} involved in the commutator can be written as

$$\mathcal{D} = \mathcal{L}_U V + L_U^{-1}(\partial_{Y^2}U^{\text{app}} - 1),$$

where the second term can be developed using the explicit expression U^{app} . Notice the commutator $\mathcal{C}[\partial_Y^2 V]$ contains some quadratic terms in V . In order to estimate these quadratic terms, we will need both preliminary estimates and estimates in L^∞ on the function V . The L^∞ estimates are derived in detail in section 3, and rely on a careful use of the maximum principle.

2.3.2 Action of \mathcal{L}_U on the remainder term \mathcal{R}

We will use the decomposition of the remainder given in Remark 2.7. The first two terms, namely \mathcal{R}_1 and \mathcal{R}_2 , are essentially polynomials. Therefore, in order to deal with them, we will need to get explicit formulas for terms such as $\mathcal{L}_U(Y^4) = 12L_U^{-1}(Y^2)$, and more generally, to understand the asymptotic behavior of $L_U^{-1}(Y^k)$ for $Y \gg 1$ and $k \geq 2$.

In order to get explicit formulas, we will use in several instances the following trick: for $k \in \mathbb{N}$, write

$$Y^k = L_U^{-1}L_U(Y^k) = L_U^{-1}(L_{U^{\text{app}}}(Y^k) + L_V Y^k).$$

Now, since U^{app} is a polynomial, $L_{U^{\text{app}}}(Y^k)$ can be easily computed, and is also a polynomial in Y . The term $L_U^{-1}L_V(Y^k)$ is expected to be of lower order. For instance, taking $k = 1$, we observe that $L_{U^{\text{app}}}(Y) = \frac{Y^2}{2} + O(bY^5)$ for $Y \ll s^{2/7}$. Hence we obtain a formula for $L_U^{-1}(Y^2)$ (up to some remainder terms).

Concerning the asymptotic behavior of $L_U^{-1}(Y^k)$ for $Y \gg 1$ and for $k \geq 4$, notice that the operator L_U^{-1} acts roughly like a division by U , as can be seen from the formula (2.14). Furthermore, the L^∞ estimates (see Proposition 2.16) will ensure that there exists a constant C such that

$$C^{-1}(Y + Y^2) \leq U(s, Y) \leq C(Y + Y^2) \quad \forall Y \lesssim s^{2/7}.$$

Therefore, L_U^{-1} behaves differently for $Y \ll 1$ and for $Y \gg 1$: for Y close to zero, applying L_U^{-1} amounts to dividing by Y , while for $Y \gg 1$, it amounts to dividing by Y^2 . We obtain that for $Y \ll s^{1/2}$ and $k \geq 4$,

$$L_U^{-1}(Y^k) = \begin{cases} O(Y^{k-1}) & \text{if } Y \ll 1, \\ O(Y^{k-2}) & \text{if } Y \gtrsim 1. \end{cases}$$

As explained in Remark 2.7, the term \mathcal{R}_3 is treated as a perturbation of the dissipation coming from the transport and the diffusion term. Eventually, since \mathcal{R}_4 is supported in $Y \gtrsim s^{2/7}$, while we use weights that have a strong polynomial decay for $Y \gtrsim s^\beta$ for some $\beta < 2/7$, the size of \mathcal{R}_4 in our energy norms will be smaller than that of $\mathcal{R}_1 + \mathcal{R}_2 + \mathcal{R}_3$.

2.3.3 Energy estimates on transport-diffusion equations of the type (2.18)

The most difficult part is proving coercivity and positivity estimates for the diffusion. We will rely on the diffusion Lemma 4.7, which makes an extensive use of weighted Hardy inequalities, see [23]. They also rely on the fact that if $U = Y + \frac{Y^2}{2} + O(bY^4)$, then $\partial_{YY}L_U^{-1}$ is ‘‘almost’’ a local differential operator (see the formulas in Lemma A.1 in the Appendix).

We will also often use the observation that if $Y \lesssim s^{1/4}$, then the diffusion term dominates, while for $Y \gtrsim s^{1/4}$, then the transport part becomes preponderant. Indeed, for $k \geq 6$, if $b \sim s^{-1}$

$$CbY^k + \frac{b}{2}Y\partial_Y Y^k \sim \left(\frac{k}{2} + C\right)\frac{1}{s}Y^k,$$

and $\mathcal{L}_U Y^k = O(Y^{k-4})$ for $Y \gg 1$.

It is easily checked that both terms are of the same order for $Y \sim s^{1/4}$, and diffusion (resp. transport) is dominant below (resp. above) that threshold.

We now turn towards the sequence of estimates on V . In the end, we seek to obtain estimates on $\partial_Y \mathcal{L}_U^2 V$ and $\partial_Y^2 \mathcal{L}_U^2 V$. We recall that $V \sim -C(b_s + b^2)Y^7$ for Y close to zero, and therefore $\partial_Y \mathcal{L}_U^2 V \sim -C(b_s + b^2)$ for Y close to zero.

Using Lemma 2.8, we infer that $\mathcal{L}_U V$ satisfies the following equation:

$$\partial_s \mathcal{L}_U V + b \mathcal{L}_U V + \frac{b}{2} Y \partial_Y \mathcal{L}_U V - \mathcal{L}_U^2 V = \mathcal{L}_U \mathcal{R} + \mathcal{C}[\partial_{YY} V]. \quad (2.19)$$

We get the following result:

Proposition 2.9. *Assume that:*

- There exists a constant $J > 0$ such that

$$\int_{s_0}^{s_1} s^{13/4} |b_s + b^2|^2 ds \leq J;$$

- $\frac{1-\bar{\epsilon}}{s} \leq b \leq \frac{1+\bar{\epsilon}}{s}$ for $s \in [s_0, s_1]$ and for some small universal constant $\bar{\epsilon}$ (say $\bar{\epsilon} = 1/50$);
- There exist constants M_1, M_2, c independent of s such that for all Y ,

$$\begin{aligned} -M_1 &\leq U_{YY} \leq 1 \quad \forall Y \geq 0, \\ 1 - M_2 b Y^2 &\leq U_{YY} \quad \forall Y \in [0, cs^{1/3}]. \end{aligned}$$

Let $w_1 := Y^{-a}(1 + s^{-\beta_1}Y)^{-m_1}$ for some $\beta_1 \in]1/4, 2/7[$, $m_1 \in \mathbb{N}$.

Let

$$\begin{aligned} E_1(s) &:= \int_0^\infty (\partial_Y^2 \mathcal{L}_U V)^2 w_1, \\ D_1(s) &:= \int_0^\infty \frac{(\partial_Y^3 \mathcal{L}_U V)^2}{U} w_1 + \int_0^\infty \frac{(\partial_Y^2 \mathcal{L}_U V)^2}{U^2} w_1. \end{aligned}$$

Then there exist universal constants $\bar{a}, \bar{c} > 0$, such that for all $a \in]0, \bar{a}[$, for all $\alpha < 6 - (11 - a)\beta_1$, for m_1 large enough, there exists $S_0, H_1 > 0$ depending on $M_1, M_2, c, \beta_1, m_1$, and a such that if $s_0 \geq \max(S_0, J^4)$,

$$E_1(s) \leq H_1(1 + E_1(s_0)s_0^\alpha)s^{-\alpha}, \quad \int_{s_0}^{s_1} s^\alpha D_1(s) ds \leq H_1(1 + E_1(s_0)s_0^\alpha) \quad \forall s \in [s_0, s_1].$$

Remark 2.10. The weight Y^{-a} in w_1 has two different roles. On the one hand, we gain a bit of decay in the remainder terms. On the other hand, we are able to control, through a simple Cauchy-Schwarz inequality, quantities of the type

$$\int_0^Y \frac{\partial_Y^2 \mathcal{L}_U V}{U^2}$$

by the diffusion term.

Remark 2.11. Notice that if we take β_1 such that

$$\frac{1}{4} < \beta_1 < \frac{1}{4} \frac{11}{11-a},$$

then we can choose α so that $\alpha > 13/4$. We will make this choice in the final energy estimates, and we will use the corresponding decay of E_1 when we apply the maximum principle.

Differentiating (2.19) with respect to Y and taking the trace of the equation at $Y = 0$, we obtain in particular

Lemma 2.12. For all $s \geq s_0$, we have

$$\partial_Y \mathcal{L}_U^2 V|_{Y=0} = -\frac{1}{2}(b_s + b^2).$$

We then derive an equation on $\mathcal{L}_U^2 V$. Applying once more the commutator result of Lemma 2.8, we deduce that $\mathcal{L}_U^2 V$ satisfies the following equation:

$$\partial_s \mathcal{L}_U^2 V + 3b \mathcal{L}_U^2 V + \frac{b}{2} Y \partial_Y \mathcal{L}_U^2 V - \mathcal{L}_U^3 V = \mathcal{L}_U^2 \mathcal{R} + \mathcal{C}[\partial_Y^2 \mathcal{L}_U V] + \mathcal{L}_U \mathcal{C}[\partial_Y^2 V]. \quad (2.20)$$

In order to have a trace estimate, and also to have nice positivity properties for the diffusion term, we will also need to use estimates on $\partial_Y^2 \mathcal{L}_U^2 V$. We therefore define the energy

$$E_2(s) := \int_0^\infty (\partial_Y^2 \mathcal{L}_U^2 V)^2 w_2,$$

together with the dissipation terms

$$D_2(s) := \int_0^\infty \frac{(\partial_Y^3 \mathcal{L}_U^2 V)^2}{U} w_2 + \int_0^\infty \frac{(\partial_Y^2 \mathcal{L}_U^2 V)^2}{U^2} w_2,$$

where the weight w_2 is defined by $w_2 = Y^{-a}(1+s^{-\beta_2}Y)^{-m_2}$ for some parameters $\beta_2 \in]1/4, \beta_1[$, $m_2 \gg m_1$ sufficiently large. The parameter a is the same as the one in Proposition 2.9.

We then claim that we have the following estimate:

Proposition 2.13. *Assume that the hypotheses of Proposition 2.9 are satisfied. Let $C_0 := \max(E_1(s_0)s_0^{13/4+\eta}, E_2(s_0)s_0^5)$ for some $\eta > 0$ such that $13/4 + \eta < 6 - (11 - a)\beta_1$. Then there exists a universal constant \bar{a} , such that for all $a \in]0, \bar{a}[$, for a suitable choice of β_2, m_2 , there exist $S_0, H_2 > 0$ (depending on $a, \beta_i, m_i, M_1, M_2$) such that if $s_0 \geq \max(S_0, J^4, C_0^8)$,*

$$E_2(s) \leq H_2(1 + C_0) \exp(H_2(1 + C_0))s^{-5} \quad \forall s \in [s_0, s_1]. \quad (2.21)$$

Let us now go back to the definition of the semi-norm \mathcal{N} . We need a new type of trace estimate, taking advantage of the fact that E_2 has a stronger decay than E_1 (notice that the sole decay of E_1 is not sufficient to close the bootstrap argument, since $E_1 \lesssim s^{-13/4-\eta}$ and we need $|b_s + b^2| \lesssim s^{-2-\eta/2}$, while $13/4 < 4$).

We will use the following trace estimate, which is proved in the Appendix:

Lemma 2.14. *There exists a universal constant \bar{C} , such that for all $L \geq 1$, for any smooth function f ,*

$$|f(0)|^2 \leq \bar{C} \left(L^{1+a} \int_0^L (\partial_Y f)^2 Y^{-a} dY + \frac{1}{L^{3-a}} \int_0^L |f(Y)|^2 (Y + Y^2) Y^{-a} dY \right).$$

In particular, taking $f = \partial_Y \mathcal{L}_U^2 V$ and $L = s^{1/4}$, under the assumptions of Proposition 2.13,

$$|b_s + b^2|^2 \leq \bar{C} \left(s^{\frac{1+a}{4}} E_2 + s^{-\frac{3-a}{4}} \int_0^{s^{1/4}} U (\partial_Y \mathcal{L}_U^2 V)^2 \right).$$

Let us now go back to the definition of the semi-norm \mathcal{N} . According to the above Lemma, we can take for instance

$$\mathcal{N}(V) := \left(s^{\frac{1+a}{4}} E_2 + s^{-\frac{3-a}{4}} \int_0^\infty U (\partial_Y \mathcal{L}_U^2 V)^2 \tilde{w}_1 \right)^{1/2},$$

where $\tilde{w}_1 := Y^{-a}(1 + s^{-\beta_1}Y)^{-m_1-2} = w_1(1 + s^{-\beta_1}Y)^{-2}$, so that $\mathcal{N}(V) \geq \bar{C}|b_s + b^2|$ for some universal constant \bar{C} .

However there remains to prove that with this definition, $\mathcal{N}(V)$ is sufficiently small. According to the above Lemma, we also need to find a bound for $\int_0^\infty U (\partial_Y \mathcal{L}_U^2 V)^2 \tilde{w}_1$. We claim that we have the following estimate:

Lemma 2.15. *Assume that the assumptions of Proposition 2.9 are satisfied. Let $P > 0$ be arbitrary. There exist $S_0, H_1 > 0$ depending on M_1, M_2, β_1, m_1 , and a , and a function ρ such that $\int_{s_0}^{s_1} \rho(s) s^{-1/2} ds \leq 1$, such that if m_1 is large enough (depending on P) and if $s_0 \geq \max(S_0, J^4)$,*

$$\int_0^\infty U (\partial_Y \mathcal{L}_U^2 V)^2 \tilde{w}_1 \leq H_1 s^{(3-a)\beta_1} D_1 + s^{-1} E_1 + s^{-P} \rho(s). \quad (2.22)$$

Gathering Lemmas 2.14 and 2.15 and using Proposition 2.9 and Proposition 2.13, we find that if $s_0 \geq \max(S_0, J^4, C_0^8)$,

$$|b_s + b^2|^2 \leq H_2(1 + C_0) \exp(H_2(1 + C_0)) s^{-\frac{19-a}{4}} + H_1 s^{(3-a)(\beta_1 - \frac{1}{4})} D_1 + s^{-P} \rho(s).$$

Recall that $\int_{s_0}^{s_1} s^\alpha D_1(s) ds \leq H_1(1 + C_0)$ for $s_0 \geq \max(S_0, J^4)$ and for $\alpha = 13/4 + \eta < 6 - (11 - a)\beta_1$. A short computation¹ shows that we can choose β_1 and a so that

$$\frac{13}{4} + (3 - a) \left(\beta_1 - \frac{1}{4} \right) < 6 - (11 - a)\beta_1.$$

We obtain eventually that

$$|b_s + b^2| \leq \bar{C} \mathcal{N}(V),$$

and

$$\int_{s_0}^{s_1} s^{13/4} \mathcal{N}(V(s))^2 ds \leq H(1 + C_0) \exp(H(1 + C_0)),$$

for some constant H depending on $a, \beta_i, m_i, M_1, M_2$, provided $s_0 \geq \max(S_0, J^4, C_0^8)$. Thus b satisfies the assumptions of Lemma B.1 with $\gamma = 13/4 > 3$.

We gather the estimates of Propositions 2.9 and 2.13 and Lemma 2.14 in the following Theorem:

Theorem 2. *Let $a \in]0, \bar{a}[$, and choose the parameters β_1 and β_2 such that*

$$\frac{1}{4} < \beta_2 < \beta_1 < \frac{1}{4} \inf \left(\frac{11}{11 - a}, \frac{14 - a}{14 - 2a} \right) \quad (2.23)$$

and $m_2 \gg m_1 \gg 1$.

Let $\eta = \eta(a, \beta_1)$ such that $0 < \eta < (3 - a)(\beta_1 - 1/4)$.

Assume that the following assumptions are satisfied:

¹It is enough to choose

$$\beta_1 < \frac{1}{4} \left(1 + \frac{a}{14 - 2a} \right).$$

- There exists a constant $J > 0$ such that

$$\int_{s_0}^{s_1} s^{13/4} |b_s + b^2|^2 ds \leq J;$$

- $(1 - \bar{\epsilon})/s \leq b \leq (1 + \bar{\epsilon})/s$ for $s \in [s_0, s_1]$ for some small enough constant $\bar{\epsilon}$ (say $\bar{\epsilon} = 1/50$);
- There exist constants M_1, M_2, c independent of s , such that

$$\begin{aligned} -M_1 &\leq U_{YY} \leq 1 \quad \forall Y \geq 0, \\ 1 - M_2 b Y^2 &\leq U_{YY} \quad \forall Y \in [0, cs^{1/3}]. \end{aligned}$$

- There exists a constant C_0 , independent of s_0 and λ_0 , such that for some $\eta > 0$

$$E_2(s_0) s_0^5 + E_1(s_0) s_0^{\eta+13/4} \leq C_0.$$

Then there exists constants H, S_0 , depending on $a, m_1, m_2, \beta_1, \beta_2, M_1$ and M_2 , such that for all $s_0 \geq \max(S_0, J^4, C_0^8)$,

$$\int_{s_0}^{s_1} s^{13/4} |b_s + b^2|^2 \leq \exp(H(1 + C_0)).$$

In particular, setting $J' := \exp(H(1 + C_0))$, we have

$$s_0 \geq \max(S_0, J^4, C_0^8) \Rightarrow \int_{s_0}^{s_1} s^{13/4} |b_s + b^2|^2 \leq J',$$

and the constant J' is independent of J .

2.4 Construction of sub- and super-solutions

The other ingredient in the proof of Theorem 1 is the use of the maximum principle in order to control the growth of U and its derivatives on the one hand, and the size of some non-linear terms on the other hand. Indeed, the assumptions of Propositions 2.9 and 2.13 require estimates on $\partial_Y^2 U$. These estimates are obtained by careful applications of a comparison principle. We emphasize that this principle is not applied to equation (2.3) directly, but rather to an equation derived from (2.3) after a non-linear change of variables. More precisely, we use the von Mises variables

$$\psi := \int_0^Y U, \quad W := U^2. \quad (2.24)$$

The tangential variable remains s , the normal variable is now ψ (instead of Y), and the new unknown function is W . This change of variables transforms (2.3) into a non-linear transport-diffusion equation which is more suited for maximum principle techniques, namely

$$\partial_s W - 2bW + \frac{3b}{2} \psi \partial_\psi W - \sqrt{W} \partial_\psi^2 W = -2 \quad (2.25)$$

Equation (2.25) is the third and last form of equation (2.3). We refer to [29] and to section 3 of the present paper for more details. Since the new equation is local and parabolic, it enjoys maximum principle properties. Therefore we construct sub- and super-solutions for W and

its derivatives and thereby derive estimates on W . These estimates are then translated in terms of the former variables s, Y .

One of the key points lies in the construction of a sub-solution for W (see Lemma 3.6). Actually, the Sobolev estimates of Proposition 2.9 provide a very good pointwise control of U up to $Y \sim s^{\beta_1} \gg s^{1/4}$, but this control degenerates for $Y \gtrsim s^{\beta_1}$. Hence it is sufficient to construct sub-solutions for $Y \gtrsim s^{1/4}$, or equivalently (since $\psi \sim Y^3/6$ for $Y \gg 1$) for $\psi \gtrsim s^{3/4}$. On this zone, the sub-solutions will be linear combinations of powers of ψ . Furthermore, we define a *regularized modulation rate* \tilde{b} , whose role is to remove some oscillations from b while keeping the same asymptotic behavior. We take

$$\tilde{b}_s + b\tilde{b} = 0, \quad \tilde{b}|_{s=s_0} = \frac{1}{s_0}.$$

The sub-solutions for W are defined by

$$\underline{W} := \frac{(6\psi)^{4/3}}{4} - A\psi^{7/3}\tilde{b}^{5/4},$$

where A is chosen sufficiently large. Notice that the main order term $(6\psi)^{4/3}/4$ is the same as in the original solution. It corresponds to the main order term $Y^2/2$ in $U(s, Y)$.

The regularized modulation rate \tilde{b} is also used in the construction of a sub-solution for $U_{YY} - 1$ (see Lemma 3.7).

The final result is the following (we refer to Remark 2.11 regarding the assumption on E_1):

Proposition 2.16. *Assume that there exist constants $J > 0$, $\eta > 0$ and $\epsilon > 0$ such that for all $s \in [s_0, s_1]$, assumption (2.12) is satisfied. Assume furthermore that there exists a constant M_0 such that*

$$\begin{aligned} -M_0 \inf(1, s_0^{-1}Y^2) \leq U_{YY}(s_0, Y) - 1 \leq 0 \quad \forall Y > 0, \\ \lim_{Y \rightarrow \infty} U(s_0, Y) \leq M_0 s_0, \end{aligned}$$

and that there exists $C_1 > 0$ such that

$$E_1(s) \leq C_1 s^{-13/4} \quad \forall s \in [s_0, s_1].$$

Then there exist universal constants $\bar{M}, \bar{C} > 0$, and S_0 depending on C_1, β_1, m_1 , such that if $s_0 \geq \max(S_0, \bar{C}(J\epsilon^{-2})^{1/2\eta})$, then, setting $M' = \bar{M} \max(1, M_0)$,

$$\begin{aligned} -M'bY^2 \leq U_{YY}(s, Y) - 1 \leq 0 \quad \forall s \in [s_0, s_1], \quad \forall Y \in [0, s^{1/3}], \\ \text{and } -M' \leq U_{YY}(s, Y) - 1 \leq 0 \quad \forall s \in [s_0, s_1], \quad \forall Y \geq s^{1/3}. \end{aligned}$$

Notice that the above estimates are precisely the ones that are required in Proposition 2.9.

2.5 Bootstrap argument

The bootstrap argument consists in bringing together Theorem 2 on the one hand, and Proposition 2.16 on the other. In the rest of this section, we will assume that $U(s_0)$ satisfies

$$\begin{aligned} E_1(s_0) \leq C_0 s_0^{-13/4-\eta/2}, \quad E_2(s_0) \leq C_0 s_0^{-5}, \\ \left| b(s_0) - \frac{1}{s_0} \right| \leq \frac{\bar{\epsilon}}{2s_0}, \end{aligned} \quad (2.26)$$

$$\text{and } -M_0 \inf(1, s_0^{-1}Y^2) \leq U_{YY}(s_0, Y) - 1 \leq 0 \quad \forall Y > 0, \quad \lim_{Y \rightarrow \infty} U(s_0, Y) \leq M_0 s_0.$$

where C_0, M_0 are constants independent of s_0 , and η is such that $\eta - 3/4 < 2 - (11 - a)\beta_1$. Without loss of generality, we also assume that $M_0 \geq 1$. Such an initial data is “well-prepared” in the sense that it is close to the blow-up profile.

Assumption (2.26) involves three different types of estimates. In order to propagate these estimates, we will apply three different results:

1. The energy estimates from Propositions 2.9 and 2.13, which are gathered in Theorem 2;
2. The maximum principle estimates from Proposition 2.16;
3. Lemma B.1 on the modulation rate b .

Note that the maximum principle will propagate the third estimate of (2.26) without improving it (in fact, we will change the constant M_0 into $\bar{M}M_0$); however the energy estimates will transform $\eta/2$ into η , and will therefore improve the estimates on E_1 .

The argument goes as follows: let S_0, H be the constants from Theorem 2 with $\alpha = 13/4 + \eta$ and $M_1 = M_2 = \bar{M}M_0$ (recall that S_0 depends in particular on β_1, β_2, η and a). Let $J := 2\exp(H(1 + C_0))$. Assume that $s_0 \geq \max(S_0, \bar{C}J^4, C_0^8)$ for the large universal constant \bar{C} from Proposition 2.16.

If the initial data satisfies (2.26), by continuity, there exists $s_1 > s_0$ such that for all $s \in [s_0, s_1]$,

$$E_1(s) \leq 2H_1(1 + C_0)s^{-13/4 - \frac{\eta}{2}}, \quad \left| b(s) - \frac{1}{s} \right| \leq \frac{\bar{\epsilon}}{s}, \quad \int_{s_0}^{s_1} |b_s + b^2|^2 s^{13/4} ds \leq J. \quad (2.27)$$

Then for all $s \in [s_0, s_1]$, according to Proposition 2.16 with $C_1 := 2H_1(1 + C_0)$, we infer that up to choosing a larger S_0 (depending on H_1 and C_0),

$$\begin{aligned} -\bar{M}M_0 &\leq U_{YY} \leq 1 \quad \forall Y \geq 0, \\ 1 - \bar{M}M_0 b Y^2 &\leq U_{YY} \quad \forall Y \in [0, s^{1/3}]. \end{aligned}$$

The assumptions of Propositions 2.9 and 2.13 are satisfied (with $M_1 = M_2 = \bar{M}M_0$), and we infer that if $s_0 \geq \max(S_0, \bar{C}J^4, C_0^8)$, for all $s \in [s_0, s_1]$,

$$E_1(s) \leq H_1(1 + C_0 s_0^{\eta/2}) s^{-13/4 - \eta}, \quad E_2(s) \leq H_2(1 + C_0) \exp(H_2(1 + C_0)) s^{-5}.$$

We have in particular for all $s \in [s_0, s_1]$

$$E_1(s) \leq H_1(1 + C_0) s^{-13/4 - \eta/2},$$

and using Lemmas 2.14 and 2.15,

$$\int_{s_0}^{s_1} |b_s + b^2|^2 s^{13/4} ds \leq \exp(H(1 + C_0)) = \frac{J}{2}.$$

Using Lemma B.1 in the Appendix, we infer that for all $s \in [s_0, s_1]$,

$$\left| b(s) - \frac{1}{s} \right| \leq \frac{1 + \bar{\epsilon}}{1 - \bar{\epsilon}} \left| \frac{1}{s_0} - b(s_0) \right| \frac{s_0^2}{s^2} + \frac{1 + \bar{\epsilon}}{(1 - \bar{\epsilon})^2} s^{-9/8} \sqrt{\frac{2J}{7}}.$$

Without loss of generality, we can always assume that $\bar{\epsilon}, \eta, s_0$ are such that

$$\frac{1 + \bar{\epsilon}}{1 - \bar{\epsilon}} \leq \frac{5}{4}, \quad \frac{1 + \bar{\epsilon}}{(1 - \bar{\epsilon})^2} \sqrt{\frac{2J}{7}} s_0^{-1/8} \leq \frac{\bar{\epsilon}}{4}.$$

Then for all $s \in [s_0, s_1]$, we have

$$\left| b(s) - \frac{1}{s} \right| \leq \frac{7\bar{\epsilon}}{8s}. \quad (2.28)$$

Gathering (2.27) and (2.28), we infer that

$$s_1 := \inf \{ s \geq s_0, E_1(s) = 2H_1(1 + C_0)s^{-13/4 - \eta/2} \text{ or } |b(s) - 1/s| = \bar{\epsilon}/s \\ \text{or } \int_{s_0}^s \tau^{13/4} |b_\tau + b^2|^2 d\tau = J \} = +\infty.$$

As a consequence, we infer that for some constant J depending only on C_0 and M_0 , if $s_0 \geq \max(S_0, C J^4, C_0^8)$,

$$\int_{s_0}^{\infty} s^{13/4} |b_s + b^2|^2 ds \leq J, \quad \left| b - \frac{1}{s} \right| \leq \frac{\bar{\epsilon}}{s}.$$

We therefore obtain the following Theorem in the rescaled variables (s, Y) :

Theorem 3. *Let η be such that $0 < \eta < (3 - a)(\beta_1 - 1/4)$, where β_1 satisfies (2.23). Assume that U_0 satisfies the hypotheses (2.26), and consider the solution U of equation (2.3) with $U(s_0, Y) = U_0$. Then there exists a constant $S_0 > 0$, depending on η, C_0, M_0 , such that if $s_0 \geq S_0$, then for all $s \geq s_0$,*

$$\int_{s_0}^{\infty} s^{13/4} |b_s + b^2|^2 ds < +\infty, \quad \left| b - \frac{1}{s} \right| \leq \frac{\bar{\epsilon}}{s}. \quad (2.29)$$

Let us now go back to the original variables and prove Theorem 1. First, we set

$$\lambda_0 := \partial_y u_0|_{y=0}, \quad b_0 := -\lambda_0^2 \partial_y^4 u_0|_{y=0}, \quad s_0 := b_0^{-1}.$$

The assumption (H1) entails that

$$c_0^{-1} \lambda_0^2 \leq s_0^{-1} \leq c_0 \lambda_0^2.$$

Assumption (H2) implies

$$-M_0 \inf(1, s_0^{-1} Y^2) \leq U_{YY}(s_0, Y) - 1 \leq 0 \quad \forall Y > 0, \\ U(s_0, Y) \leq M_0 \lambda_0^{-2} \leq M_0 c_0 s_0$$

for a suitable constant M_0 . Of course, without loss of generality, we can assume that $M_0 \geq 1$. Furthermore, assumption (H3) becomes, in the rescaled variables and after a few easy computations,

$$V(s_0, Y) = O(s_0^{-9/4})(Y^7 + c_8 Y^8) + O(s_0^{-3}(Y^{10} + Y^{11}))$$

for $Y \leq s_0^{2/7}$. Here the constant c_8 is defined so that

$$\partial_Y \mathcal{L}_U^2 V(s_0) = O(s_0^{-9/4}) + O(s_0^{-3} Y^2).$$

It can be easily checked that these assumptions ensure that U is a well-prepared initial data. As a consequence, if s_0 is large enough (i.e. if λ_0 is small enough), (2.29) holds. Using Lemma 2.3, we infer that $x^* < +\infty$ and that $\lambda(x) \sim C\sqrt{x^* - x}$. Theorem 1 follows.

Furthermore, we deduce from the maximum principle estimates some pointwise control on u . Indeed, we have

$$Y + \frac{Y^2}{2} - \frac{M_2}{12}bY^4 \leq U(s, Y) \leq Y + \frac{Y^2}{2} \quad \text{for } 0 \leq Y \leq cs^{1/3}.$$

Going back to the original variables, we find that there exist constants C, c such that

$$\lambda(x)y + \frac{y^2}{2} - Cy^4 \leq u(x, y) \leq \lambda(x)y + \frac{y^2}{2} \quad \forall y \leq (x^* - x)^{1/6}.$$

Remark 2.17 (Comparison with the result by Caffarelli and E). *Let us now plug the change of variables in the result announced by Caffarelli and E in [4] into the asymptotic expansion above. We recall that*

$$u^\mu(\xi, z) = \frac{1}{\mu^{1/2}}u(x^* - \xi\mu, \mu^{1/4}z).$$

It follows that in the zone $z \leq \mu^{-1/12}\xi^{1/6}$, $\xi \lesssim 1$,

$$u^\mu(\xi, z) = O(\mu^{1/4}\sqrt{\xi})z + \frac{z^2}{2} + O(\mu^{1/2}z^4) \rightarrow \frac{z^2}{2} \quad \text{as } \mu \rightarrow 0.$$

2.6 Organization of the rest of the paper

The rest of the paper is dedicated to the proof of Theorem 3, or more specifically, to the proofs of Proposition 2.9, Proposition 2.13 and Proposition 2.16. Since the maximum principle estimates are easier to derive than the energy estimates, we start with the proof of Proposition 2.16 in section 3. We then lay the ground for the derivation of the energy estimates by proving several important intermediate results in section 4. Eventually, we prove Proposition 2.9 and Proposition 2.13 in section 5.

Let us also explain here the order in which the parameters are chosen. We first pick $a \in (0, \bar{a})$, where \bar{a} is the universal constant in Proposition 2.9. We then choose $\beta_1 > \beta_2$ satisfying (2.23), and $\eta > 0$ such that $\eta - \frac{3}{4} < 2 - (11 - a)\beta_1$. We then pick m_1, m_2 large enough and such that $m_2 \gg m_1$. Eventually, we take s_0 large, depending on all other parameters.

Notation

We will use indifferently f_Y and $\partial_Y f$ to denote the Y derivative of an arbitrary function f . The constants with a bar ($\bar{a}, \bar{C}, \bar{M}, \bar{\epsilon}$) denote universal constants, that do not depend on any of the parameters. All constants with a zero subscript (M_0, C_0, s_0) refer to the initial data. Constants involving the letter M (M_0, M_1, M_2, \bar{M}) are related to the maximum principle.

3 Derivation of L^∞ estimates and construction of sub and super solutions

This section is devoted to the proof of Proposition 2.16, which consists in the derivation of pointwise estimates on U, U_Y and U_{YY} , provided b satisfies the assumptions of Lemma B.1 and $E_1(s) = O(s^{-13/4})$. Throughout this section, we will use the von Mises formulation of the rescaled Prandtl equation, namely (2.24)-(2.25). The idea is to use the maximum and comparison principles for these equations (see Lemmas 2.1.3 and 2.1.4 in [29]), together with Sobolev estimates coming from the bound on $E_1(s)$.

Let us first recall some useful formulas regarding the von Mises formulation of the equation in the original variables and in the rescaled variables. If u is the solution of (P), we set

$$\phi(x, y) = \int_0^y u, \quad w = u^2.$$

We recall that (P) is equivalent to the following equation, written in the variables (x, ϕ)

$$w_x - \sqrt{w} \partial_\phi^2 w = -2. \quad (3.1)$$

Furthermore, notice that if W, ψ are defined by (2.24)

$$\psi(s, Y) = \lambda(x(s))^{-3} \phi(x(s), \lambda(x(s))Y), \quad W(s, \psi) = \lambda(x(s))^{-4} w(x(s), \lambda^3 \psi).$$

It follows that some qualitative properties of equation (2.25) (growth with respect to ϕ , local bounds) can be inherited directly from equation (3.1). More precisely, we have the following result:

Lemma 3.1. *Let $w_0 \in C^{3,\alpha}(\mathbb{R}_+)$ such that $w_0(0) = 0$ and $w_0'(0) > 0$. Assume that w_0 is increasing. Then for all $x \in [0, x^*[$, $w(x, \phi)$ is increasing with respect to ϕ . Furthermore, for any $X \in [0, x^*[$, there exists $C_X > 0$ such that for all $x \in [0, X]$,*

$$\begin{aligned} |w_\phi(x, \phi)| &\leq C_X \quad \forall \phi \geq 0, \\ |\partial_\phi^2 w(x, \phi)|, |\partial_\phi^3 w(x, \phi)| &\leq C_X \quad \forall \phi \geq 1. \end{aligned}$$

As a consequence, W is increasing in ψ (or equivalently, U is increasing in Y) for all $s \in [s_0, s_1]$, and

$$\lim_{\psi \rightarrow \infty} W_\psi = \lim_{\psi \rightarrow \infty} W_{\psi\psi} = 0 \quad \forall s \in [s_0, s_1].$$

Proof. The bounds on $w_\phi, w_{\phi\phi}$ are explicitly written in [29] (see Lemmas 2.1.9 and 2.1.11). The bound on $\partial_\phi^3 w$ follows from the same arguments as [29, Lemma 2.1.11], writing down the equation on w_ϕ . Since $\lim_{\phi \rightarrow \infty} w(x, \phi) = \bar{U}(x)^2$, it follows that $\lim_{\phi \rightarrow \infty} w_\phi = 0$ for all $x \in (0, x^*)$. Therefore we also have $\lim_{\phi \rightarrow \infty} w_{\phi\phi} = 0$. Whence $\lim_{\psi \rightarrow \infty} W_\psi = \lim_{\psi \rightarrow \infty} W_{\psi\psi} = 0$.

Furthermore, the equation satisfied by w_ϕ is

$$\partial_x w_\phi - \frac{w_\phi}{2\sqrt{w}} \partial_\phi w_\phi - \sqrt{w} \partial_{\phi\phi} w_\phi = 0,$$

with boundary conditions $w_\phi|_{x=0} = w_0'(\phi) \geq 0$, $\lim_{\phi \rightarrow \infty} w_\phi = 0$, and $w_\phi|_{\phi=0} = 2\lambda(x) > 0$ for all $x \in (0, x^*)$. According to the maximum principle, we have $w_\phi \geq 0$ in $(0, x^*) \times (0, \infty)$. Hence $W_\psi \geq 0$, and W is increasing in ψ . □

3.1 Uniform bounds on $\partial_{YY}U$

The first step of the proof of Proposition 2.16 is the derivation of uniform L^∞ bounds on $\partial_{YY}U$. The result we prove in this paragraph is the following

Lemma 3.2. *Let U be a solution of (2.3) on $(s_0, s_1) \times (0, +\infty)$ such that $U_{Y|Y=0} = 1$ for all $s \in [s_0, s_1]$, and such that U is strictly increasing in Y for all s , with $\lim_{Y \rightarrow \infty} U(s, Y) = U_\infty(s) < +\infty$.*

Assume that there exists M_2 such that

$$-M_2 \leq \partial_{YY}U(s_0, Y) - 1 \leq 0 \quad \forall Y > 0,$$

and that

$$\partial_{YY}U(s_0, Y) = 1 - 12a_4b_0Y^2 + O(Y^5) \text{ for } Y \ll 1. \quad (3.2)$$

Then

$$-\max(M_2, 1) \leq \partial_{YY}U(s, Y) - 1 \leq 0 \quad \forall Y > 0 \quad \forall s \geq s_0.$$

Remark 3.3. *Assumption (3.2) is a compatibility condition at a high order at $s = s_0$. It is propagated by the equation.*

Proof. We rely on the equation on W in the (s, ψ) variables. We recall that $\partial_Y U = \partial_\psi W/2$, and therefore

$$\partial_{YY}U(s, Y) = \frac{1}{2} \sqrt{W(s, \psi(s, Y))} \partial_{\psi\psi} W(s, \psi(s, Y)) \quad \forall s \geq s_0, \quad \forall Y > 0.$$

Therefore we derive estimates on the quantity

$$F(s, \psi) := \sqrt{W} \partial_{\psi\psi} W - 2.$$

Notice that the assumptions on U imply that

$$-2M_2 \leq F(s_0, \psi) \leq 0 \quad \forall \psi > 0.$$

On $\{Y = 0\}$, we have $U_{YY} = 1$, and therefore $F|_{\psi=0} = 0$. Using Lemma 3.1, we also have $\lim_{\psi \rightarrow \infty} F(s, \psi) = -2$.

Furthermore, F satisfies

$$\partial_s F = \frac{\partial_s W}{2\sqrt{W}} \partial_{\psi\psi} W + \sqrt{W} \partial_{\psi\psi} \partial_s W.$$

Using the equation on W (2.25) and writing $\partial_{\psi\psi} W = (F + 2)/\sqrt{W}$, we infer that

$$\partial_s F = \frac{1}{2W} F(F + 2) + \frac{1}{2\sqrt{W}} \partial_\psi^2 W \left(2bW - \frac{3b}{2} \psi \partial_\psi W \right) + \sqrt{W} \left(-b \partial_\psi^2 W - \frac{3b}{2} \psi \partial_\psi^3 W + \partial_\psi^2 F \right).$$

Gathering all the terms and using the formula

$$\partial_\psi F = \frac{1}{2\sqrt{W}} \partial_\psi W \partial_{\psi\psi} W + \sqrt{W} \partial_\psi^3 W,$$

we obtain eventually

$$\partial_s F - \frac{1}{2W} F(F+2) + \frac{3b}{2} \psi \partial_\psi F - \sqrt{W} \partial_{\psi\psi} F = 0. \quad (3.3)$$

▷ *First step: Lower bound on F and consequences.*

We start with the lower bound, which is easier. Assume that F has an interior minimum F_{min} at some point (s, ψ) for some $s \in (s_0, s_1]$, $\psi > 0$. Then according to equation (3.3), $F_{min}(F_{min} + 2) \leq 0$, and therefore $F_{min} \in (-2, 0)$. Thus $F(s, \psi) \geq \min(-2, \inf F(s_0)) \geq \min(-2, -2M_2)$.

We infer from this lower bound on F some non-degeneracy estimates for W for ψ close to zero. Indeed, it follows from the inequality $U_{YY} \geq -M'_2$ with $M'_2 = \max(M_2 - 1, 0)$, that

$$1 - M'_2 Y \leq U_Y \quad \forall Y > 0.$$

In particular, if $Y \leq Y_0 := (2M'_2)^{-1}$, then $U_Y(s, Y) \geq 1/2$ and $U(s, Y) \geq Y/2$. As a consequence, if $\psi \leq \psi(s, Y_0)$, then $W_\psi \geq 1$. Now, the lower bound on U_{YY} also entails that

$$\psi(s, Y_0) = \int_0^{Y_0} U(s, Y) dY \geq \frac{Y_0^2}{4} = \frac{1}{16M_2'^2}.$$

Hence in particular, for all $s \geq s_0$,

$$\psi \leq \frac{1}{16M_2'^2} \Rightarrow W(s, \psi) \geq \psi \text{ and } W_\psi \geq 1. \quad (3.4)$$

▷ *Second step: Upper bound on F .*

The derivation of the upper-bound is a little more involved. The main difficulty comes from the nonlinear term $F(F+2)/W$, which is also singular near $\psi = 0$. In order to deal with it, we use a bootstrap type argument. Notice first that the preliminary bounds of Lemma 3.1 entail that W is Lipschitz continuous, uniformly in ψ and locally uniformly in s and that $\partial_{\psi\psi} W$, $\partial_\psi^3 W$ are bounded (locally in s , uniformly in ψ) in $s \geq s_0$, $\psi \geq \delta$, for any $\delta > 0$. Considering eventually equation (3.3), we deduce that $\partial_s F$ is bounded in a neighbourhood of $s = s_0$, uniformly in ψ for $\psi \geq \delta$. Furthermore, using assumption (3.2) on $U(s_0)$, both $F(s_0)/W(s_0)$ and $\sqrt{W}(s_0)\partial_{\psi\psi} F(s_0)$ are bounded in a neighbourhood of $Y = 0$, and therefore $\partial_s F|_{s=s_0}$ is bounded in $L^\infty(\mathbb{R}_+)$.

We now set

$$s'_0 := \inf\{s \in [s_0, s_1], \exists \psi > 0, F(s, \psi) \geq 1\}.$$

It follows from the above arguments that $s'_0 > s_0$. On the interval $[s_0, s'_0]$, we have $F(s, \psi) \in [-2 \max(M_2, 1), 1]$. As a consequence, we multiply (3.3) by $F_+ p$, where $p \in C^\infty(\mathbb{R})$ is a non-increasing weight function such that $p \equiv 1$ for ψ close to zero and $p(\psi) = O(\psi^{-k})$ for some $k > 1$ for $\psi > 1$, with $|p'|/p \in L^\infty$, $p''/p \in L^\infty$. Since $F_+|_{\psi=0} = 0$, we obtain, as long as $s \leq s'_0$,

$$\frac{d}{ds} \int_{\mathbb{R}_+} F_+^2 p + \int_{\mathbb{R}_+} \sqrt{W} (\partial_\psi F_+)^2 p - \frac{1}{2} \int_{\mathbb{R}_+} F_+^2 \partial_\psi^2 (\sqrt{W} p) \leq \frac{3}{2} \int_{\mathbb{R}_+} \frac{F_+^2}{W} p + \frac{3b}{4} \int_{\mathbb{R}_+} F_+^2 p.$$

An easy computation gives

$$\partial_\psi^2 (\sqrt{W} p) = \frac{F+2}{2W} p - \frac{1}{4} \frac{W_\psi^2}{W^{3/2}} p + \frac{W_\psi}{\sqrt{W}} p' + \sqrt{W} p''.$$

Using the assumptions on p , the upper-bound $\sqrt{W} \leq U_\infty(s)$ and the bound on F for $s \leq s'_0$, we deduce eventually that

$$\begin{aligned} \frac{d}{ds} \int_{\mathbb{R}_+} F_+^2 p + \int_{\mathbb{R}_+} \sqrt{W} (\partial_\psi F_+)^2 p + \frac{1}{2} \int_{\mathbb{R}_+} F_+^2 \frac{W_\psi}{\sqrt{W}} |p'| + \frac{1}{8} \int_{\mathbb{R}_+} F_+^2 \frac{W_\psi^2}{W^{3/2}} p \\ \leq C \int_{\mathbb{R}_+} \frac{F_+^2}{W} p + C(1 + U_\infty(s)) \int_{\mathbb{R}_+} F_+^2 p. \end{aligned}$$

The second term in the right-hand side will be handled thanks to a Gronwall type argument. The singularity of the first term in the right-hand side will be absorbed in the dissipation term. Indeed, let us first decompose the integral into two pieces depending on the value of W . First,

$$\int_{\mathbb{R}_+} \mathbf{1}_{W \geq \inf(1, (5M'_2)^{-2})} \frac{F_+^2}{W} p \leq C \int_{\mathbb{R}_+} F_+^2 p,$$

and as before, that part can be handled thanks to a Gronwall type argument. We thus focus of the values of W below $\inf(1, (5M'_2)^{-2})$. In that case, according to (3.4), we have $\psi \leq \inf(1, (5M'_2)^{-2}) =: \psi_0$ and

$$\frac{W_\psi}{W} \geq W_\psi \geq 1.$$

Let us choose p so that $p(\psi) = 1$ for $\psi \in [0, 1]$. We deduce that there exists an explicit constant C such that for all $\psi \in (0, \psi_0)$,

$$\sqrt{W} (\partial_\psi F_+)^2 + \frac{1}{8} F_+^2 \frac{W_\psi^2}{W^{3/2}} \geq C \left(\partial_\psi \left(W^{1/4} F_+ \right) \right)^2.$$

Therefore, using the Hardy inequality, there exists a constant C such that

$$D(s) := \int_0^{\psi_0} \sqrt{W} (\partial_\psi F_+)^2 p + \frac{1}{2} \int_0^{\psi_0} F_+^2 \frac{W_\psi}{\sqrt{W}} |p'| + \frac{1}{8} \int_0^{\psi_0} F_+^2 \frac{W_\psi^2}{W^{3/2}} p \geq C \int_0^{\psi_0} \frac{W^{1/2} F_+^2}{\psi^2} p.$$

Using once again the non-degeneracy of W for ψ close to zero (see (3.4)), we infer that up to choosing a smaller ψ_0 ,

$$\int_{\mathbb{R}_+} \mathbf{1}_{W \leq \inf(1, (5M'_2)^{-2})} \frac{F_+^2}{W} p \leq \frac{1}{2} D(s).$$

Eventually, we obtain

$$\frac{d}{ds} \int_{\mathbb{R}_+} F_+^2 p \leq C(1 + U_\infty(s)) \int_{\mathbb{R}_+} F_+^2 p \quad \forall \in [s_0, s'_0].$$

Now, since $F|_{s=s_0} \leq 0$, we have $F|_{s=s_0} \equiv 0$. The Gronwall Lemma implies that $F_+ \equiv 0$ for $s \leq s'_0$. Therefore $F(s, \psi) \leq 0 < 1$ for all $s \leq s'_0$. It follows that $s'_0 = s_1$, and thus $F(s, \psi) \leq 0$ for all $s \in [s_0, s_1]$ and for all $\psi > 0$. □

Under the assumptions of Lemma 3.2, we therefore have

$$\begin{aligned} \sup \left(Y - M'_2 \frac{Y^2}{2}, 0 \right) \leq U(s, Y) \leq Y + \frac{Y^2}{2}, \quad \forall Y > 0, \forall s \geq s_0. \\ \sup(1 - M'_2 Y, 0) \leq U_Y \leq 1 + Y \end{aligned} \quad (3.5)$$

Notice that these estimates are independent of s , and that the constant M'_2 depends only on M_2 .

3.2 Construction of sub and super solutions for W

We now derive pointwise estimates on U , which will be used in the last paragraph of this section to obtain a refined lower bound on $\partial_{YY}U$. We distinguish between different zones:

- On the zone $Y \ll s^{\beta_1}$, where $\beta_1 > 1/4$ is the parameter entering the definition of w_1 (see Proposition 2.9), the energy estimate $E_1(s) \lesssim s^{-13/4}$ actually provides a very good pointwise estimate of U . However, this estimate degenerates when $Y \gtrsim s^{\beta_1}$. Let us emphasize that we do need estimates on U, U_Y, U_{YY} on the zone $Y \geq s^{\beta_1}$ in order to prove Proposition 2.9 and therefore close the bootstrap argument.
- On the zone $Y \geq Cs^{1/4}$ for some large enough constant C , which corresponds to $\psi \gtrsim s^{3/4}$, we construct sub and super solutions for U (or rather, for W) by using maximum principle arguments. Note that this requires to have a good control of W on the lower boundary of that zone, i.e. on the line $\psi = C's^{3/4}$. This is achieved thanks to the pointwise control coming from the bound on E_1 .

Let us start with the following Lemma:

Lemma 3.4. *Assume that U satisfies the assumptions of Lemma 3.2 and that*

$$E_1(s) \leq C_1 s^{-13/4} \quad \forall s \in [s_0, s_1],$$

where E_1 is defined in Proposition 2.9.

Let $c > 0$ be arbitrary. Then for all $Y \in [0, cs^{1/4}]$, provided s_0 is large enough (depending on β_1, m_1 and c),

$$\begin{aligned} U_{YY}(s, Y) &= 1 - 12a_4bY^2 + O(s^{-13/8}Y^{\frac{5+a}{2}}(1+Y)), \\ U(s, Y) &= Y + \frac{Y^2}{2} - a_4bY^4 + O(s^{-13/8}Y^{\frac{9+a}{2}}(1+Y)). \end{aligned}$$

As a consequence, if s_0 is large enough (depending on β_1, m_1, c and C_1),

$$U_{YY}(s, Y) \geq 1 - \frac{1}{2}bY^2 \quad \forall Y \in [0, cs^{1/4}].$$

Proof. We recall that

$$E_1(s) = \int_0^\infty (\partial_Y^2 \mathcal{L}_U V)^2 w_1,$$

with $w_1 = Y^{-a}(1 + s^{-\beta_1}Y)^{-m_1}$. Therefore, choosing s_0 sufficiently large (depending on β_1, m_1 and c), we have, for all $Y \in [0, cs^{1/4}]$,

$$w_1 \geq Y^{-a} \left(1 + cs_0^{\frac{1}{4} - \beta_1}\right)^{-m_1} \geq \frac{1}{2}Y^{-a}.$$

Using a simple Cauchy-Schwartz inequality, it follows that

$$|\partial_Y \mathcal{L}_U V(s, Y)| = \left| \int_0^Y \partial_Y^2 \mathcal{L}_U V \right| \leq \sqrt{2}Y^{\frac{1+a}{2}} E_1(s)^{1/2} \quad \forall s \in [s_0, s_1], \quad \forall Y \in [0, cs^{1/4}].$$

Integrating twice, we obtain estimates on $\mathcal{L}_U V$ and $\int_0^Y \mathcal{L}_U V$.

Now

$$\partial_Y^2 V = L_U \mathcal{L}_U V = U \mathcal{L}_U V - U_Y \int_0^Y \mathcal{L}_U V.$$

Using (3.5), we infer that

$$|\partial_Y^2 V| \leq \bar{C}(1+Y)Y^{\frac{5+a}{2}}E_1^{1/2},$$

where \bar{C} is an explicit and computable constant. Therefore

$$|V| \leq \bar{C}C_1^{1/2}(1+Y)Y^{\frac{9+a}{2}}s^{-13/8} \quad \forall Y \in [0, cs^{1/4}].$$

Writing $U = U^{\text{app}} + V$ and recalling the definition of U^{app} , we notice that $U^{\text{app}} = Y + Y^2/2 - a_4bY^4 + O(s^{-2}Y^7 + s^{-3}(Y^{10} + Y^{11}))$ for $Y \leq cs^{1/4}$, and we obtain the estimate announced in the statement of the Lemma. Notice that the remainder terms in U^{app} (namely $O(s^{-2}Y^7 + s^{-3}(Y^{10} + Y^{11}))$) are smaller than $(1+Y)Y^{\frac{9+a}{2}}s^{-13/8}$ in the region $Y \leq cs^{1/4}$. \square

Let us now deduce from the above Lemma an asymptotic expansion of W for $1 \ll \psi \lesssim s^{3/4}$. Indeed, a precise pointwise estimate on W is necessary in order to build sub and super solutions.

By definition of W and ψ , we have, in terms of Y ,

$$\begin{aligned} W &= Y^2 + Y^3 + \frac{Y^4}{4} - a_4bY^6 + O(s^{-1}Y^5 + s^{-13/8}Y^{\frac{15+a}{2}}) \\ &= \frac{Y^4}{4} \left(1 + 4Y^{-1} + 4Y^{-2} - 4a_4bY^2 + O(s^{-1}Y + s^{-13/8}Y^{\frac{7+a}{2}}) \right) \\ \psi &= \frac{1}{6}Y^3 \left(1 + 3Y^{-1} - \frac{6}{5}a_4bY^2 + O(s^{-13/8}Y^{\frac{7+a}{2}}) \right). \end{aligned}$$

Above, the notation $A = O(B)$ means the following: there exists a constant C , depending only on C_1 , and there exists $S_0 > 0$ depending on $c, C_1, M_2, \beta_1, m_1$ such that for all $s \geq s_0 \geq S_0$, for all $Y \in [1, cs^{1/4}]$, $|A| \leq CB$.

It follows that

$$\begin{aligned} W(s, \psi) &= \frac{(6\psi)^{4/3}}{4} \left(1 + 4Y^{-1} + 4Y^{-2} - 4a_4\tilde{b}Y^2 + O(s^{-1}Y + s^{-13/8}Y^{\frac{7+a}{2}}) \right) \\ &\quad \times \left(1 + 3Y^{-1} - \frac{6}{5}a_4bY^2 + O(s^{-13/8}Y^{\frac{7+a}{2}}) \right)^{-4/3}. \end{aligned}$$

Performing an asymptotic expansion of the right-hand side for $1 \ll Y \lesssim s^{1/4}$, we find that

$$W(s, \psi) = \frac{(6\psi)^{4/3}}{4} \left(1 + 2Y^{-2} - \frac{12}{5}a_4bY^2 + O(s^{-2}Y^5 + s^{-13/8}Y^{\frac{5+a}{2}} + Y^{-3}) \right).$$

Since $Y \sim (6\psi)^{1/3}$, we obtain eventually, for $1 \ll \psi \lesssim s^{3/4}$,

$$W(s, \psi) = \frac{(6\psi)^{4/3}}{4} \left(1 + 2(6\psi)^{-2/3} - \frac{12}{5}a_4b(6\psi)^{2/3} + O(s^{-1}\psi^{1/3} + s^{-13/8}\psi^{\frac{7+a}{6}} + \psi^{-1}) \right). \quad (3.6)$$

We are now ready to construct a sub-solution for W beyond $cs^{1/4}$, for some constant $c > 0$ large but fixed, that will be determined later on. To that end, we introduce a *regularized*

modulation rate \tilde{b} , that has the same asymptotic behavior as b , but whose role is to remove some time oscillations. More precisely, define \tilde{b} by the ODE

$$\tilde{b}_s + \tilde{b}\tilde{b}' = 0, \quad \tilde{b}|_{s=s_0} = \frac{1}{s_0}. \quad (3.7)$$

We then have the following result (see Appendix B for a proof):

Lemma 3.5. *Assume that there exist constants $J > 0$, and $\epsilon > 0$ such that for all $s \in [s_0, s_1]$ such that for all $s \in [s_0, s_1]$*

$$\int_{s_0}^{s_1} |b_s + b^2|^2 s^{13/4} ds \leq J, \quad (3.8)$$

$$\frac{1 - \epsilon}{s} \leq b(s) \leq \frac{1 + \epsilon}{s}.$$

Then if s_0 is large enough (depending on K and ϵ), for all $s \geq s_0$,

$$\frac{1 - 2\epsilon}{s} \leq \tilde{b}(s) \leq \frac{1 + 2\epsilon}{s}.$$

We then define our subsolution and super-solutions in the following way:

Lemma 3.6. *Assume that:*

- There exist constants $J > 0$, $\eta \in (0, 1)$ and $\epsilon > 0$ such that (3.8) is satisfied;
- U satisfies the assumptions of Lemma 3.2;
- There exists a constant M_0 such that

$$U_{YY}(s_0) - 1 \geq -M_0 s_0^{-1} Y^2 \quad \forall Y \geq 0 \quad (3.9)$$

and such that $\lim_{Y \rightarrow \infty} U(s_0, Y) \leq M_0 s_0$;

- $E_1(s) \leq C_1 s^{-13/4}$ for all $s \in [s_0, s_1]$.

Then there exist a universal constant \bar{C} and a constant A_0 , depending only on M_0 , such that the following properties are satisfied:

- *Sub-solution:* For $A_- > 0$ define²

$$\underline{W}(s, \psi) := \frac{(6\psi)^{4/3}}{4} - A_- \psi^{7/3} \tilde{b}^{5/4} \quad \forall s \in [s_0, s_1], \quad \forall \psi \in \left[C_- \tilde{b}^{-3/4}, C_+ \tilde{b}^{-5/4} \right].$$

If $A_- \geq A_0$, $C_- \geq \bar{C}$ and if s_0 is large enough, then

$$\underline{W}(s, \psi) \leq W(s, \psi) \quad \forall s \in [s_0, s_1], \quad \psi \geq C_- \tilde{b}^{-3/4}.$$

- *Super-solution:* For $A_+ > 0$, define

$$\bar{W}(s, \psi) := \frac{(6\psi)^{4/3}}{4} + A_+ \psi^{10/3} \tilde{b}^2 \quad \forall s \in [s_0, s_1], \quad \forall \psi \geq C_- \tilde{b}^{-3/4}.$$

If $A_- \geq A_0$, $C_- \geq \bar{C}$ and if s_0 is large enough, then

$$W(s, \psi) \leq \bar{W}(s, \psi) \quad \forall s \in [s_0, s_1], \quad \psi \geq C_- \tilde{b}^{-3/4}.$$

The proof of Lemma 3.6 is postponed to the Appendix.

²The constant C_+ is such that $W(s, C_+ \tilde{b}^{-5/4}) = 0$. It can be determined explicitly, depending on A_- ; however its precise value is irrelevant.

3.3 Refined lower bound on $\partial_{YY}U$

Lemma 3.6 allowed us to extend the lower bound on W coming from the estimation of E_1 beyond $\psi \simeq s^{3/4}$. Thanks to this extension, we now construct a sub-solution for $U_{YY} - 1$ (or rather, for the function F introduced in Lemma 3.2). Eventually, the lower bound on $U_{YY} - 1$ will yield a finer lower bound on U .

Lemma 3.7. *Assume that the hypotheses of Lemma 3.6 are satisfied. Then, setting $M_2 := \max(M_0, \bar{M})$ for some universal constant \bar{M} , there exists a constant $c > 0$ such that*

$$U_{YY} - 1 \geq -M_2 b Y^2 \quad \forall Y \in [0, cs^{1/3}].$$

The proof is postponed to the Appendix.

Putting together the results of this section, we obtain Proposition 2.16.

4 Main tools for the energy estimates

This section is devoted to the derivation of several independent intermediate results which play an important role in the proof of energy estimates. We first prove the result on the decomposition of the diffusion term and on the remainder term, namely Lemma 2.6. We then turn to the commutator Lemma 2.8. We study the structure of the diffusion term (see Lemma 4.7). Eventually, we state some estimates allowing to perform a systematic treatment of some remainder terms.

For the sake of brevity, we adopt the following notation, which we will use extensively in the next two sections: for any $\alpha > 0$, and for quantities A and B that depend on s , we say that $A = O_\alpha(B)$ if there exists a constant C and a function $Q = Q(s, Y)$ with at most polynomial growth in s and Y , such that

$$\begin{aligned} |A(s, Y)| &\leq C|B(s, Y)| \quad \text{for } Y \leq s^\alpha, \\ |A(s, Y)| &\leq Q(s, Y) \quad \text{for } Y \geq s^\alpha. \end{aligned} \tag{4.1}$$

This notation will be useful because we work with weights of the form $w(s, Y) = Y^{-a}(1 + s^{-\beta}Y)^{-m}$, where m is an arbitrarily large integer. Therefore, the contribution of any function having at most polynomial growth in s and Y can be made as small as desired on the set $Y \geq s^\alpha$, in the following sense: if $\alpha > \beta$, for any integers $n, P \in \mathbb{N}$, if m is large enough (depending on α, β, n and P),

$$\int_{s^\alpha}^{\infty} (s^n + Y^n) w(s, Y) dY \leq s^{-P}.$$

In other words, when we estimate functions in $L^2(w)$, their behavior for $Y \gg s^\beta$ is unimportant, as long as these functions are polynomially bounded (with an explicit and computable bound).

4.1 Proof of Lemma 2.6

Since U^{app} is essentially a polynomial in b and Y (at least in the zone $Y \lesssim s^{2/7}$), the computation of the transport term $\partial_s U^{\text{app}} - bU^{\text{app}} + bY/2\partial_Y U^{\text{app}}$ is straightforward. Difficulties

stem from $L_U^{-1}(\partial_{YY}U^{\text{app}} - 1)$, which is also present in the diffusion term \mathcal{D} . Hence we start with a decomposition of the diffusion term

$$\mathcal{D} := L_U^{-1}(\partial_{YY}U - 1),$$

which will be useful in other occurrences. Writing $U = U^{\text{app}} + V$, we decompose \mathcal{D} into four parts:

- the biggest term, which we compute explicitly, and which is equal to $-b/2Y$. This term comes from U^{app} .
- a second order term \mathcal{L}_UV ;
- a first order term $\frac{b}{2}L_U^{-1}L_VY$;
- additional error terms coming from U^{app} , which we will treat as perturbations in all occurrences.

Our precise result concerning the diffusion term is the following:

Lemma 4.1. *We recall that $\mathcal{D} = L_U^{-1}(\partial_{YY}U - 1)$. Then*

$$\begin{aligned} \mathcal{D} &= -\frac{b}{2}Y + \mathcal{L}_UV + \frac{b}{2}L_U^{-1}L_VY \\ &\quad + L_U^{-1} \left(\left(\left(\frac{5}{4}a_7 - 90a_{10} \right) b^3Y^8 - 110a_{11}b^3Y^9 + 2a_{10}b^4Y^{11} + \frac{9}{4}a_{11}b^4Y^{12} \right) \chi \left(\frac{Y}{s^{2/7}} \right) \right) \\ &\quad + L_U^{-1} \left(P(s, Y)(1 - \bar{\chi}) \left(\frac{Y}{s^{2/7}} \right) \right) \\ &= -\frac{b}{2}Y + \mathcal{D}_{NL} + \tilde{\mathcal{D}} \end{aligned}$$

where $\chi, \bar{\chi} \in C_0^\infty(\mathbb{R})$ are cut-off functions such that $\chi, \bar{\chi} \equiv 1$ in a neighbourhood of zero, and P is a function that has at most polynomial growth in s and Y .

We have set

$$\begin{aligned} \mathcal{D}_{NL} &:= \mathcal{L}_UV + \frac{b}{2}L_U^{-1}L_VY, \\ \tilde{\mathcal{D}} &:= L_U^{-1} \left(\left(\left(\frac{5}{4}a_7 - 90a_{10} \right) b^3Y^8 - 110a_{11}b^3Y^9 + 2a_{10}b^4Y^{11} + \frac{9}{4}a_{11}b^4Y^{12} \right) \chi \left(\frac{Y}{s^{2/7}} \right) \right) \\ &\quad + L_U^{-1} \left(P(s, Y)(1 - \bar{\chi}) \left(\frac{Y}{s^{2/7}} \right) \right). \end{aligned}$$

Remark 4.2. *The decomposition of Lemma 4.1 will be used in two different occurrences:*

- First, we will use it to decompose the total diffusion term \mathcal{D} into a dissipation operator acting on the error term V , namely \mathcal{L}_UV , and remainder terms, namely $-\frac{b}{2}Y$, $\frac{b}{2}L_U^{-1}(L_VY)$ and $\tilde{\mathcal{D}}$. As we derive an equation on V , the diffusion term \mathcal{L}_UV will be kept in the left-hand side of the equation, while the remainder terms will be added to the terms stemming from U^{app} in the left-hand side.

- Additionally, \mathcal{D} will appear in the commutator of \mathcal{L}_U with $\partial_s + \frac{b}{2}Y\partial_Y$. We will then isolate the term in \mathcal{D} which bears the highest number of derivatives on V , namely $\mathcal{L}_U V$, which we will need to estimate separately in some instances.

Proof. Throughout the proof, since we are not interested in the specific definitions of the functions $P, \chi, \bar{\chi}$, we keep uniform notations for these three objects, even though they are used to group together different terms.

Recalling the definition of U^{app} (2.11), we have

$$\begin{aligned}
\mathcal{D} &= L_U^{-1}(\partial_{YY}U - 1) = L_U^{-1}(\partial_{YY}U^{\text{app}} - 1) + \mathcal{L}_U V \\
&= -L_U^{-1} \left((12a_4bY^2 + 42a_7b^2Y^5 + 90a_{10}b^3Y^8 + 110a_{11}b^3Y^9)\chi \left(\frac{Y}{s^{2/7}} \right) \right) \\
&\quad + L_U^{-1}(F''(\sqrt{bY}) - 1) \\
&\quad + L_U^{-1} \left((Y - a_4bY^4 - a_7b^2Y^7 - a_{10}b^3Y^{10} - a_{11}b^3Y^{11}) \frac{1}{s^{4/7}} \chi'' \left(\frac{Y}{s^{2/7}} \right) \right) \\
&\quad + 2L_U^{-1} \left((1 - 4a_4bY^3 - 7a_7b^2Y^6 - 10a_{10}b^3Y^9 - 11a_{11}b^3Y^{10}) \chi' \left(\frac{Y}{s^{2/7}} \right) \right) \\
&\quad + \mathcal{L}_U V.
\end{aligned}$$

We now examine each of the terms in the right-hand side separately.

- The term $F''(\sqrt{bY}) - 1$ and all the terms involving at least one derivative of χ are identically zero up to $Y \sim s^{2/7}$. Therefore they can all be written as $P(s, Y)(1 - \bar{\chi}) \left(\frac{Y}{s^{2/7}} \right)$.
- We therefore focus on

$$L_U^{-1} \left((12a_4bY^2 + 42a_7b^2Y^5 + 90a_{10}b^3Y^8 + 110a_{11}b^3Y^9)\chi \left(\frac{Y}{s^{2/7}} \right) \right),$$

and in particular on the value of this term on $Y \leq s^{2/7}$. Indeed, all values for $Y \geq s^{2/7}$ can be written as $L_U^{-1}(P(s, Y)(1 - \bar{\chi}) \left(\frac{Y}{s^{2/7}} \right))$.

We recall that $12a_4 = 1/4$, and that $42a_7 = a_4/2$. We compute

$$\begin{aligned}
L_U(Y) &= UY - U_Y \frac{Y^2}{2} \\
&= U^{\text{app}}Y - U_Y^{\text{app}} \frac{Y^2}{2} + L_V Y \\
&= \left(\frac{Y^2}{2} + a_4bY^5 + \frac{5}{2}a_7b^2Y^8 + 4a_{10}b^3Y^{11} + \frac{9}{2}a_{11}b^3Y^{12} \right) \chi \left(\frac{Y}{s^{2/7}} \right) \quad (4.2) \\
&\quad + L_V Y + P(s, Y)(1 - \bar{\chi}) \left(\frac{Y}{s^{2/7}} \right).
\end{aligned}$$

Multiplying (4.2) by $b/2$ and applying L_U^{-1} , we deduce that

$$\begin{aligned}
& L_U^{-1} \left((12a_4bY^2 + 42a_7b^2Y^5 + 90a_{10}b^3Y^8 + 110a_{11}b^3Y^9) \chi \left(\frac{Y}{s^{2/7}} \right) \right) \\
&= \frac{b}{2}Y - \frac{b}{2}L_U^{-1}L_VY \\
&+ L_U^{-1} \left(\left((90a_{10} - \frac{5}{4}a_7)b^3Y^8 + 110a_{11}b^3Y^9 - 2a_{10}b^4Y^{11} - \frac{9}{4}a_{11}b^4Y^{12} \right) \chi \left(\frac{Y}{s^{2/7}} \right) \right) \\
&+ L_U^{-1} \left(P(s, Y)(1 - \bar{\chi}) \left(\frac{Y}{s^{2/7}} \right) \right).
\end{aligned}$$

Gathering all the terms, we obtain the decomposition announced in the Lemma. □

Corollary 4.3. *Assume that there exist constants M, c such that*

$$\begin{aligned}
|\partial_{YY}U - 1| &\leq MbY^2 \quad \forall Y \in [0, cs^{1/3}], \quad \forall s \in [s_0, s_1], \\
\frac{1}{2s} &\leq b \leq \frac{2}{s}, \\
|U_{YY}| &\leq M \quad \forall Y > 0, \quad \forall s \in [s_0, s_1].
\end{aligned}$$

Then

$$\mathcal{D} = O_{1/3}(bY), \quad \partial_Y \mathcal{D} = \partial_Y \mathcal{L}_U V + O_{2/7}(b).$$

Proof. Note first that under these assumptions,

$$\begin{aligned}
Y + \frac{Y^2}{2} - \frac{M}{12}bY^4 &\leq U(s, Y) \leq Y + \frac{Y^2}{2} + \frac{M}{12}bY^4, \\
1 + Y - \frac{M}{4}bY^3 &\leq U_Y(s, Y) \leq 1 + Y + \frac{M}{4}bY^3,
\end{aligned} \quad \forall Y \in [0, cs^{1/3}].$$

The estimate on \mathcal{D} follows simply from writing

$$\mathcal{D} = U_Y \int_0^Y \frac{\partial_{YY}U - 1}{U^2} + \frac{\partial_{YY}U - 1}{U}$$

and using the bounds on $\partial_{YY}U, U_Y$ and U . As for the second one, notice that

$$\partial_Y \mathcal{D} = \partial_Y \mathcal{L}_U V - \frac{b}{2} + \partial_Y L_U^{-1} Z,$$

where

$$\begin{aligned}
Z &= \frac{b}{2}(YV - \frac{Y^2}{2}V_Y) \\
&+ \left(\left(\frac{5}{4}a_7 - 90a_{10} \right) b^3Y^8 - 110a_{11}b^3Y^9 + 2a_{10}b^4Y^{11} + \frac{9}{4}a_{11}b^4Y^{12} \right) \chi \left(\frac{Y}{s^{2/7}} \right) \\
&+ P(s, Y)(1 - \bar{\chi}) \left(\frac{Y}{s^{2/7}} \right)
\end{aligned}$$

where P has at most polynomial growth in s and Y . The estimate then follows from the bounds

$$V = O_{2/7}(bY^4), \quad V_Y = O_{2/7}(bY^3), \quad V_{YY} = O_{2/7}(bY^2)$$

and from the formula giving $\partial_Y L_U^{-1}$ in Lemma A.1. □

We deduce that V is a solution of equation (2.16), i.e.

$$\partial_s V - bV + \frac{b}{2}Y \partial_Y V - \mathcal{L}_U V = \mathcal{R},$$

with a remainder

$$\begin{aligned} \mathcal{R} &:= - \left(\partial_s U^{\text{app}} - bU^{\text{app}} + \frac{b}{2}Y \partial_Y U^{\text{app}} \right) - \frac{b}{2}Y \\ &+ L_U^{-1} \left(\left(\left(\frac{5}{4}a_7 - 90a_{10} \right) b^3 Y^8 - 110a_{11} b^3 Y^9 + 2a_{10} b^4 Y^{11} + \frac{9}{4}a_{11} b^4 Y^{12} \right) \chi \left(\frac{Y}{s^{2/7}} \right) \right) \\ &+ \frac{b}{2} L_U^{-1} (L_V Y) \\ &+ L_U^{-1} \left(P(s, Y)(1 - \bar{\chi}) \left(\frac{Y}{s^{2/7}} \right) \right), \end{aligned}$$

which we now compute.

Lemma 4.4 (Computation of \mathcal{R}). *The remainder term \mathcal{R} can be written as*

$$\begin{aligned} \mathcal{R} &= (b_s + b^2)(a_4 Y^4 + 2a_7 b Y^7 + 3a_{10} b^2 Y^{10} + 3a_{11} b^2 Y^{11}) \chi \left(\frac{Y}{s^{2/7}} \right) \\ &+ (a_{10} b^4 Y^{10} + 3a_{11} b^4 / 2 Y^{11}) \chi \left(\frac{Y}{s^{2/7}} \right) \\ &+ \frac{b}{2} L_U^{-1} (L_V Y) + \frac{1}{2} b^3 a_7 L_U^{-1} \left(\left(Y^7 V - V_Y \frac{Y^8}{8} \right) \chi \left(\frac{Y}{s^{2/7}} \right) \right) \\ &+ L_U^{-1} \left[(d_{11} b^4 Y^{11} + d_{12} b^5 Y^{12} + d_{14} b^5 Y^{14} + d_{17} b^6 Y^{17} + d_{18} b^6 Y^{18}) \chi \left(\frac{Y}{s^{2/7}} \right) \right] \\ &+ L_U^{-1} \left(P(s, Y)(1 - \bar{\chi}) \left(\frac{Y}{s^{2/7}} \right) \right), \end{aligned}$$

for some explicit constants $d_{11}, d_{12}, d_{14}, d_{17}, d_{18} \in \mathbb{R}$.

Remark 4.5. *Notice that this remainder term is essentially (up to a small error depending on V)*

$$L_U^{-1} \left(U^{\text{app}} \partial_s U^{\text{app}} - \partial_Y U^{\text{app}} \int_0^Y U^{\text{app}} - 2b(U^{\text{app}})^2 + \frac{3b}{2} \partial_Y U^{\text{app}} \int_0^Y U^{\text{app}} - \partial_{YY} U^{\text{app}} + 1 \right),$$

and therefore has been computed (up to the application of the operator L_U^{-1}) when the approximate solution U^{app} was defined. However, it is actually easier to do over the computations rather than to apply L_U^{-1} to the remainder that has already been computed.

Proof. We start with

$$\begin{aligned}
& \partial_s U^{\text{app}} - bU^{\text{app}} + \frac{b}{2}Y\partial_Y U^{\text{app}} + \frac{b}{2}Y \\
&= -a_4(b_s + b^2)Y^4\chi\left(\frac{Y}{s^{2/7}}\right) \\
&+ \chi\left(\frac{Y}{s^{2/7}}\right)\left[-a_7b\left(2b_s + \frac{5b^2}{2}\right)Y^7 - a_{10}Y^{10}(3b_sb^2 + 4b^4) - a_{11}Y^{11}\left(3b_sb^2 + \frac{9}{2}b^4\right)\right] \\
&+ \chi'\left(\frac{Y}{s^{2/7}}\right)\frac{Y}{s^{2/7}}\left(\frac{b}{2} - \frac{2/7}{s}\right)\left[Y - a_4bY^4 - a_7b^2Y^7 - a_{10}b^3Y^{10} - a_{11}b^3Y^{11}\right] \\
&+ b^{-2}(b_s + b^2)\left(\frac{1}{2}Z\Theta'(Z) - \Theta(Z)\right)\Big|_{Z=\sqrt{b}Y} \\
&+ \frac{b}{2}Y(1 - \chi)\left(\frac{Y}{s^{2/7}}\right).
\end{aligned}$$

The last three terms in the right-hand side are supported in $Y \geq c_1s^{2/7}$. They can be written as a linear combination of terms of the type $P(s, Y)(1 - \bar{\chi})\left(\frac{Y}{s^{2/7}}\right)$.

We thus focus on the second term, which we group with the other terms in the definition of \mathcal{R} . We first isolate the factor $(b_s + b^2)$ in Y^7, Y^{10}, Y^{11} , which we group with the first term. There remains to study

$$\begin{aligned}
& \frac{1}{2}a_7b^3Y^7\chi\left(\frac{Y}{s^{2/7}}\right) \\
&+ b^3L_U^{-1}\left(\left(\left(\frac{5}{4}a_7 - 90a_{10}\right)Y^8 - 110a_{11}Y^9 + 2a_{10}bY^{11} + \frac{9}{4}a_{11}bY^{12}\right)\chi\left(\frac{Y}{s^{2/7}}\right)\right).
\end{aligned}$$

We use the same trick as in Lemma 4.1 and we write

$$Y^7\chi\left(\frac{Y}{s^{2/7}}\right) = L_U^{-1}\left((L_{U^{\text{app}}} + L_V)Y^7\chi\left(\frac{Y}{s^{2/7}}\right)\right)$$

Notice that $L_V Y^7 = VY^7 - V_Y Y^8/8$. On the other hand, a lengthy but straightforward computation yields

$$L_{U^{\text{app}}}Y^7 = \frac{7}{8}Y^8 + \frac{3}{8}Y^9 - \frac{a_4}{2}bY^{11} - \frac{a_7}{8}b^2Y^{14} + \frac{a_{10}}{4}b^3Y^{17} + \frac{3a_{11}}{8}b^3Y^{18}.$$

Gathering all the terms and recalling the values of a_{10}, a_{11} (2.10), we obtain the decomposition announced in the Lemma. □

4.2 Proof of the commutator result (Lemma 2.8)

We compute separately $[L_U^{-1}, \partial_s]$ and $[L_U^{-1}, Y\partial_Y]$, and then check that cancellations occur between the two commutators.

Using the formulas in Lemma A.1 in the Appendix, we have

$$\begin{aligned}
[L_U^{-1}, Y\partial_Y]W &= \left(U \int_0^Y \frac{Y\partial_Y W}{U^2} \right)_Y - Y\partial_Y L_U^{-1}W \\
&= U_Y \int_0^Y \frac{Y\partial_Y W}{U^2} + \frac{Y\partial_Y W}{U} - Y \left(\partial_Y^2 U \int_0^Y \frac{W}{U^2} + \frac{\partial_Y W}{U} \right) \\
&= U_Y \int_0^Y \frac{Y\partial_Y W}{U^2} - Y\partial_Y^2 U \int_0^Y \frac{W}{U^2}.
\end{aligned}$$

We introduce the quantity

$$\Gamma := YU_Y - 2U,$$

so that $\Gamma_Y = Y\partial_Y^2 U - U_Y$. Then

$$\begin{aligned}
[L_U^{-1}, Y\partial_Y]W &= -\Gamma_Y \int_0^Y \frac{W}{U^2} + U_Y \int_0^Y \frac{\partial}{\partial Y} \left(\frac{W}{Y} \right) \frac{Y^2}{U^2} \\
&= -\Gamma_Y \int_0^Y \frac{W}{U^2} + \frac{YU_Y}{U^2} W - 2U_Y \int_0^Y W \frac{U - YU_Y}{U^3} \\
&= 2L_U^{-1}W - \Gamma_Y \int_0^Y \frac{W}{U^2} + \frac{\Gamma}{U^2} W + 2U_Y \int_0^Y W \frac{\Gamma}{U^3}. \tag{4.3}
\end{aligned}$$

We now address the commutator with ∂_s . To that end, we recall that U satisfies (2.15), so that, with the previous definitions of f and \mathcal{D} ,

$$U_s = -\frac{b}{2}\Gamma + \mathcal{D}. \tag{4.4}$$

It follows that

$$\begin{aligned}
[L_U^{-1}, \partial_s]W &= -\left(U_s \int_0^Y \frac{W}{U^2} \right)_Y + 2\left(U \int_0^Y \frac{W}{U^3} U_s \right)_Y \\
&= \frac{b}{2} \left(\Gamma \int_0^Y \frac{W}{U^2} \right)_Y - b \left(U \int_0^Y \frac{W}{U^3} \Gamma \right)_Y - \left(\mathcal{D} \int_0^Y \frac{W}{U^2} \right)_Y + 2 \left(U \int_0^Y \frac{W}{U^3} \mathcal{D} \right)_Y.
\end{aligned}$$

Using (4.3), we infer that

$$\begin{aligned}
&\left[L_U^{-1}, \partial_s + \frac{b}{2}Y\partial_Y \right] W \\
&= bL_U^{-1}W - \frac{b}{2}\Gamma_Y \int_0^Y \frac{W}{U^2} + \frac{b}{2}\frac{\Gamma}{U^2}W + bU_Y \int_0^Y W \frac{\Gamma}{U^3} \\
&+ \frac{b}{2} \left(\Gamma \int_0^Y \frac{W}{U^2} \right)_Y - b \left(U \int_0^Y \frac{W}{U^3} \Gamma \right)_Y - \left(\mathcal{D} \int_0^Y \frac{W}{U^2} \right)_Y + 2 \left(U \int_0^Y \frac{W}{U^3} \mathcal{D} \right)_Y \\
&= bL_U^{-1}W - \left(\mathcal{D} \int_0^Y \frac{W}{U^2} \right)_Y + 2 \left(U \int_0^Y \frac{W}{U^3} \mathcal{D} \right)_Y.
\end{aligned}$$

This completes the proof of the commutator Lemma.

4.3 Structure of the diffusion term

We will use in several instances the following weighted Hardy inequality (see [23]):

Lemma 4.6. *Let p_1, p_2 be measurable functions such that $p_1, p_2 > 0$ almost everywhere. Let $0 < R \leq \infty$ and let*

$$C_H := 4 \sup_{0 < r < R} \left(\int_r^R p_1 \right) \left(\int_0^r \frac{1}{p_2} \right).$$

Assume that $C_H < +\infty$. Then for any function $f \in H_{loc}^1(\mathbb{R}_+)$ such that $f(0) = 0$, there holds

$$\int_0^R f^2 p_1 \leq C_H \int_0^R (\partial_Y f)^2 p_2.$$

In this paragraph, we state and prove the coercivity inequality that will be used to control (E_1, D_1) and (E_2, D_2) , up to small remainder terms.

Lemma 4.7. *Let $s_0 < s_1$, and let $\delta > 0$ be arbitrary. Assume that U is increasing in Y for all $s \geq s_0$, with $U|_{Y=0} = 0$, $\partial_Y U|_{Y=0} = 1$, and that there exists constants M_2, c such that*

$$\begin{aligned} 1 - M_2 b Y^2 &\leq \partial_{YY} U \leq 1 \quad \forall Y \in [0, cs^{1/3}], \quad \forall s \in [s_0, s_1], \\ -M_2 &\leq \partial_{YY} U \leq 1 \quad \forall Y \geq cs^{1/3} \quad \forall s \in [s_0, s_1]. \end{aligned} \quad (4.5)$$

Assume furthermore that

$$\frac{1}{2s} \leq b \leq \frac{2}{s} \quad \forall s \in [s_0, s_1].$$

For $a > 0, \beta \in]1/4, 2/7[, m \in \mathbb{N}$, define the weight $w(s, Y) := Y^{-a}(1 + s^{-\beta}Y)^{-m}$.

There exist universal constants $\bar{a}, \bar{c} > 0$, independent of δ, s, β, m , such that if s_0 is large enough (depending on δ, β, a), then for all $f \in W^{1,\infty}(\mathbb{R}_+)$ such that $\partial_Y f = O(Y)$ for $Y \ll 1$, for all $0 < a \leq \bar{a}$, for any cut-off function $\chi \in \mathcal{C}_0^\infty(\mathbb{R}_+)$ such that $\chi \equiv 1$ in $[0, 1]$,

$$\begin{aligned} - \int_0^\infty (\partial_{YY} L_U^{-1} f) f w &\geq \bar{c} \left(\int_0^\infty \frac{(\partial_Y f)^2}{U} w + \int_0^\infty \frac{f^2}{U^2} w \right) - \delta b \int_0^\infty f^2 w \\ &\quad - C \int_{cs^{1/3}}^\infty U \left(\int_0^Y \left(1 - \chi \left(\frac{Y}{s^{1/4}} \right) \right) \frac{f}{U^2} \right)^2 w. \end{aligned}$$

Remark 4.8. *Note that the estimate we prove here is not as strong as one would like to have. Indeed, we only control f and $\partial_Y f$, and not $L_U^{-1} f$. However, another way of writing the inequality, setting $h = L_U^{-1} f$, is*

$$- \int_0^\infty \partial_{YY} h L_U h w \geq \bar{c} \int_0^\infty \frac{(\partial_Y(L_U h))^2}{U} w + \int_0^\infty \frac{(L_U h)^2}{U^2} w + \text{remainder terms.}$$

But notice that if $h = U_Y$, then $L_U h = 0$ while $h \neq 0$, $\partial_Y h \neq 0$. Hence it is hopeless to control h and $\partial_Y h$ by the left-hand side of the above inequality without any further assumption that would discard the case $h = U_Y$.

Remark 4.9. *The last term in the right-hand side of the inequality, namely*

$$C \int_{cs^{1/3}}^{\infty} U \left(\int_0^Y \left(1 - \chi \left(\frac{Y}{s^{1/4}} \right) \right) \frac{f}{U^2} \right)^2 w$$

will be handled in Lemma 4.15. Heuristically, since it is supported in a zone where w is strongly decaying, it can be made “as small as desired”, i.e. $O(s^{-P})$ for any $P > 0$, provided m is chosen sufficiently large (depending on P and β).

Remark 4.10. *Notice that under the assumptions of the Lemma, there exists a constant C such that for $1 \leq Y \leq cs^{1/3}$,*

$$\frac{C^{-1}}{Y^2} \leq \frac{1}{U} \leq \frac{C}{Y^2}, \quad \frac{C^{-2}}{Y^4} \leq \frac{1}{U^2} \leq \frac{C^2}{Y^4}.$$

Hence, for $1 \leq Y \leq cs^{1/3}$, the two integrals in the diffusion term

$$\int_0^{\infty} \frac{(\partial_Y f)^2}{U} w + \int_0^{\infty} \frac{f^2}{U^2} w$$

have the same scaling.

However, if $Y \leq 1$, then

$$\frac{C^{-1}}{Y} \leq \frac{1}{U} \leq \frac{C}{Y}, \quad \frac{C^{-2}}{Y^2} \leq \frac{1}{U^2} \leq \frac{C^2}{Y^2}.$$

Hence for $Y \leq 1$, the control given by the first integral is stronger. Indeed, the Hardy inequality implies that

$$\int_0^1 \frac{(\partial_Y f)^2}{Y^{1+a}} \geq C \int_0^1 \frac{f^2}{Y^{3+a}}.$$

Therefore

$$\int_0^{cs^{1/3}} \left(\frac{(\partial_Y f)^2}{U} + \frac{f^2}{U^2} \right) w \geq C \left(\int_1^{cs^{1/3}} \frac{f^2}{Y^4} w + \int_0^1 \frac{f^2}{Y^{3+a}} \right).$$

We will often use this control close to zero when we deal with some non-local terms.

Proof of Lemma 4.7. Starting with a simple integration by parts,

$$- \int_0^{\infty} (\partial_{YY} L_U^{-1} f) f w = \int_0^{\infty} (\partial_Y L_U^{-1} f) (\partial_Y f w + f \partial_Y w).$$

Using the formula in lemma A.1 and integrating by parts once again, we obtain

$$\begin{aligned} - \int_0^{\infty} (\partial_{YY} L_U^{-1} f) f w &= \int_0^{\infty} \frac{(\partial_Y f)^2}{U} w - \int_0^{\infty} \frac{f^2}{U^2} w \\ &\quad + \int_0^{\infty} (U_{YY} - 1) \left(\int_0^Y \frac{f}{U^2} \right) (f \partial_Y w + \partial_Y f w) \\ &\quad + \int_0^{\infty} \frac{1}{U} \partial_Y f f \partial_Y w. \end{aligned}$$

The first two terms are the main order terms. We now prove the coercivity thanks to a weighted Hardy inequality for which we compute the constant explicitly. Using the assumptions on U , we have

$$U \geq Y + \frac{Y^2}{2} - \frac{M_2}{12} b Y^4 \quad \text{for } Y \in [0, cs^{1/3}].$$

Therefore, since U is increasing,

$$U(s, Y) \geq \frac{K^2}{4} s^{1/2} \quad \forall Y \geq K s^{1/4}$$

provided s_0 is sufficiently large. It follows that if K is chosen large enough (depending on δ),

$$\int_{Ks^{1/4}}^{\infty} \frac{f^2}{U^2} w \leq \frac{16}{K^4} s^{-1} \int_{Ks^{1/4}}^{\infty} f^2 w \leq \delta b \int_0^{\infty} f^2 w.$$

On the set $[0, Ks^{1/4}]$, we use a weighted Hardy inequality (see Lemma 4.6), namely

$$\int_0^{Ks^{1/4}} \frac{1}{U^2} f^2 w \leq C_a \int_0^{Ks^{1/4}} \frac{(\partial_Y f)^2}{U} w$$

where the constant C_a satisfies

$$C_a \leq 4 \sup_{0 < r < Ks^{1/4}} \left(\int_r^{Ks^{1/4}} \frac{w}{U^2} \right) \left(\int_0^r \frac{U}{w} \right).$$

On the set $[0, Ks^{1/4}]$, we have

$$\begin{aligned} \mu Y + \frac{Y^2}{2} \leq U \leq Y + \frac{Y^2}{2} \quad \text{with } \mu = 1 - \frac{M_2}{24} K^3 s^{-1/4}, \\ \text{and } Y^{-a}(1 - \delta) \leq w \leq Y^{-a} \end{aligned}$$

for any $\delta > 0$ provided s_0 is sufficiently large. Therefore

$$C_a \leq \frac{1}{1 - \delta} \bar{C}_{a, \mu},$$

where

$$\bar{C}_{a, \mu} := 4 \sup_{r > 0} \left(\int_r^{\infty} \frac{Y^{-a}}{\left(\mu Y + \frac{Y^2}{2} \right)^2} dY \right) \left(\int_0^r Y^a \left(Y + \frac{Y^2}{2} \right) dY \right).$$

We then have the following Lemma, which is proved in the Appendix:

Lemma 4.11. *There exists universal constants $\bar{a} > 0, \mu_0 \in (0, 1)$ such that for all $a \in (0, \bar{a})$, for all $\mu \in (\mu_0, 1)$,*

$$C_{a, \mu} \leq \frac{9}{10}.$$

Therefore, for any $a \in (0, \bar{a})$, provided δ is small enough (say $\delta < 1/50$) and s_0 is sufficiently large,

$$C_a \leq 1 - \frac{1}{20}$$

and thus

$$\int_0^\infty \frac{(\partial_Y f)^2}{U} w - \int_0^\infty \frac{f^2}{U^2} w \geq \frac{1}{40} \left(\int_0^\infty \frac{(\partial_Y f)^2}{U} w + \int_0^\infty \frac{f^2}{U^2} w \right) - \delta b \int_0^\infty f^2 w. \quad (4.6)$$

There remains to estimate the two lower order terms, namely

$$\int_0^\infty \frac{1}{U} \partial_Y f f \partial_Y w \quad \text{and} \quad \int_0^\infty (U_{YY} - 1) \left(\int_0^Y \frac{f}{U^2} \right) (f \partial_Y w + \partial_Y f w).$$

For the first lower order term, we distinguish once again between the zones $Y \leq Ks^{1/4}$ and $Y \geq Ks^{1/4}$. Using the Cauchy-Schwartz inequality, we have

$$\left| \int_0^\infty \frac{1}{U} \partial_Y f f \partial_Y w \right| \leq \left(\int_0^\infty \frac{(\partial_Y f)^2}{U} w \right)^{1/2} \left(\int_0^\infty \frac{f^2 (\partial_Y w)^2}{U w} \right)^{1/2}.$$

For $Y \leq Ks^{1/4}$, we have, for s_0 large enough (depending on a, m, β, K)

$$|\partial_Y w| \leq 2aY^{-1}w.$$

Using a Hardy inequality, we have

$$\int_0^{Ks^{1/4}} f^2 \frac{1}{Y^2 U} w \leq C_H \int_0^{Ks^{1/4}} \frac{(\partial_Y f)^2}{U} w \quad (4.7)$$

where the constant C_H is defined by

$$C_H = 4 \sup_{0 < r < Ks^{1/4}} \left(\int_r^{Ks^{1/4}} \frac{1}{Y^2 U} w \right) \left(\int_0^r U \frac{1}{w} \right).$$

As above, we have, provided s_0 is sufficiently large (depending on m, β, K)

$$C_H \leq 16 \sup_{r > 0} \left(\int_r^\infty \frac{1}{Y^{2+a}(Y+Y^2)} dY \right) \left(\int_0^r Y^a (Y+Y^2) dY \right).$$

Studying separately the cases $r < 1$ and $r > 1$, it can be easily proved that C_H is bounded uniformly in a and s , so that there exists a universal constant \bar{C} such that for s large enough

$$\int_0^{Ks^{1/4}} \frac{f^2 (\partial_Y w)^2}{U w} \leq \bar{C} a \int_0^{Ks^{1/4}} \frac{(\partial_Y f)^2}{U} w.$$

This term is absorbed in the main order diffusion term for a small enough. Now, for $Y \geq Ks^{1/4}$, we have

$$|\partial_Y w| \leq (a+m)Y^{-1}w \leq (a+m)K^{-1}s^{-1/4}w,$$

and $U(s, Y) \geq \frac{K^2}{4}s^{1/2}$. As a consequence,

$$\int_{Ks^{1/4}}^\infty \frac{f^2 (\partial_Y w)^2}{U w} \leq 4 \frac{(a+m)^2}{K^4} s^{-1} \int_0^\infty f^2 w \leq \delta b \int_0^\infty f^2 w$$

for K large enough (depending on m, δ). We infer that

$$\left| \int_0^\infty \frac{1}{U} \partial_Y f f \partial_Y w \right| \leq \bar{C} a^{1/2} \int_0^\infty \frac{(\partial_Y f)^2}{U} w + \delta b \int_0^\infty f^2 w. \quad (4.8)$$

We now address the second lower order term. We focus on the term involving $\partial_Y f w$, since the one with $f \partial_Y w$ can be treated with similar ideas combined with the same estimates as above. Using assumption (4.5) on $\partial_Y U$, we have

$$\begin{aligned} & \left| \int_0^{cs^{1/3}} (U_{YY} - 1) \left(\int_0^Y \frac{f}{U^2} \right) \partial_Y f w \right| \\ & \leq M_2 b \left(\int_0^{cs^{1/3}} \frac{(\partial_Y f)^2}{U} w \right)^{1/2} \left(\int_0^{cs^{1/3}} Y^4 \left(\int_0^Y \frac{f}{U^2} \right)^2 U w \right)^{1/2}. \end{aligned}$$

We separate the last integral in the right-hand side into three zones: $Y \leq 1$, $1 \leq Y \leq Ks^{1/4}$ for some large constant K , and $Y \geq Ks^{1/4}$. Notice that if $Y \leq Ks^{1/4}$, then

$$w(s, Y) \geq (1 + Ks_0^{1/4-\beta})^{-m} Y^{-a} \geq \frac{1}{2} Y^{-a}$$

for s_0 large enough (depending on K, m and β). For $Y \leq 1$, we have, using (4.7)

$$\left| \int_0^Y \frac{f}{U^2} \right| \leq \left(\int_0^Y \frac{f^2}{Y^{2+a}U} \right)^{1/2} \left(\int_0^Y \frac{Y^{2+a}}{U^3} \right) \leq \frac{\bar{C}}{a^{1/2}} \left(\int_0^1 \frac{(\partial_Y f)^2}{U} w \right)^{1/2}.$$

And if $1 \leq Y \leq Ks^{1/4}$, using a simple Cauchy-Schwartz inequality,

$$\int_0^Y \frac{|f|}{U^2} \leq \frac{\bar{C}}{a^{1/2}} \left(\int_0^1 \frac{(\partial_Y f)^2}{U} w \right)^{1/2} + \bar{C} \left(\int_1^Y \frac{f^2}{U^2} w \right)^{1/2}.$$

Therefore, we have

$$\begin{aligned} \int_0^{Ks^{1/4}} Y^4 \left(\int_0^Y \frac{f}{U^2} \right)^2 U w & \leq \frac{\bar{C}}{a} \left[\int_0^\infty \frac{(\partial_Y f)^2}{U} w + \int_0^\infty \frac{f^2}{U^2} w \right] \int_0^{Ks^{1/4}} Y^{4-a} U \\ & \leq \frac{\bar{C} K^7}{a} \left[\int_0^\infty \frac{(\partial_Y f)^2}{U} w + \int_0^\infty \frac{f^2}{U^2} w \right] s^{(7-a)/4}. \end{aligned}$$

It follows that

$$\begin{aligned} M_2 b \left(\int_0^{cs^{1/3}} \frac{(\partial_Y f)^2}{U} w \right)^{1/2} \left(\int_0^{Ks^{1/4}} Y^4 \left(\int_0^Y \frac{f}{U^2} \right)^2 U w \right)^{1/2} \\ \leq \bar{C} M_2 \frac{K^{7/2}}{a^{1/2}} s^{-\frac{1+a}{8}} \left(\int_0^\infty \frac{(\partial_Y f)^2}{U} w + \int_0^\infty \frac{f^2}{U^2} w \right). \end{aligned}$$

Therefore, for s_0 sufficiently large (depending on K and a), this term can be absorbed in the main order diffusion term. On the other hand, using the same estimates as above,

$$\begin{aligned}
& \int_{Ks^{1/4}}^{cs^{1/3}} Y^4 \left(\int_0^Y \frac{f}{U^2} \right)^2 Uw \\
& \leq \frac{C}{a} \left[\int_0^\infty \frac{(\partial_Y f)^2}{U} w + \int_0^\infty \frac{f^2}{U^2} w \right] \int_{Ks^{1/4}}^\infty Y^4 U w + \int_{Ks^{1/4}}^{cs^{1/3}} Y^4 U \left(\int_{Ks^{1/4}}^Y \frac{|f|}{U^2} \right)^2 w \\
& \leq C_{a,m} s^{(7-a)\beta} \left[\int_0^\infty \frac{(\partial_Y f)^2}{U} w + \int_0^\infty \frac{f^2}{U^2} w \right] + C_{m,\beta}(s) \int_{Ks^{1/4}}^{cs^{1/3}} \frac{f^2}{U^4} w
\end{aligned}$$

where

$$\begin{aligned}
C_{m,\beta}(s) &= 4 \sup_{r \in (Ks^{1/4}, cs^{1/3})} \left(\int_r^{cs^{1/3}} Y^4 U w \right) \left(\int_{Ks^{1/4}}^r \frac{1}{w} \right) \\
&\leq 8s^{8\beta} \sup_{r \in (Ks^{\frac{1}{4}-\beta}, cs^{\frac{1}{3}-\beta})} \left(\int_r^{cs^{\frac{1}{3}-\beta}} Z^{6-a} (1+Z)^{-m} dZ \right) \left(\int_{Ks^{\frac{1}{4}-\beta}}^r Z^a (1+Z)^m dZ \right) \\
&\leq C_m s^{8\beta} s^{8(\frac{1}{3}-\beta)} \leq C_m s^{8/3}.
\end{aligned}$$

It follows that, if $\beta < 2/7$, for s_0 sufficiently large,

$$\begin{aligned}
& M_2 b \left(\int_0^{cs^{1/3}} \frac{(\partial_Y f)^2}{U} w \right)^{1/2} \left(\int_{Ks^{1/4}}^{cs^{1/3}} Y^4 \left(\int_0^Y \frac{f}{U^2} \right)^2 Uw \right)^{1/2} \\
& \leq C_{a,m} M_2 s^{(7-a)\frac{\beta}{2}-1} \left[\int_0^\infty \frac{(\partial_Y f)^2}{U} w + \int_0^\infty \frac{f^2}{U^2} w \right] \\
& \quad + \delta \left[\int_0^\infty \frac{(\partial_Y f)^2}{U} w + \int_0^\infty \frac{f^2}{U^2} w \right] \\
& \quad + C_{\delta,m} b^2 s^{8/3} K^{-8} s^{-2} \int_0^\infty f^2 w \\
& \leq \delta \left[\int_0^\infty \frac{(\partial_Y f)^2}{U} w + \int_0^\infty \frac{f^2}{U^2} w + b \int_0^\infty f^2 w \right].
\end{aligned}$$

There remains to consider the part of the second lower order term for $Y \geq cs^{1/3}$. Using the cut-off function χ , we have

$$\begin{aligned}
& \left| \int_{cs^{1/3}}^\infty (U_{YY} - 1) \left(\int_0^Y \frac{f}{U^2} \right) \partial_Y f w \right| \\
& \leq \delta \int_0^\infty \frac{(\partial_Y f)^2}{U} w + \frac{(M_2 + 1)^2}{4\delta} \int_{cs^{1/3}}^\infty U \left(\int_0^Y \left(1 - \chi \left(\frac{Y}{s^{1/4}} \right) + \chi \left(\frac{Y}{s^{1/4}} \right) \right) \frac{f}{U^2} \right)^2 w \\
& \leq \delta \int_0^\infty \frac{(\partial_Y f)^2}{U} w + \frac{C}{\delta} \left(\int_0^{Ks^{1/4}} \frac{|f|}{U^2} \right)^2 \int_{cs^{1/3}}^\infty U w \\
& \quad + \frac{C}{\delta} \int_{cs^{1/3}}^\infty U \left(\int_0^Y \left(1 - \chi \left(\frac{Y}{s^{1/4}} \right) \right) \frac{f}{U^2} \right)^2 w.
\end{aligned}$$

The second term in the right-hand side is estimated as above. Notice that for any $P > 0$,

$$\int_{cs^{1/3}}^{\infty} U w \leq \int_{cs^{1/3}}^{\infty} \left(Y + \frac{Y^2}{2} \right) w \leq s^{-P}$$

provided m is sufficiently large (depending on β and P). Therefore we obtain, for any $\delta > 0$, provided $s \geq s_0$ with s_0 sufficiently large,

$$\begin{aligned} & \left| \int_0^{\infty} (U_{YY} - 1) \left(\int_0^Y \frac{f}{U^2} \right) (f \partial_Y w + \partial_Y f w) \right| \\ & \leq \delta \left(\int_0^{\infty} \frac{(\partial_Y f)^2}{U} w + \int_0^{\infty} \frac{f^2}{U^2} w + b \int_0^{\infty} f^2 w \right) + \frac{C}{\delta} \int_{cs^{1/3}}^{\infty} U \left(\int_0^Y \left(1 - \chi \left(\frac{Y}{s^{1/4}} \right) \right) \frac{f}{U^2} \right)^2 w. \end{aligned} \quad (4.9)$$

Gathering (4.6), (4.8) and (4.9), we conclude that there exists a positive universal constant \bar{c} (say for instance $\bar{c} = 1/50$), and $\bar{a} > 0$, $\bar{\delta} > 0$, such that for all $0 < a < \bar{a}$ and for all $\beta \in (1/4, 2/7)$, $m \geq 1$, $\delta \in (0, \bar{\delta})$, there exists $s_0 > 0$ such that if $s \geq s_0$, then

$$\begin{aligned} \int_0^{\infty} (\partial_Y^2 L_U^{-1} f) f w & \geq \bar{c} \left(\int_0^{\infty} \frac{(\partial_Y f)^2}{U} w + \int_0^{\infty} \frac{f^2}{U^2} w \right) - \delta b \int_0^{\infty} f^2 w \\ & \quad - C \int_{cs^{1/3}}^{\infty} U \left(\int_0^Y \left(1 - \chi \left(\frac{Y}{s^{1/4}} \right) \right) \frac{f}{U^2} \right)^2 w. \end{aligned} \quad (4.10)$$

This completes the proof of Lemma 4.7. \square

4.4 Structure of the commutator

We record here some formulas and a few estimates that will be useful in the estimation of commutator terms. We recall that $\mathcal{D} = L_U^{-1}(\partial_{YY}U - 1) = \mathcal{L}_U V + L_U^{-1}(\partial_{YY}U^{\text{app}} - 1)$, and that a decomposition of $\partial_{YY}U^{\text{app}} - 1$ is given in Lemma 4.1. We also recall that the commutator \mathcal{C} is defined by $\mathcal{C} = [L_U^{-1}, \partial_s + b/2Y\partial_Y] - bL_U^{-1}$, and that according to Lemma 2.8,

$$\mathcal{C}[W] = - \left(\mathcal{D} \int_0^Y \frac{W}{U^2} \right)_Y + 2 \left(U \int_0^Y \frac{W}{U^3} \mathcal{D} \right)_Y.$$

Lemma 4.12. *Let $W \in \mathcal{C}^2(\mathbb{R}_+)$ be arbitrary and such that $W = O(Y^2)$ for $Y \ll 1$. Then*

$$\begin{aligned} \mathcal{C}[W] & = 2L_U^{-1} \left(\frac{\mathcal{D}}{U} W \right) - \left(\frac{\mathcal{D}}{U} \int_0^Y L_U^{-1} W \right)_Y, \\ \partial_{YY} \mathcal{C}[W] & = \frac{\mathcal{D}}{U} \partial_Y^2 L_U^{-1} W + \partial_Y \frac{\mathcal{D}}{U} \left[\partial_Y L_U^{-1} W - 2 \frac{U_Y}{U^2} W - 4U_{YY} \int_0^Y \frac{W}{U^2} \right] \\ & \quad + \partial_Y^2 \frac{\mathcal{D}}{U} \left[-3L_U^{-1} W + 2 \frac{W}{U} \right] - \partial_Y^3 \frac{\mathcal{D}}{U} \int_0^Y L_U^{-1} W \\ & \quad + 2\partial_Y^3 U \int_0^Y \frac{W \mathcal{D}}{U^3} - 2\partial_Y^3 U \frac{\mathcal{D}}{U} \int_0^Y \frac{W}{U^2}. \end{aligned}$$

Remark 4.13. In the estimations of E_1 and E_2 , we will use the form of $\partial_{YY}\mathcal{C}[W]$ with $W = \partial_Y^2 V$ and $W = \partial_Y^2 \mathcal{L}_U V$ respectively. Notice in particular that using Corollary 4.3, for any weight w with a strong polynomial decay for $Y \geq s^\beta$ for some $\beta \in [1/4, 2/7]$,

$$\begin{aligned} & \left| \int_0^\infty \frac{\mathcal{D}}{U} (\partial_Y^2 L_U^{-1} W) (\partial_Y^2 L_U^{-1} W) w \right| \\ & \leq Cb \int_0^\infty \frac{1}{1+Y} (\partial_Y^2 L_U^{-1} W)^2 w + \left\| \frac{\mathcal{D}}{U} \right\|_{L^\infty(Y \gtrsim s^{1/3})} \int_{cs^{1/3}}^\infty (\partial_Y^2 L_U^{-1} W)^2 w. \end{aligned}$$

In order to control the tail of the integral, we will use lower order estimates. More precisely, we will use a control of $\partial_Y W$ in $L_{s,Y}^2$ (with appropriate weights in s and Y). We refer to Lemma 4.15 for details.

Proof. The first formula follows easily by recalling the definition of L_U^{-1} and noticing that

$$\int_0^Y \frac{W}{U^2} = \frac{1}{U} \int_0^Y L_U^{-1} W.$$

The second formula is a consequence of Lemma A.1. Notice indeed that

$$\begin{aligned} \partial_Y^2 L_U^{-1} \left(\frac{\mathcal{D}}{U} W \right) &= \partial_Y^3 U \int_0^Y \frac{\mathcal{D}W}{U^3} + \partial_Y^2 U \frac{\mathcal{D}W}{U^3} - \frac{U_Y}{U^2} \partial_Y \left(\frac{\mathcal{D}W}{U} \right) + \frac{1}{U} \partial_Y^2 \left(\frac{\mathcal{D}W}{U} \right) \\ &= \partial_Y^3 U \int_0^Y \frac{\mathcal{D}W}{U^3} + \frac{\mathcal{D}}{U} \partial_Y^2 W - \partial_Y^3 U \frac{\mathcal{D}}{U} \int_0^Y \frac{W}{U^2} \\ &\quad - \frac{U_Y}{U^2} \partial_Y \left(\frac{\mathcal{D}}{U} \right) W + \frac{2}{U} \partial_Y \left(\frac{\mathcal{D}}{U} \right) \partial_Y W + \frac{1}{U} \partial_Y^2 \left(\frac{\mathcal{D}}{U} \right) W. \end{aligned}$$

Then, writing

$$\frac{\partial_Y W}{U} = \partial_Y L_U^{-1} W - U_{YY} \int_0^Y \frac{W}{U^2}$$

and gathering the terms, we obtain the expression announced in Lemma 4.12 □

4.5 Useful inequalities

4.5.1 Evaluation of some remainder terms

In equations (2.19), (2.20), several terms behave heuristically like

$$b \frac{Y}{U} f,$$

where $f = \mathcal{L}_U V$ in (2.19) and $f = \mathcal{L}_U^2 V$ in (2.20). This is for instance the case for the main order commutator term in (2.19) (see Remark 4.13) or for the remainder term

$$\frac{b}{2} \mathcal{L}_U L_U^{-1} \left(YV - \frac{Y^2}{2} V_Y \right).$$

(Recall that L_U^{-1} behaves like a division by U and ∂_Y like a division by Y .) Therefore it is useful to have a systematic way to estimate such remainder terms. To that end, let us first recall that the L^∞ estimates given by Proposition 2.16 ensure that

$$Y + \frac{Y^2}{2} - \frac{M_2}{12}bY^4 \leq U \leq Y + \frac{Y^2}{2} \quad \forall Y \in [0, cs^{1/3}],$$

so that there exist constants $C, c > 0$ such that

$$\frac{Y}{U} \leq \frac{C}{1+Y} \quad \forall Y \leq cs^{1/3}.$$

We then have the following result:

Lemma 4.14. *Assume that*

$$\frac{1}{2s} \leq b \leq \frac{2}{s} \quad \forall s \in [s_0, s_1].$$

Define the weight $w(s, Y) := Y^{-a}(1 + s^{-\beta}Y)^{-m}$ for some $a > 0, m > 1, \beta > 1/4$.

Let $\delta > 0$ arbitrary. Then there exists $s_0 > 0$, depending on δ, β, m , such that the following properties holds: For all $f \in L^\infty(\mathbb{R}_+)$ such that $f = O(Y)$ for $Y \ll 1$, for all $s \geq s_0$,

$$b \int_0^\infty \frac{1}{1+Y} f^2 w \leq \delta b \int_0^\infty f^2 w + \delta \int_0^\infty \frac{f^2}{U^2} w.$$

Proof. We split the integral in two, distinguishing between $Y \leq \delta^{-1}$ and $Y \geq \delta^{-1}$. First, it is clear that

$$\int_{\delta^{-1}}^\infty \frac{1}{1+Y} f^2 w_1 \leq \delta \int_{\delta^{-1}}^\infty f^2 w_1.$$

Thus we focus on the set $Y \leq \delta^{-1}$. On this set, we have

$$\frac{1}{U^2} \geq \frac{1}{Y^2 (1 + \frac{Y}{2})^2},$$

so that

$$s \geq 2\delta^{-3}(1 + \delta^{-1}) \Rightarrow \frac{b}{1+Y} \leq \frac{\delta}{U^2} \quad \forall Y \in [0, \delta^{-1}].$$

□

4.5.2 Control of integral tails

Lemma 4.15. *Assume that U satisfies the following assumptions:*

$$\begin{aligned} U(s, Y) &\leq Y + \frac{Y^2}{2}, \\ U_Y &\leq 1 + Y, \\ |U_{YY}| &\leq C. \end{aligned}$$

Let p, p_0 be positive weights given by

$$p(s, Y) = Y^{-k}(1 + s^{-\beta}Y)^{-m}, \quad p_0(s, Y) = Y^{-k_0}(1 + s^{-\beta_0}Y)^{-m_0},$$

for some $k, k_0 \geq 0, m \geq m_0, \beta < \beta_0$. Let $\alpha_0 > \beta_0$.

Let $P > 0$ be arbitrary. Then there exists $m_P \geq 1$ (depending on $\alpha_0, \beta, \beta_0, k, k_0$) such that if and $m_0, m - m_0 \geq m_P$, then for all $\delta > 0$:

- For $a > 0$, for all $W \in W^{3,\infty}$ such that $W = O(Y^2)$ for $Y \ll 1$,

$$\begin{aligned} & \int_{cs^{\alpha_0}}^{\infty} (\partial_Y^2 L_U^{-1} W)^2 p \\ & \leq \delta \int_0^{\infty} \frac{(\partial_Y^3 L_U^{-1} W)^2}{U} p + \frac{s^{-P}}{\delta} \left(\int_0^{\infty} \frac{(\partial_Y W)^2}{U} p_0 + \int_0^{\infty} W^2 p_0 + C_a \int_0^1 \frac{W^2}{Y^{3+a}} \right); \end{aligned}$$

- Let $\alpha_1 \leq \beta$, and let $\chi \in C_0^\infty(\mathbb{R})$ be a cut-off function such that $\chi \equiv 1$ in a neighbourhood of zero. Then

$$\begin{aligned} & \int_{cs^{\alpha_0}}^{\infty} U \left(\int_0^Y (1 - \chi) \left(\frac{Y}{s^{\alpha_1}} \right) \frac{\partial_Y^2 L_U^{-1} W}{U^2} \right)^2 p \\ & \leq C s^{-P} \left(\int_0^{\infty} W^2 p_0 + \int_0^1 \frac{W^2}{Y^{3+a}} + \int_0^{\infty} \frac{(\partial_Y W)^2}{U} p_0 \right). \end{aligned}$$

,

Remark 4.16. Notice that a similar estimate also holds for quantities such as

$$\int_{cs^{\alpha_0}}^{\infty} (\partial_Y^3 L_U^{-1} W) (\partial_Y^2 L_U^{-1} W) p.$$

Indeed, the integral above can be transformed after a straightforward integration by parts into a quantity similar to the one handled in the Lemma (provided $p(cs^{\alpha_0}) = 0$, which we can always assume without loss of generality, up to the addition of a cut-off function.)

Remark 4.17. We will use these estimates in the next section and we will make the following choices

- $\alpha_0 = 1/3, \alpha_1 = 1/4$;
- $W = \partial_Y^2 V$, $p = w_1$, $p_0 = w_0 = Y^{-a}(1 + s^{-\beta_0})^{-m_0}$, (estimate on E_1 from Proposition 2.9);
- $W = \partial_Y^2 \mathcal{L}_U V$, $p = w_2$, $p_0 = w_1$ (estimate on E_2 from Proposition 2.13).

Proof. Let $\chi \in C_0^\infty(\mathbb{R})$ be a non-negative cut-off function such that $\chi \equiv 1$ in $\subset [-c/2, c/2]$ and $\text{Supp } \chi \subset [-c, c]$. Then

$$\begin{aligned} \int_{cs^{\alpha_0}}^{\infty} (\partial_Y^2 L_U^{-1} W)^2 p & \leq \int_0^{\infty} (1 - \chi) \left(\frac{Y}{s^{\alpha_0}} \right) (\partial_Y^2 L_U^{-1} W)^2 p \\ & = - \int_0^{\infty} \partial_Y L_U^{-1} W \partial_Y \left(\partial_Y^2 L_U^{-1} W (1 - \chi) \left(\frac{Y}{s^{\alpha_0}} \right) p \right). \end{aligned}$$

We then estimate each term in the right-hand side separately. For instance,

$$\begin{aligned} & \int_0^{\infty} |\partial_Y L_U^{-1} W| |\partial_Y^3 L_U^{-1} W| (1 - \chi) \left(\frac{Y}{s^{\alpha_0}} \right) p \\ & \leq \delta \int_0^{\infty} \frac{(\partial_Y^3 L_U^{-1} W)^2}{U} p + \frac{1}{4\delta} \int_0^{\infty} (\partial_Y L_U^{-1} W)^2 U (1 - \chi) \left(\frac{Y}{s^{\alpha_0}} \right) p. \end{aligned}$$

Then, recalling that

$$|\partial_Y L_U^{-1} W| \leq \int_0^Y \frac{|W|}{U^2} + \frac{|\partial_Y W|}{U},$$

we infer that

$$\int_0^\infty (\partial_Y L_U^{-1} W)^2 U (1 - \chi) \left(\frac{Y}{s^{\alpha_0}} \right) p \leq 2 \int_{\frac{cs^{\alpha_0}}{2}}^\infty \frac{(\partial_Y W)^2}{U} p + \int_{\frac{cs^{\alpha_0}}{2}}^\infty \left(\int_0^Y \frac{|W|}{U^2} \right)^2 U p.$$

If $Y \gtrsim s^{\alpha_0}$, write

$$\int_0^Y \frac{|W|}{U^2} \leq C_a \left(\int_0^1 \frac{W^2}{Y^{3+a}} \right)^{1/2} + C \left(\int_0^Y W^2 p_0 \right)^{1/2} \left(\int_0^Y p_0^{-1} \right)^{1/2}.$$

Using the fact that $\|U(s)\|_\infty = O(s)$ together with the assumptions on p and p_0 , we obtain the desired result.

As for the other term, we have

$$\begin{aligned} & - \int_0^\infty \partial_Y L_U^{-1} W \partial_Y^2 L_U^{-1} W \partial_Y \left((1 - \chi) \left(\frac{Y}{s^{\alpha_0}} \right) p \right) \\ &= \frac{1}{2} \int_0^\infty (\partial_Y L_U^{-1} W)^2 \partial_{YY} \left((1 - \chi) \left(\frac{Y}{s^{\alpha_0}} \right) p \right) \\ &\leq C s^{-2/3} \int_{cs^{\alpha_0}/2}^\infty (\partial_Y L_U^{-1} W)^2 p, \end{aligned}$$

which is evaluated with the same estimate as above.

As for the second estimate, notice that thanks to the cut-off function χ , we can integrate by parts without having to deal with boundary terms, so that

$$\int_0^Y (1 - \chi) \left(\frac{Y}{s^{\alpha_1}} \right) \frac{\partial_Y^2 L_U^{-1} W}{U^2} = - \int_0^Y \partial_Y L_U^{-1} W \partial_Y \left[(1 - \chi) \left(\frac{Y}{s^{\alpha_1}} \right) \frac{1}{U^2} \right]. \quad (4.11)$$

Recall that

$$\partial_Y L_U^{-1} W = \frac{\partial_Y W}{U} + U_{YY} \int_0^Y \frac{W}{U^2}.$$

Then the integral in the right-hand side of (4.11) is bounded by

$$\left(\int_0^\infty W^2 p_0 + \int_0^1 \frac{W^2}{Y^{3+a}} + \int_0^\infty \frac{(\partial_Y W)^2}{U} p_0 \right)^{1/2} \left(\int_0^Y Q(s, Y) p_0^{-1} \right)^{1/2},$$

for some function Q that can be computed explicitly and that has at most polynomial growth in s and Y . We conclude as before by choosing m sufficiently large. \square

4.5.3 A special case of Hardy inequalities

We will often use the weighted Hardy inequality from Lemma 4.6 in the following case:

Lemma 4.18. *Let $k \geq 2$ be arbitrary, and let $w = w_i$ for $i = 1, 2$. Then there exists a constant $C = C(k, m_i, a)$, independent of s , such that the following inequalities hold: if s is large enough, then for any $f \in H_{loc}^1(\mathbb{R}_+)$ such that $f(0) = 0$,*

$$\int_0^\infty \frac{1}{(1+Y)^k} f^2 w \leq C \int_0^\infty \frac{1}{(1+Y)^{k-2}} (\partial_Y f)^2 w,$$

$$\int_0^\infty \frac{1}{Y^k} f^2 w \leq C \int_0^\infty \frac{1}{Y^{k-2}} (\partial_Y f)^2 w \quad (\text{provided } f = O(Y^{k/2}) \text{ for } 0 < Y \ll 1).$$

Remark 4.19. *Obviously the Lemma can be extended to weights of the form $Y^{-k}(1+Y)^{-l}w$ with $k+l \geq 2$.*

Proof. We focus on the first inequality, since the second one goes along the same lines (and is in fact slightly easier). Lemma 4.6 entails that

$$\int_0^\infty \frac{1}{(1+Y)^k} f^2 w \leq C_H \int_0^\infty \frac{1}{(1+Y)^{k-2}} (\partial_Y f)^2 w$$

where

$$C_H = 4 \sup_{r>0} \left(\int_r^\infty \frac{w}{(1+Y)^k} \right) \left(\int_0^r (1+Y)^{k-2} w^{-1} \right).$$

We distinguish between the cases $r < 1$ and $r > 1$. If $Y \leq r < 1$, then for s large enough $w^{-1} \leq 2Y^a$, and

$$\left(\int_r^\infty \frac{w}{(1+Y)^k} \right) \left(\int_0^r (1+Y)^{k-2} w^{-1} \right) \leq C_{k,a}.$$

If $r > 1$, then writing $\int_0^r = \int_0^1 + \int_1^r$, we obtain

$$\begin{aligned} & \left(\int_r^\infty \frac{w}{(1+Y)^k} \right) \left(\int_0^r (1+Y)^{k-2} w^{-1} \right) \\ & \leq 2 \left(\int_1^\infty \frac{Y^{-a}}{(1+Y)^k} \right) \left(\int_0^1 (1+Y)^{k-2} Y^a \right) + 2^{k-2} \left(\int_r^\infty \frac{w}{Y^k} \right) \left(\int_1^r Y^{k-2} w^{-1} \right) \\ & \leq C_{k,a} + C_k \left(\int_{s^{-\beta r}}^\infty Z^{-k-a} (1+Z)^{-m} dZ \right) \left(\int_{s^{-\beta}}^{s^{-\beta r}} Z^{k+a-2} (1+Z)^m dZ \right), \end{aligned}$$

so that

$$C_H \leq C_{k,a} + C_k \sup_{r'>0} \left(\int_{r'}^\infty Z^{-k-a} (1+Z)^{-m} dZ \right) \left(\int_0^{r'} Z^{k+a-2} (1+Z)^m dZ \right) \leq C_{m,k,a}.$$

□

5 Sequence of estimates on V

The goal of this section is to prove the energy estimates of Propositions 2.9 and 2.13. To that end, we rely on the transport/diffusion version of the rescaled Prandtl equation, namely (2.16). We will use extensively the tools introduced in section 4. Throughout the section, we

will assume that U satisfies the following pointwise L^∞ estimates: there exists constants c, C such that for all $Y \in [0, cs^{1/3}]$, for all $s \in [s_0, s_1]$,

$$\begin{aligned} Y + \frac{Y^2}{2} - \frac{M_2}{12}bY^4 &\leq U(s, Y) \leq Y + \frac{Y^2}{2}, \\ 1 + Y - \frac{M_2}{4}bY^3 &\leq U_Y(s, Y) \leq 1 + Y, \\ -M_2bY^2 &\leq U_{YY} - 1 \leq 0. \end{aligned} \quad (5.1)$$

Furthermore, we assume that there exists a constant M_1 such that

$$-M_1 \leq U_{YY} \leq 1 \quad \forall Y \geq cs^{1/3}$$

It follows in particular that there exists a universal constant \bar{C} such that for all $Y \in [0, cs^{1/3}]$,

$$|V| \leq \bar{C}(1 + M_2)bY^4, \quad |V_Y| \leq \bar{C}(1 + M_2)bY^3, \quad |V_{YY}| \leq \bar{C}(1 + M_2)bY^2. \quad (5.2)$$

We will also assume that

$$\frac{1 - \bar{\epsilon}}{s} \leq b(s) \leq \frac{1 + \bar{\epsilon}}{s}, \quad (5.3)$$

for some small universal $\bar{\epsilon}$ ($\bar{\epsilon} = 1/50$ is sufficient), and that

$$\int_{s_0}^{s_1} s^{13/4}(b_s + b^2)^2 ds \leq J. \quad (5.4)$$

The L^∞ estimates (5.2) imply in particular the following estimates, which will be used repeatedly in the sequel

$$|\mathcal{L}_U V| \leq \bar{C}(1 + M_2)^2 bY, \quad \left| \int_0^Y \mathcal{L}_U V \right| \leq \bar{C}(1 + M_2)bY^2 \quad \forall Y \leq cs^{1/3}, \quad (5.5)$$

and $\mathcal{L}_U V$ is bounded by a polynomial in Y for $Y \geq s^{1/3}$.

Let us recall the definition of the notation O_α (see (4.1)): there exists a constant $C > 0$ such that

$$\begin{aligned} |A(s, Y)| &\leq C|B(s, Y)| \quad \text{for } Y \leq s^\alpha, \\ |A(s, Y)| &\leq Q(s, Y) \quad \text{for } Y \geq s^\alpha, \end{aligned}$$

for some function Q that has at most polynomial growth in s and Y .

Remark 5.1. *This section contains rather heavy calculations: in particular, terms such as $\partial_Y \mathcal{L}_U^2(L_U^{-1}(L_V Y))$ can be expressed as a linear combination of derivatives of V from order zero up to order 6, with coefficients that are rational expressions involving U and its derivatives. However, most of the terms in this expression will often have the same scaling, in the sense that they can all be bounded in L^2 by the leading order term, i.e. the one that has the largest number of derivatives. To obtain such estimates, we use the weighted Hardy inequalities from the previous section (see Lemma 4.18) together with the pointwise bounds on U (5.1). For instance, if $w = w_j$, $j = 1, 2$, then for any $P > 0$, provided m_j is large enough,*

$$\left\| \frac{U_Y}{U} \mathcal{L}_U V \right\|_{L^2(w)} \leq C \left\| \frac{1}{1+Y} \mathcal{L}_U V \right\|_{L^2(w)} + Cs^{-P} \leq C \|\partial_Y \mathcal{L}_U V\|_{L^2(w)} + Cs^{-P}$$

and

$$\left\| \frac{\partial_{YY} V}{U} \right\|_{L^2(w)} = \left\| \mathcal{L}_U V - \frac{U_Y}{U} \int_0^Y \mathcal{L}_U V \right\|_{L^2(w)} \leq C \|\mathcal{L}_U V\|_{L^2(w)} + C s^{-P}.$$

As a consequence, we adopt the following convention: we will write

$$f = g + \text{l.o.t.} \quad \text{in } L^2(w)$$

if $f = g + h$ and there exists a constant C such that $\|h\|_{L^2(w)} \leq C\|g\|_{L^2(w)} + C s^{-P}$ for any $M > 0$ provided m_1, m_2 are chosen large enough.

Eventually, let us recall the definition of the different weights that will be used in the sequel. We will use parameters

$$\frac{1}{4} < \beta_2 < \beta_1 < \beta_0 < \frac{2}{7}$$

and integers $m_2 \gg m_1 \gg m_0$. For $i \in \{0, 1, 2\}$, we set $w_i := Y^{-a}(1 + s^{-\beta_i} Y)^{-m_i}$. The parameter $a > 0$ is a fixed number such that $a \in (0, \bar{a})$, where \bar{a} is the universal constant in Lemma 4.7. The need for the weight w_0 is explained in the following remark:

Remark 5.2 (Role of the different estimates). *In this section, we derive estimates on E_0, E_1, E_2 , where, for $i = 0, 1, 2$,*

$$E_i(s) := \int_0^\infty (\partial_Y^2 \mathcal{L}_U^i V(s))^2 w_i,$$

$$D_i(s) := \int_0^\infty \frac{(\partial_Y^3 \mathcal{L}_U^i V(s))^2}{U} w_i + \int_0^\infty \frac{(\partial_Y^2 \mathcal{L}_U^i V(s))^2}{U^2} w_i.$$

- The estimate on D_0 allows us to have an L^2 control on $\partial_Y^3 V$, which is useful to bound the integral tails stemming from the estimates of E_1, D_1 (see Lemma 4.15). Note that the integral tails in the estimate of (E_0, D_0) are merely handled thanks to the pointwise L^∞ estimates (5.1).
- The estimate on E_1 will be used in the maximum principle argument (see Proposition 2.16).
- The estimate on D_1 will be used (in a quantitative fashion) when we bound the remainder terms on (E_2, D_2) .
- Eventually, E_2, E_1 and D_1 will control $b_s + b^2$ thanks to Lemmas 2.14 and 2.15.

Since the equations on $\partial_Y^2 \mathcal{L}_U^k V$ for $k = 0, 1, 2$ have the same structure, the estimates on E_0, E_1, E_2 go along the same lines. The differences between the three estimates come from the right-hand side of the equation:

- Commutator terms may or may not be present;
- The estimates on the remainder terms are different for each energy estimate.

The reader who merely wishes to understand the energetic structure of the equation may go through paragraph 5.1 only.

Notation

Throughout this section, we will denote by H_1 a constant depending only on $a, m_1, \beta_1, M_1, M_2$, and by H_2 a constant depending on the same parameters and also on m_2, β_2 .

5.1 Preliminary step: estimates on (E_0, D_0)

Let us recall that the purpose of this paragraph is to have an L^2 control of $\partial_Y^3 V$ through D_0 .

First, notice that $\partial_Y^2 V$ is a solution of

$$\partial_s \partial_Y^2 V + \frac{b}{2} Y \partial_Y (\partial_Y^2 V) - \partial_Y^2 L_U^{-1} \partial_Y^2 V = \partial_Y^2 \mathcal{R}.$$

Consider the weight $w_0 := Y^{-a}(1 + s^{-\beta_0} Y)^{-m_0}$, for some $a \in]0, \bar{a}]$, where \bar{a} is defined in Lemma 4.7, $\beta_0 \in]1/4, 2/7[$, and $m_0 \geq 1$. The diffusion term

$$- \int_0^\infty (\partial_Y^2 L_U^{-1} \partial_Y^2 V) \partial_Y^2 V w_0$$

is handled by Lemma 4.7, up to a remainder term which we estimate now: we have, for any $P > 0$,

$$\begin{aligned} & \int_{cs^{1/3}}^\infty U \left(\int_0^Y (1 - \chi) \left(\frac{Y}{s^{1/4}} \right) \frac{\partial_Y^2 V}{U^2} \right)^2 w_0 \\ & \leq \bar{C} (1 + M_1)^2 \int_{cs^{1/3}}^\infty \left(Y + \frac{Y^2}{2} \right) \frac{Y^2}{s^2} w_0 \\ & \leq \bar{C} (1 + M_1)^2 s^{-P} \end{aligned}$$

provided m_0 is sufficiently large.

Hence, according to Lemma 4.7, setting

$$\begin{aligned} E_0(s) &:= \int_0^\infty (\partial_Y^2 V)^2 w_0, \\ D_0(s) &:= \int_0^\infty \frac{(\partial_Y^3 V)^2}{U} w_0 + \int_0^\infty \frac{(\partial_Y^2 V)^2}{U^2} w_0 \end{aligned}$$

we have, for any $\delta > 0$, provided s_0 is large enough,

$$- \int_0^\infty (\partial_Y^2 L_U^{-1} \partial_Y^2 V) \partial_Y^2 V w_0 \geq \bar{c} D_0(s) - \delta b E_0(s) - \bar{C} (1 + M_1)^2 s^{-P}$$

Concerning the transport term, we have

$$\int_0^\infty (\partial_s \partial_Y^2 V) \partial_Y^2 V w_0 = \frac{1}{2} \frac{d}{ds} \int_0^\infty (\partial_Y^2 V)^2 w_0 - \frac{1}{2} \int_0^\infty (\partial_Y^2 V)^2 \partial_s w_0$$

and

$$\int_0^\infty Y \partial_Y^3 V \partial_Y^2 V w_0 = -\frac{1}{2} \int_0^\infty (\partial_Y^2 V)^2 (Y w_0)_Y.$$

Combining the two identities and using the expression of w_0 , we infer that

$$\begin{aligned} \int_0^\infty \left(\partial_s \partial_Y^2 V + \frac{b}{2} Y \partial_Y (\partial_Y^2 V) \right) \partial_Y^2 V w_0 &= \frac{1}{2} \frac{d}{ds} \int_0^\infty (\partial_Y^2 V)^2 w_0 - \frac{1-a}{4} b \int_0^\infty (\partial_Y^2 V)^2 w_0 \\ &\quad + m \left(\frac{b}{2} - \frac{\beta_0}{2s} \right) \int_0^\infty (\partial_Y^2 V)^2 \frac{s^{-\beta_0} Y}{1 + s^{-\beta_0} Y} w_0. \end{aligned}$$

Using assumption (5.3) with $\bar{\epsilon} < 1/7$ and $\beta_0 < 2/7$, we infer that the last term is non-negative.

It follows that for all $\delta > 0$, if s_0 is large enough,

$$\frac{dE_0}{ds} - \frac{1-a-\delta}{2} b E_0(s) + 2\bar{c} D_0(s) \leq 2 \int_0^\infty (\partial_Y^2 \mathcal{R}) \partial_Y^2 V w_0 + \bar{C} (1 + M_1)^2 s^{-P}.$$

We now evaluate $\partial_Y^2 \mathcal{R}$. Using Lemma 4.4 together with assumption (5.3), we have

$$\begin{aligned} \partial_Y^2 \mathcal{R} &= (b_s + b^2) \left(\frac{Y^2}{4} + a_4 b Y^5 + 270 a_{10} b^2 Y^8 + 330 a_{11} b^2 Y^9 \right) \chi \left(\frac{Y}{s^{2/7}} \right) \\ &\quad + O_{2/7}(b^4(Y^8 + Y^9)) + \partial_Y^2 L_U^{-1} (O_{2/7}(bY)V + O_{2/7}(bY^2)V_Y + O_{2/7}(b^4Y^{11})), \end{aligned}$$

where the $O(\cdot)$ in the last term of the right-hand side must be understood³ in $W^{2,\infty}$. Recalling the expression of $\partial_Y^2 L_U^{-1}$ (see Lemma A.1) together with the L^∞ estimates (5.2), we obtain, for $Y \leq cs^{2/7}$,

$$\partial_Y^2 \mathcal{R} = O((b_s + b^2)Y^2) + O(b^4(Y^8 + Y^9)) + O\left(\frac{bY^2 \partial_Y^3 V}{U}\right) + \partial_Y^3 U \int_0^Y \frac{O(b^2 Y^5)}{U^2}.$$

All constants appearing in the $O(\cdot)$ depend on M_1, M_2 . Notice that for $Y \leq cs^{2/7}$

$$\int_0^Y \frac{b^2 Y^5}{U^2} = O(b^2 Y^2) \ll \frac{bY^2}{U},$$

and $\partial_Y^3 U = \partial_Y^3 U^{\text{app}} + \partial_Y^3 V = \partial_Y^3 V + O(bY)$. Hence the last term is smaller than the first two on $Y \leq cs^{2/7}$.

Now, for all $\delta > 0$, for $s \geq s_\delta$ large enough and for $m_0 \geq 6 - a$,

$$\begin{aligned} \int_0^\infty |b_s + b^2| Y^2 |\partial_Y^2 V| w_0 &\leq \delta b E_0 + \frac{H_1}{\delta} (b_s + b^2)^2 s^{1+(5-a)\beta_0}, \\ \int_0^{cs^{2/7}} b^2 Y^2 |\partial_Y^2 V| w_0 &\leq \delta b E_0 + \frac{H_1}{\delta} b^3 s^{(5-a)\beta_0}, \\ \int_0^{cs^{2/7}} b Y^2 \frac{|\partial_Y^3 V|}{U} |\partial_Y^2 V| w_0 &\leq \delta D_0 + \frac{1}{4\delta} \int_0^{cs^{2/7}} b^2 \frac{Y^4}{U} |\partial_Y^2 V|^2 w_0 \leq \delta D_0 + \delta b E_0. \end{aligned}$$

We now focus on the set $Y \geq s^{2/7}$. Recalling that $\beta_0 < 2/7$, for any function $\phi = \phi(s, Y)$ such that

$$\phi(s, Y) \leq C Y^{\gamma_1} s^{\gamma_2} \quad \forall Y \geq 0, s \geq s_0,$$

for some explicit $\gamma_1, \gamma_2 \geq 0$, we have, for any $P > 0$,

$$\int_{cs^{2/7}}^\infty \phi(s, Y) w_0 \leq s^{-P}$$

³We say that “ $f = O_\alpha(g)$ in $W^{k,\infty}$ ” if $\partial_Y^j f = O_\alpha(\partial_Y^j g)$ for $0 \leq j \leq k$.

provided m_0 is large enough (depending on $\beta_0, \gamma_1, \gamma_2$ and P). We use a simple Cauchy-Schwartz inequality and control the terms involving $\partial_Y^3 V$ by the dissipation term D_0 . Since we have L^∞ estimates on $V, \partial_Y V, \partial_Y^2 V$, it follows that for all $\delta > 0$

$$\int_{cs^{2/7}}^\infty |\partial_Y^2 \mathcal{R}| |\partial_Y^2 V| w_0 \leq \delta D_0 + \frac{H_1}{\delta} s^{-P}.$$

Eventually, choosing δ small enough (depending on a), we obtain

$$\frac{dE_0}{ds} - \frac{1}{2s} E_0 + \bar{c} D_0 \leq H_1 s^{-3+(5-a)\beta_0} + H_1 (b_s + b^2)^2 s^{1+(5-a)\beta_0}.$$

Thus there exists S_0 , depending on M_1, M_2, a, β_0 , such that if $s \geq S_0$,

$$E_0(s) s^{-1/2} + \int_{s_0}^s t^{-1/2} D_0(t) dt \leq E_0(s_0) s_0^{-1/2} + H_1 + H_1 J s_0^{-\frac{11}{4}+(5-a)\beta_0}. \quad (5.6)$$

We will use this inequality in the sequel in order to have a control of $\partial_Y^3 V$, which is not given by the L^∞ estimates. This will allow us in particular to control the tail of some commutators in the estimate of Proposition 2.9.

Remark 5.3. Notice that the assumption $|\partial_Y^2 V(s_0)| \leq 1 + M_1$ implies that

$$E_0(s_0) \leq (1 + M_1)^2 s_0^{(1-a)\beta_0} \leq s_0^{1/2}$$

is s_0 is sufficiently large (depending on M_1 and β_0). Therefore there is no need to include any additional assumption on $E_0(s_0)$ in the bootstrap argument.

5.2 Estimate on $\partial_Y^2 \mathcal{L}_U V$: proof of Proposition 2.9

We now turn towards the estimate on $\mathcal{L}_U V$. Notice that the main order term in the right-hand side of (2.19) is now $(b_s + b^2)Y$. The lack of decay of this remainder prompts us to differentiate twice (2.19), in order to cancel the linear term. We therefore perform estimates on $g := \partial_Y^2 \mathcal{L}_U V$. Using Lemma 2.8, we have

$$\partial_s g + 2bg + \frac{b}{2} Y \partial_Y g - \partial_{YY} L_U^{-1} g = \partial_Y^2 \mathcal{L}_U \mathcal{R} + \partial_Y^2 \mathcal{C}[\partial_Y^2 V]. \quad (5.7)$$

We multiply (5.7) by $g w_1$, where $w_1 = Y^{-a}(1 + s^{-\beta_1} Y)^{-m_1}$ for some $\beta_1 \in]1/4, \beta_0[$, $a > 0$, and $m_1 \gg m_0 \gg 1$. Using the same computations as in the previous paragraph, we have

$$\int_0^\infty \left(\partial_s g + 2bg + \frac{b}{2} Y \partial_Y g \right) g w_1 \geq \frac{1}{2} \frac{d}{ds} E_1(s) + \frac{7+a}{4} b E_1(s),$$

where

$$E_1(s) := \int_0^\infty g^2 w_1. \quad (5.8)$$

Using Lemma 4.7 and Lemma 4.15, we also have, for any $P > 0$, provided $a \leq \bar{a}$, s_0 large enough (depending on a and β_1) and m_1 large enough,

$$- \int_0^\infty (\partial_{YY} L_U^{-1} g) g w_1 \geq \bar{c} D_1(s) - \frac{1}{8} b E_1(s) - s^{-P} D_0(s),$$

where

$$D_1(s) := \int_0^\infty \frac{(\partial_Y g)^2}{U} w_1 + \int_0^\infty \frac{g^2}{U^2} w_1. \quad (5.9)$$

We now treat independently the other terms, namely

- The commutator term $\partial_Y^2 \mathcal{C}[\partial_Y^2 V]$;
- The remainder term $\partial_Y^2 \mathcal{L}_U \mathcal{R}$.

The commutator term

The goal of this paragraph is to prove that for all $\delta, P > 0$, there exists $s_0 > 0$ such that is $s \geq s_0$,

$$\left| \int_0^\infty \partial_Y^2 \mathcal{C}[\partial_Y^2 V] \partial_Y^2 \mathcal{L}_U V w_1 \right| \leq \delta (bE_1(s) + D_1(s)) + s^{-P} D_0(s) + s^{-P}.$$

We use the formula in Lemma 4.12. We recall that $\mathcal{D} = L_U^{-1}(\partial_{YY} U^{\text{app}} - 1) + \mathcal{L}_U V$, so that \mathcal{D} “contains” two derivatives of V through the term $\mathcal{L}_U V$. Hence each term in

$$\int_0^\infty \partial_Y^2 \mathcal{C}[\partial_Y^2 V] \partial_Y^2 \mathcal{L}_U V w_1$$

is a product of three derivatives of V .

As a consequence, we arrange the terms in $\partial_Y^2 \mathcal{C}[\partial_Y^2 V]$ in the following way:

- The terms with the highest number of derivatives of V are estimated by E_1 or D_1 ;
- The terms with a number of derivatives lower than or equal to 2 are estimated in L^∞ thanks to (5.2).

This strategy will work as long as we do not end up with a product of three terms of the type

$$\partial_Y^{k_1} V \partial_Y^{k_2} V \partial_Y^{k_3} V,$$

with $k_1, k_2, k_3 \geq 3$. Such terms will need to be re-arranged thanks to an integration by parts. However, a quick look at the formula in Lemma 4.12 shows that this situation occurs only for the second term in the right-hand side of the formula giving $\partial_Y^2 \mathcal{C}[\partial_Y^2 V]$, namely

$$\int_0^\infty \partial_Y \frac{\mathcal{D}}{U} \partial_Y \mathcal{L}_U V \partial_Y^2 \mathcal{L}_U V w_1. \quad (5.10)$$

But as we will see, it is easy to overcome the difficulty raised by this term.

We now examine the terms in $\partial_Y^2 \mathcal{C}[\partial_Y^2 V]$ one by one.

- Using the L^∞ estimates (5.2) together with Corollary 4.3, it can be easily checked that

$$\frac{\mathcal{D}}{U} = O_{1/3} \left(b \frac{1}{1+Y} \right).$$

It follows that

$$\int_0^{cs^{1/3}} \left| \frac{\mathcal{D}}{U} \right| (\partial_Y^2 \mathcal{L}_U V)^2 w_1 \leq H_1 b \int_0^{cs^{1/3}} \frac{1}{1+Y} (\partial_Y^2 \mathcal{L}_U V)^2 w_1,$$

which is estimated thanks to Lemma 4.14. The part of the integral for $Y \geq cs^{1/3}$ is estimated thanks to Lemma 4.15, with $p = Y^k w_1$ for some integer k , $p_0 = w_0$ with $m_1 \gg m_0$. It follows from Lemma 4.14 that for all $\delta > 0$, $P \geq 1$, provided s_0 is large enough and m_1 is large enough,

$$\int_0^\infty \left| \frac{\mathcal{D}}{U} \right| (\partial_Y^2 \mathcal{L}_U V)^2 w_1 \leq \delta b E_1 + \delta D_1 + s^{-P} D_0(s) + s^{-P}. \quad (5.11)$$

- We then address the problematic term (5.10). We integrate by parts and obtain

$$\begin{aligned} I_1 &= - \int_0^\infty \frac{\mathcal{D}}{U} (\partial_Y^2 \mathcal{L}_U V)^2 w_1 - \int_0^\infty \frac{\mathcal{D}}{U} \partial_Y \mathcal{L}_U V \partial_Y^3 \mathcal{L}_U V w_1 \\ &\quad - \int_0^\infty \frac{\mathcal{D}}{U} \partial_Y \mathcal{L}_U V \partial_Y^2 \mathcal{L}_U V \partial_Y w_1. \end{aligned}$$

The first term in the right-hand side is the same as in (5.11). In the third term, we use the fact that $|\partial_Y w_1| \leq C_{a,m} Y^{-1} w_1$ together with a weighted Hardy inequality from Lemma 4.18, so that

$$\begin{aligned} &\left| \int_0^\infty \frac{\mathcal{D}}{U} \partial_Y \mathcal{L}_U V \partial_Y^2 \mathcal{L}_U V \partial_Y w_1 \right| \\ &\leq H_1 b \left(\int_0^\infty \frac{(\partial_Y^2 \mathcal{L}_U V)^2}{1+Y} w_1 \right)^{1/2} \left(\int_0^\infty \frac{(\partial_Y \mathcal{L}_U V)^2}{Y^2(1+Y)} w_1 \right)^{1/2} + s^{-P} D_0 + s^{-P} \\ &\leq H_1 b \int_0^\infty \frac{(\partial_Y^2 \mathcal{L}_U V)^2}{1+Y} w_1 + s^{-P} D_0 + s^{-P}. \end{aligned}$$

We eventually evaluate the last term in I_1 . We have, for all $\delta > 0$

$$\left| \int_0^\infty \frac{\mathcal{D}}{U} \partial_Y \mathcal{L}_U V \partial_Y^3 \mathcal{L}_U V w_1 \right| \leq \delta D_1 + \frac{1}{4\delta} \int_0^\infty \frac{\mathcal{D}^2}{U} (\partial_Y \mathcal{L}_U V)^2 w_1.$$

Using Corollary 4.3, we have

$$\frac{\mathcal{D}^2}{U} = O_{1/3} \left(\frac{b^2 Y}{1+Y} \right) = O_{1/3} \left(\frac{b}{Y^2(1+Y)} \right).$$

Using a Hardy inequality from Lemma 4.18 together with Lemma 4.14, we deduce that the part of the integral bearing on $Y \lesssim s^{1/3}$ is lower than $\delta D_1 + \delta b E_1$ for $s \geq s_0$ large enough. The part of the integral bearing on $Y \gtrsim s^{1/3}$ is handled thanks to Lemma 4.15, recalling the form of $\partial_Y \mathcal{L}_U^{-1}$ (see Lemma A.1). It is therefore smaller than $\delta D_1 + s^{-P} D_0 + s^{-P}$, for any $\delta, P > 0$, provided s_0 is large enough.

- We then treat simultaneously the next three terms, namely

$$\begin{aligned} &\int_0^\infty \partial_Y \frac{\mathcal{D}}{U} \left(-2 \frac{U_Y}{U^2} \partial_Y^2 V - 4 U_{YY} \int_0^Y \frac{\partial_Y^2 V}{U^2} \right) \partial_Y^2 \mathcal{L}_U V w_1, \\ &\int_0^\infty \partial_Y^2 \frac{\mathcal{D}}{U} \left[-3 \mathcal{L}_U V + 2 \frac{\partial_Y^2 V}{U} \right] \partial_Y^2 \mathcal{L}_U V w_1 \text{ and } \int_0^\infty \partial_Y^3 \frac{\mathcal{D}}{U} \left(\int_0^Y \mathcal{L}_U V \right) \partial_Y^2 \mathcal{L}_U V w_1. \end{aligned} \tag{5.12}$$

The overall idea is to decompose $\partial_Y^k (\mathcal{D}/U)$ for $k = 1, 2, 3$ into a part that is controlled in L^∞ and a part that involves derivatives of V of order 3 or higher (or equivalently, derivatives of $\mathcal{L}_U V$ of order one or higher). Concerning the part of $\partial_Y^k \mathcal{D}/U$ that is controlled in L^∞ , we use weighted Hardy inequalities to upper-bound $\partial_Y^2 V, \mathcal{L}_U V$, etc. by $\partial_Y^2 \mathcal{L}_U V$ in L^2 . As for the part of $\partial_Y^k (\mathcal{D}/U)$ that is not controlled in L^∞ , we observe that in the three terms in (5.12), $\partial_Y^2 V, \mathcal{L}_U V$ and $\int_0^Y \mathcal{L}_U V$ are controlled in L^∞ , and we use this L^∞ control to conclude.

Let us now be more specific: it can be easily checked that for $k = 1, 2, 3$,

$$\partial_Y^k \frac{\mathcal{D}}{U} = O_{1/3} \left(b \frac{1}{Y^k(1+Y)} \right) + O_{1/3} \left(\frac{1}{Y(1+Y)} \right) \partial_Y^k \mathcal{L}_U V + \text{l.o.t.}$$

Since we also have L^∞ estimates on $\partial_Y^2 V$, U_Y, U_{YY} and U^{-1} , using Lemma 4.15, we infer that the part of the integrals in (5.12) bearing on $Y \gtrsim s^{1/3}$ is bounded by $s^{-P} + s^{-P} D_0(s) + \delta D_1$ for some $P > 0$ arbitrary provided m_1 is large enough.

We now address the part of the integral bearing on $Y \lesssim s^{1/3}$, and we start with the part of $\partial_Y^k(\mathcal{D}/U)$ that is bounded in L^∞ . We focus on the first integral in (5.12), since the other two are treated in a similar fashion. We recall that

$$\partial_Y^2 V = U \mathcal{L}_U V - U_Y \int_0^Y \mathcal{L}_U V, \quad \int_0^Y \frac{\partial_Y^2 V}{U^2} = \frac{1}{U} \int_0^Y \mathcal{L}_U V,$$

and therefore

$$-2 \frac{U_Y}{U^2} \partial_Y^2 V - 4U_{YY} \int_0^Y \frac{\partial_Y^2 V}{U^2} = O_{1/3}(Y^{-1}) \mathcal{L}_U V + \text{l.o.t.}$$

Using several weighted Hardy inequalities (see Lemma 4.18), it follows that

$$\begin{aligned} \left| \int_0^{cs^{1/3}} O_{1/3} \left(b \frac{1}{Y(1+Y)} \right) \left(-2 \frac{U_Y}{U^2} \partial_Y^2 V - 4U_{YY} \int_0^Y \frac{\partial_Y^2 V}{U^2} \right) \partial_Y^2 \mathcal{L}_U V w_1 \right| \\ \leq H_1 b \int_0^{cs^{1/3}} \frac{1}{1+Y} |\partial_Y^2 \mathcal{L}_U V|^2 w_1. \end{aligned}$$

We then focus on the part of $\partial_Y(\mathcal{D}/U)$ that is not controlled in L^∞ , and that involves $\partial_Y \mathcal{L}_U V$. For that part, we use the L^∞ estimates on V , which entail

$$\left| -2 \frac{U_Y}{U^2} \partial_Y^2 V - 4U_{YY} \int_0^Y \frac{\partial_Y^2 V}{U^2} \right| \leq H_1 b \quad \forall Y \leq cs^{1/3}.$$

It follows that

$$\begin{aligned} \left| \int_0^{cs^{1/3}} O_{1/3} \left(\frac{1}{Y(1+Y)} \right) \partial_Y \mathcal{L}_U V \left(-2 \frac{U_Y}{U^2} \partial_Y^2 V - 4U_{YY} \int_0^Y \frac{\partial_Y^2 V}{U^2} \right) \partial_Y^2 \mathcal{L}_U V w_1 \right| \\ \leq H_1 b \int_0^{cs^{1/3}} \frac{1}{Y(1+Y)} |\partial_Y \mathcal{L}_U V| |\partial_Y^2 \mathcal{L}_U V| w_1. \end{aligned}$$

We conclude once again using a weighted Hardy inequality.

We then treat the other two integrals in (5.12) using similar arguments. We conclude that for any $\delta, P > 0$, provided $m_1 \gg m_0$ and s_0 is large enough, the three integrals in (5.12) are bounded by

$$\delta D_1 + \frac{H_1}{\delta} \left(b \int_0^\infty \frac{1}{1+Y} |\partial_Y^2 \mathcal{L}_U V|^2 w_1 + s^{-P} + s^{-P} D_0 \right),$$

where the term D_1 stems from the bound of the third integral in (5.12), which involves $\partial_Y^3 \mathcal{L}_U V$.

- Eventually, we address

$$2 \int_0^\infty \partial_Y^3 U \left(\int_0^Y \frac{\partial_Y^2 V \mathcal{D}}{U^3} - \frac{\mathcal{D}}{U} \int_0^Y \frac{\partial_Y^2 V}{U^2} \right) \partial_Y^2 \mathcal{L}_U V w_1. \quad (5.13)$$

Now, \mathcal{D} is controlled in L^∞ . As in the previous step, we decompose $\partial_Y^3 U$ into a part that is controlled in L^∞ and a part over which we have no L^∞ control. More precisely,

$$\partial_Y^3 U = U \partial_Y \mathcal{L}_U V + O_{1/3}(b(Y + Y^2)).$$

Let us start with the contribution of $O_{1/3}(b(Y + Y^2))$. For that part, we use the control of V and \mathcal{D} in L^∞ to prove that the integral tails for $Y \geq cs^{1/3}$ are $O(s^{-P})$, and Hardy inequalities on the set $Y \leq cs^{1/3}$. We infer that the contribution of this part of the integral to (5.13) is bounded by

$$H_1 b E_1^{1/2} \left(\int_0^\infty \frac{1}{1+Y} (\partial_Y^2 \mathcal{L}_U V)^2 w_1 + s^{-P} \right).$$

We now address the part of (5.13) where $\partial_Y^3 U$ is replaced by $U \partial_Y \mathcal{L}_U V$. For that part, we use the L^∞ control of $\partial_Y^2 V$, \mathcal{D} and U in L^∞ , together with Lemma 4.15 and the control of D_0 to estimate the tails. We infer that this part of (5.13) is bounded by

$$\int_0^{s^{2/7}} b^2 Y^2 |\partial_Y \mathcal{L}_U V| |\partial_Y^2 \mathcal{L}_U V| w_1 + s^{-P} D_0 + s^{-P}.$$

Now, for $Y \leq Cs^{2/7}$, we have $b^2 Y^2 \leq CbY^{-1}(1+Y)^{-1/2}$, so that the integral above is bounded by

$$H_1 b E_1^{1/2} \left(\int_0^\infty \frac{1}{1+Y} (\partial_Y^2 \mathcal{L}_U V)^2 w_1 \right)^{1/2}.$$

Gathering the estimates above and using Lemma 4.14, we conclude that for any $P, \delta > 0$, if m_1, s_0 are large enough, the total commutator term satisfies

$$\left| \int_0^\infty \partial_Y^2 \mathcal{C}[\partial_Y^2 V] \partial_Y^2 \mathcal{L}_U V w_1 \right| \leq \delta b E_1(s) + \delta D_1(s) + s^{-P} + s^{-P} D_0(s). \quad (5.14)$$

The remainder term

We now evaluate

$$\int_0^\infty (\partial_Y^2 \mathcal{L}_U \mathcal{R}) (\partial_Y^2 \mathcal{L}_U V) w_1.$$

We claim that for all $\delta > 0$, for all $P > 0$, provided m_1 is large enough and s_0 is large enough,

$$\begin{aligned} & \left| \int_0^\infty (\partial_Y^2 \mathcal{L}_U \mathcal{R}) (\partial_Y^2 \mathcal{L}_U V) w_1 \right| \\ & \leq \delta (D_1 + bE_1) + \frac{H_1}{\delta} \left(s^{-P} D_0 + s^{-7+(11-a)\beta_1} + (b_s + b^2)^2 s^{-3+(11-a)\beta_1} \right). \end{aligned}$$

We follow the decomposition of Lemma 2.6 and write $\mathcal{R} = \sum_{i=1}^4 \mathcal{R}_i$, as suggested in remark 2.7.

- Recalling that $a_4 = 1/48$, $a_7 = a_4/84$, and a_{10}, a_{11} are defined by (2.10), we have

$$\begin{aligned}\partial_Y^2 \mathcal{R}_1 &= (b_s + b^2) \left(\frac{1}{4} Y^2 + a_4 b Y^5 + \frac{81}{16} a_7 b^2 Y^8 + \frac{9}{10} a_7 b^2 Y^9 \right) \chi \left(\frac{Y}{s^{2/7}} \right) \\ &\quad + \tilde{P}_1(s, Y) (1 - \chi_1) \left(\frac{Y}{s^{2/7}} \right),\end{aligned}$$

for some function \tilde{P}_1 that has at most polynomial growth in s and Y , and some cut-off functions $\chi, \chi_1 \in C_0^\infty(\mathbb{R}_+)$ such that $\chi, \chi_1 \equiv 1$ in a neighbourhood of zero. Using the identity (4.2), we have, up to terms supported in $Y \gtrsim s^{2/7}$,

$$\begin{aligned}\mathcal{L}_U \mathcal{R}_1 &= \frac{1}{2} (b_s + b^2) Y + L_U^{-1} \left[\left(\frac{a_4}{2} (b_s + b^2) b Y^5 \right) \chi \left(\frac{Y}{s^{2/7}} \right) \right] \\ &\quad - (b_s + b^2) L_U^{-1} \left(\left(\frac{61}{16} a_7 b^2 Y^8 - \frac{9}{10} a_7 b^2 Y^9 + 2a_{10} b^3 Y^{10} + \frac{9}{4} a_4 b^3 Y^{12} \right) \chi \left(\frac{Y}{s^{2/7}} \right) \right) \\ &\quad - \frac{1}{2} (b_s + b^2) L_U^{-1} L_V Y - L_U^{-1} \left[\tilde{P}_1(s, Y) (1 - \chi_1) \left(\frac{Y}{s^{2/7}} \right) \right],\end{aligned}$$

and therefore

$$\begin{aligned}\partial_Y^2 \mathcal{L}_U \mathcal{R}_1 (b_s + b^2)^{-1} &= \partial_Y^2 L_U^{-1} \left[\left(\frac{a_4}{2} b Y^5 - \frac{61}{16} a_7 b^2 Y^8 + \frac{9}{10} a_7 b^2 Y^9 - 2a_{10} b^3 Y^{10} - \frac{9}{4} a_4 b^3 Y^{12} \right) \chi \left(\frac{Y}{s^{2/7}} \right) \right] \\ &\quad - \frac{1}{2} \partial_Y^2 L_U^{-1} L_V Y - \partial_Y^2 L_U^{-1} \left[\tilde{P}_1(s, Y) (1 - \chi_1) \left(\frac{Y}{s^{2/7}} \right) \right].\end{aligned}\tag{5.15}$$

Recalling the expression of $\partial_Y^2 L_U^{-1}$ from Lemma A.1, we infer that for $k \geq 5$,

$$\partial_Y^2 L_U^{-1} Y^k = \partial_Y^3 V \int_0^Y \frac{Y^k}{U^2} + O_{1/3} \left(\begin{cases} Y^{k-3} & \text{if } Y \ll 1, \\ Y^{k-4} & \text{if } 1 \lesssim Y. \end{cases} \right)$$

In a similar way, we have

$$\begin{aligned}\partial_Y^2 L_U^{-1} (L_V Y) &= \partial_Y^3 U \int_0^Y \frac{L_V Y}{U^2} - \frac{Y^2}{2} \partial_Y \mathcal{L}_U V \\ &\quad + V_{YY} \frac{Y^2 U_Y - 2Y U}{2U^2} + V_Y \frac{2U - Y^2 U_{YY}}{2U^2} + V \frac{U_{YY} Y - U_Y}{U^2} \\ &\quad + \frac{Y^2 U_{YY}}{2U} \int_0^Y \mathcal{L}_U V.\end{aligned}\tag{5.16}$$

We write $\partial_Y^3 U = \partial_Y^3 U^{\text{app}} + \partial_Y^3 V$ and replace $\partial_Y^3 V$ by the formula (A.2) in Appendix A. Using the L^∞ estimate on $\partial_Y^2 V$, we obtain

$$\partial_Y^2 L_U^{-1} (L_V Y) = O_{2/7}(Y^2) \partial_Y \mathcal{L}_U V + O_{2/7}(Y) \mathcal{L}_U V + O_{2/7}(1) \int_0^Y \mathcal{L}_U V + \text{l.o.t.}$$

Gathering all the terms, we get, for $Y \gg 1$,

$$\partial_Y^2 \mathcal{L}_U \mathcal{R}_1 = (b_s + b^2) \left(O_{2/7}(bY) + \sum_{i=0}^2 O_{2/7}(Y^i) \partial_Y^i \int_0^Y \mathcal{L}_U V + \text{l.o.t.} \right).$$

Then, noticing that for $Y \leq s^{2/7}$ we have $bY^2 \leq Y^{-1}(1+Y)^{-1/2}$ and using Hardy inequalities from Lemma 4.18, we infer that for any $P > 0$, provided m_1 and S_0 are large enough,

$$\begin{aligned}
& \left| \int_0^\infty \partial_Y^2 \mathcal{L}_U \mathcal{R}_1 \partial_Y^2 \mathcal{L}_U V w_1 \right| \\
& \leq H_1 |b_s + b^2| E_1^{1/2} \left(b s^{(3-a)\beta_1/2} + b^2 s^{(11-a)\beta_1/2} \right) \\
& \quad + H_1 s |b_s + b^2| E_1^{1/2} \left(\int_0^{s^{2/7}} \frac{1}{Y^2(1+Y)} (\partial_Y \mathcal{L}_U V)^2 w_1 + s^{-P} + s^{-P} D_0 \right)^{1/2} \\
& \leq \delta b E_1 + \frac{H_1}{\delta} (b_s + b^2)^2 s^{-3+(11-a)\beta_1} \\
& \quad + \frac{H_1}{\delta} s^3 |b_s + b^2|^2 \int_0^\infty \frac{1}{1+Y} (\partial_Y^2 \mathcal{L}_U V)^2 w_1 + s^{-P} D_0.
\end{aligned}$$

- We use the same type of estimates for the term \mathcal{R}_2 , and we find

$$\begin{aligned}
& \left| \int_0^\infty \partial_Y^2 \mathcal{L}_U \mathcal{R}_2 \partial_Y^2 \mathcal{L}_U V w_1 \right| \\
& \leq H_1 b^4 E_1^{1/2} s^{(11-a)\beta_1/2} + \delta b E_1 \\
& \leq \delta b E_1 + \frac{H_1}{\delta} s^{-7+(11-a)\beta_1} + s^{-P} D_0.
\end{aligned}$$

- We then address the term \mathcal{R}_3 . We recall that using (5.16) and (A.1), for $Y \leq cs^{1/3}$, for $k \in \{2, 3, 4\}$

$$\begin{aligned}
\partial_Y L_U^{-1}(L_V Y) &= O_{1/3}(bY^2), \quad L_U^{-1}(L_V Y) = O_{1/3}(bY^3), \\
\partial_Y^k L_U^{-1}(L_V Y) &= \sum_{i=0}^k O_{1/3}(Y^{i-k+2}) \partial_Y^i \int_0^Y \mathcal{L}_U V + O_{1/3} \left(\frac{1}{Y^{k-1}(1+Y)} \right) V_Y + \text{l.o.t.}
\end{aligned}$$

Notice also that $b^3 Y^7 \lesssim bY$ for $Y \leq cs^{1/3}$, so that we can treat $b^3 L_V Y^7$ as a perturbation of $b L_V Y$. In a similar way, for $k \in \{0, \dots, 3\}$,

$$\partial_Y^k (-a_4 b Y^4 - a_7 b^2 Y^7 + a_{10} b^3 Y^{10} + a_{11} b^3 Y^{11}) = O(bY^{4-k}) \quad \text{for } Y \lesssim s^{2/7},$$

so that

$$b^3 \partial_Y^2 L_U^{-1} L_{-a_4 b Y^4 - a_7 b^2 Y^7 + a_{10} b^3 Y^{10} + a_{11} b^3 Y^{11}} Y^7 = O_{2/7}(b^4 Y^7) + O_{2/7}(b^4 Y^8) \partial_Y^3 V.$$

It follows that

$$\begin{aligned}
\partial_Y^2 \mathcal{L}_U \mathcal{R}_3 &= \sum_{i=0}^4 O_{2/7}(bY^{i-4}) \partial_Y^i \int_0^Y \mathcal{L}_U V \\
& \quad + O_{2/7}(b^2(Y+Y^2)) \partial_Y \mathcal{L}_U V + O_{2/7}(b^2) \int_0^Y \mathcal{L}_U V \\
& \quad + O_{2/7} \left(\frac{b}{Y^4(1+Y)^2} \right) V_Y + \text{l.o.t.} + O_{2/7}(b^4 Y^3).
\end{aligned}$$

Using Hardy inequalities together with Lemma 4.15, it is easily proved that for $0 \leq i \leq 4$, for any $P > 0$, provided m_1 is large enough,

$$\begin{aligned} \left| \int_0^\infty O_{2/7}(bY^{i-4}) \partial_Y^i \int_0^Y \mathcal{L}_U V \partial_Y^2 \mathcal{L}_U V w_1 \right| &\leq \delta(bE_1 + D_1) + s^{-P} D_0(s) + s^{-P}, \\ \left| \int_0^\infty O_{2/7}(b^2(Y + Y^2)) \partial_Y \mathcal{L}_U V \partial_Y^2 \mathcal{L}_U V w_1 \right| &\leq \delta b E_1 + s^{-P} D_0(s) + s^{-P}, \\ \left| \int_0^\infty O_{2/7}(b^2) \int_0^Y \mathcal{L}_U V \partial_Y^2 \mathcal{L}_U V w_1 \right| &\leq \delta b E_1 + s^{-P} D_0(s) + s^{-P}, \end{aligned}$$

and

$$\begin{aligned} &\left| \int_0^\infty O_{2/7} \left(\frac{b}{Y^4(1+Y)^2} \right) V_Y \partial_Y^2 \mathcal{L}_U V w_1 \right| \\ &\leq H_1 b \left(\int_0^\infty \frac{1}{1+Y} (\partial_Y^2 \mathcal{L}_U V)^2 w_1 \right)^{1/2} \left(\int_0^\infty \frac{1}{Y^8(1+Y)^3} V_Y^2 w_1 + s^{-P} \right)^{1/2} \\ &\leq H_1 b \int_0^\infty \frac{1}{1+Y} (\partial_Y^2 \mathcal{L}_U V)^2 w_1 + s^{-P}. \end{aligned}$$

Using Lemma 4.14, we end up with

$$\left| \int_0^\infty \partial_Y^2 \mathcal{L}_U \mathcal{R}_3 \partial_Y^2 \mathcal{L}_U V w_1 \right| \leq \delta(D_1 + bE_1) + (s^{-P} D_0 + s^{-P}).$$

- The term \mathcal{R}_4 is easily treated thanks to Lemma 4.15. More precisely, using Appendix A, it can be proved that

$$\partial_Y^2 \mathcal{L}_U \mathcal{R}_4 = (P_1(s, Y) + P_2(s, Y) \partial_Y \mathcal{L}_U V + P_3(s, Y) \partial_Y^2 \mathcal{L}_U V) (1 - \bar{\chi}) \left(\frac{Y}{s^{2/7}} \right),$$

where P_1, P_2, P_3 are functions that have at most polynomial growth in s and Y , and $\bar{\chi} \in C_1^\infty$ is identically equal to 1 in a neighbourhood of zero. We infer that for any $P > 0$, provided m_1 and s_0 are large enough,

$$\left| \int_0^\infty (\partial_Y^2 \mathcal{L}_U \mathcal{R}_4) \partial_Y^2 \mathcal{L}_U V w_1 \right| \leq C s^{-P} + s^{-P} D_0.$$

We now gather all the terms. Notice that since $\beta_1 > 1/4$, $5 + (3 - a)\beta_1 < -7 + (11 - a)\beta_1$. We end up with the following estimate: for all $\delta, P > 0$, if m_1 is large enough, there exists a constant H_1 , depending on M_1, M_2, a, β_1 and m_1 and a constant S_0 , depending on the same parameters and also on δ , such that if $s \geq S_0$, then

$$\begin{aligned} &\left| \int_0^\infty (\partial_Y^2 \mathcal{L}_U \mathcal{R}) \partial_Y^2 \mathcal{L}_U V w_1 \right| \\ &\leq \delta(D_1 + bE_1) + \frac{H_1}{\delta} \left[(b_s + b^2)^2 s^{-3+(11-a)\beta_1} + s^3 (b_s + b^2)^2 E_1 + s^{-7+(11-a)\beta_1} \right] + s^{-P} D_0. \end{aligned} \tag{5.17}$$

Conclusion

Gathering the estimates (5.14) and (5.17), we infer that for $a > 0$ sufficiently small, for any $P > 0$, for $1/4 < \beta_1 < 2/7$ and for m_1 sufficiently large, $s_0 \leq S_0$, we have, for $s \in [s_0, s_1]$,

$$\frac{d}{ds} E_1 + \left(\frac{7}{2} b - H_1 s^3 (b_s + b^2)^2 \right) E_1 + \bar{c} D_1 \leq H_1 (b_s + b^2)^2 s^{-3+(11-a)\beta_1} + H_1 s^{-7+(11-a)\beta_1} + s^{-P} D_0.$$

Since $\beta_1 > 1/4$, we have

$$6 - (11 - a)\beta_1 < \frac{7}{2},$$

and therefore the rate of convergence is limited by the size of the right-hand side. Let $\phi_1(s) := H_1 \int_{s_0}^s \tau^3 (b_\tau + b^2)^2 d\tau$. The assumptions of the Lemma entail that $0 \leq \phi_1(s) \leq H_1 J s_0^{-1/4}$ for all $s \in [s_0, s_1]$. As a consequence, if $s_0 \geq J^4$, we have $0 \leq \phi_1(s) \leq H_1$ for all $s \in [s_0, s_1]$. Using a Gronwall type argument and using the preliminary estimate on D_0 , we obtain, for all $\alpha < 6 - (11 - a)\beta_1$,

$$\begin{aligned} E_1(s) s^\alpha \exp(-\phi_1(s)) + \bar{c} \int_{s_0}^s \tau^\alpha \exp(-\phi_1(\tau)) D_1(\tau) d\tau \\ \leq E_1(s_0) s_0^\alpha + H_1 J s_0^{-1/4} + \frac{H_1}{6 - (11 - a)\beta_1 - \alpha} s_0^{\alpha-6+(11-a)\beta_1}. \end{aligned}$$

Hence, for $s_0 \geq \max(S_0, J^4)$ we obtain (up to a new definition of the constant H_1)

$$E_1(s) s^\alpha + \int_{s_0}^s D_1(\tau) \tau^\alpha d\tau \leq H_1 (1 + E_1(s_0) s_0^\alpha).$$

This completes the proof of the Proposition. \square

We also have the following

Corollary 5.4. *Under the assumptions of Proposition 2.9, we get the following refined L^∞ estimates on V : for all $a \in (0, \bar{a})$, for all β_1 such that*

$$\frac{1}{4} < \beta_1 \leq \frac{1}{4} \frac{11}{11 - a},$$

there holds

$$\begin{aligned} |\partial_Y \mathcal{L}_U V| &= O_{\beta_1} \left(s^{-13/8} Y^{\frac{1+a}{2}} \right), \\ |\mathcal{L}_U V| &= O_{\beta_1} \left(s^{-13/8} Y^{\frac{3+a}{2}} \right), \quad \int_0^Y \mathcal{L}_U V = O_{\beta_1} \left(s^{-13/8} Y^{\frac{5+a}{2}} \right), \\ |\partial_Y^3 V| &= O_{\beta_1} \left(C_1 s^{-13/8} Y^{\frac{3+a}{2}} (1 + Y) \right). \end{aligned}$$

Note that the constants in the O_{β_1} depend on $M_1, M_2, m_1, a, \beta_1$ and J .

As a consequence, setting $C_0 = E_1(s_0) s_0^\alpha$, where α is such that $13/4 < \alpha < 6 - (11 - a)\beta_1$, we infer that there exists a universal constant \bar{H} and an constant C_{m_1} depending only on m_1 such that if $s_0 \geq \max(S_0, J^4, C_0^8)$,

$$|\partial_Y^3 U| \leq \bar{H} b Y \quad \forall Y \in [0, C_{m_1} s^{\beta_1}].$$

Furthermore, there exists a constant C_{a,m_1} depending only on a and m_1 , such that

$$\begin{aligned} \left| \int_0^Y \frac{\partial_Y^2 \mathcal{L}_U V}{U^2} \right| &\leq C_{a,m_1} D_1^{1/2} (1 + s^{-\beta_1} Y)^{\frac{4+a+m_1}{2}} = O_{\beta_1}(D_1^{1/2}), \\ |\partial_Y^2 \mathcal{L}_U V| &\leq C_{a,m_1} D_1^{1/2} (1 + s^{-\beta_1} Y)^{\frac{m_1}{2}} (1 + Y)^{1/2} Y^{(2+a)/2} = O_{\beta_1} \left(D_1^{1/2} (1 + Y)^{1/2} Y^{(2+a)/2} \right). \end{aligned} \quad (5.18)$$

In particular,

$$\partial_Y \mathcal{L}_U^2 V = \frac{\partial_Y^3 \mathcal{L}_U V}{U} + O_{\beta_1}(D_1^{1/2})$$

Proof. As mentioned in Remark 2.11 we can pick $a > 0$, $\beta_1 \in (1/4, 2/7)$ (β_1 depends on a) so that

$$\frac{13}{4} = 6 - \frac{11}{4} < 6 - (11 - a)\beta_1.$$

With this choice of a and β_1 , we have

$$E_1(s) \leq H_1(1 + C_0)s^{-13/4}.$$

A simple Cauchy-Schwartz inequality entails

$$|\partial_Y \mathcal{L}_U V| = \left| \int_0^Y \partial_Y^2 \mathcal{L}_U V \right| \leq E_1^{1/2} \left(\int_0^Y \frac{1}{w_1} \right)^{1/2}. \quad (5.19)$$

Now, setting $C_{m_1} := 2^{1/m_1} - 1 \leq 1$, it is easily checked that for $Y \leq C_{m_1} s^{\beta_1}$, we have

$$\frac{1}{w_1} \leq 2Y^a.$$

The estimates follow, using the formula in equation (A.2) for the one of $\partial_Y^3 V$. Note in particular that for $Y \leq C_{m_1} s^{\beta_1}$,

$$\begin{aligned} |\partial_Y^3 U| &\leq |\partial_Y^3 U^{\text{app}}| + \left(Y + \frac{Y^2}{2} \right) |\partial_Y \mathcal{L}_U V| + \int_0^Y |\mathcal{L}_U V| \\ &\leq \bar{H} \left(bY + (Y + Y^2) Y^{\frac{a+1}{2}} H_1^{1/2} (1 + C_0)^{1/2} s^{-13/8} \right) \\ &\leq \bar{H} bY \left(1 + H_1^{1/2} (1 + C_0)^{1/2} s_0^{\frac{a+3}{2}\beta_1 - \frac{5}{8}} \right). \end{aligned}$$

Now, for $\beta_1 < 2/7$ and a sufficiently small, $\beta_1 < 1/(3+a)$, so that, if $s_0 \geq \max(C_0^8, H_1^8)$,

$$|\partial_Y^3 U| \leq \bar{H} bY \left(1 + H_1^{1/2} (1 + C_0)^{1/2} s_0^{-1/8} \right) \leq \bar{H} bY.$$

Since $\partial_Y^3 V$ has polynomial growth in s and Y according to (5.19), we obtain the estimate on $\partial_Y^3 U$.

The two estimates from (5.18) follow from the Cauchy-Schwartz inequality (see also Remark 4.10): observe that for $Y \leq 1$,

$$\left| \int_0^Y \frac{\partial_Y^2 \mathcal{L}_U V}{U^2} \right| \leq \left(\int_0^1 \frac{(\partial_Y^2 \mathcal{L}_U V)^2}{Y^{3+a}} \right)^{1/2} \left(\int_0^Y \frac{Y^{3+a}}{U^4} \right)^{1/2} \leq C_a D_1^{1/2}.$$

The same type of estimate holds when $Y \geq 1$. As for the estimate on $\partial_Y^2 \mathcal{L}_U V$, we have

$$|\partial_Y^2 \mathcal{L}_U V| = \left| \int_0^Y \partial_Y^3 \mathcal{L}_U V \right| \leq D_1^{1/2} \left(\int_0^Y \frac{U}{w_1} \right)^{1/2},$$

which leads to the result. □

We end this paragraph with a short proof for Lemma 2.12: differentiating once equation (2.19), we have

$$\partial_s \partial_Y \mathcal{L}_U V + \frac{3b}{2} \partial_Y \mathcal{L}_U V + \frac{b}{2} Y \partial_Y^2 \mathcal{L}_U V - \partial_Y \mathcal{L}_U^2 V = \partial_Y \mathcal{L}_U \mathcal{R} + \partial_Y \mathcal{C}[\partial_Y^2 V].$$

We now take the trace of the above equality at $Y = 0$. Since $V = O(Y^7)$ for $Y \ll 1$ by definition of the approximate solution U^{app} , we have

$$\partial_Y \mathcal{L}_U V|_{Y=0} = \partial_Y^2 \mathcal{L}_U V|_{Y=0} = 0,$$

as well as $\partial_Y \mathcal{C}[\partial_Y^2 V]|_{Y=0} = 0$. Hence there remains only

$$\partial_Y \mathcal{L}_U^2 V|_{Y=0} = - \sum_{i=1}^4 \partial_Y \mathcal{L}_U \mathcal{R}_i|_{Y=0}.$$

Once again, it can be easily checked that $\partial_Y \mathcal{L}_U \mathcal{R}_i|_{Y=0} = 0$ for $i = 2, 3, 4$, and that

$$\partial_Y \mathcal{L}_U \mathcal{R}_1|_{Y=0} = a_4(b_s + b^2) \partial_Y \mathcal{L}_U Y^4|_{Y=0}.$$

Now, using (4.2), we have

$$\partial_Y \mathcal{L}_U Y^4 = 12 \partial_Y L_U^{-1} Y^2 = 24 \partial_Y Y + O(bY^3) \text{ for } Y \ll 1.$$

We obtain eventually that

$$\partial_Y \mathcal{L}_U^2 V|_{Y=0} = -\frac{1}{2}(b_s + b^2).$$

Remark 5.5. *It also follows from equation (5.7) and from the computation of $\partial_Y^2 \mathcal{R}$ that for $Y \ll 1$,*

$$\partial_Y^2 \mathcal{L}_U^2 V = O((|b|b_s + b^2| + |\partial_s(b_s + b^2)|)Y^2).$$

In particular, $\mathcal{L}_U^3 V$ is well-defined.

5.3 Estimate on $\partial_Y^2 \mathcal{L}_U^2 V$: proof of Proposition 2.13

We now tackle the estimates on $\partial_Y \mathcal{L}_U^2 V$. The general scheme of proof is the same as the one of Proposition 2.9. There are however a few differences:

- There are now two commutator terms, namely $\partial_Y \mathcal{L}_U \mathcal{C}[\partial_Y^2 V]$ and $\partial_Y \mathcal{C}[\partial_Y^2 \mathcal{L}_U V]$;
- The estimate of the remainder term $\partial_Y^2 \mathcal{L}_U^2 \mathcal{R}$ is more technical since the explicit expression of $\partial_Y^2 \mathcal{L}_U^2$ has much more terms than the one of $\partial_Y^2 \mathcal{L}_U$.

We set $h = \partial_Y^2 \mathcal{L}_U^2 V$ in the rest of the paper, and we have

$$\partial_s h + 4bh + \frac{b}{2} Y \partial_Y h - \partial_Y^2 L_U^{-1} h = \partial_Y^2 (\mathcal{L}_U^2 \mathcal{R} + \mathcal{C}[\partial_Y^2 \mathcal{L}_U V] + \mathcal{L}_U \mathcal{C}[\partial_Y^2 V]). \quad (5.20)$$

In order to derive estimate (2.21), we multiply (5.20) by hw_2 , with $w_2 := Y^{-a}(1 + s^{-\beta_2} Y)^{-m_2}$. We recall that we choose $m_2 \gg m_1 \gg 1$ and $\beta_2 < \beta_1$.

Using the same computations as in the previous paragraphs and recalling the definitions of E_2, D_2 , we have

$$\begin{aligned} & \frac{1}{2} \frac{dE_2}{ds} + \frac{15}{4} bE_2 + \bar{c}D_2 \\ & \leq \int_0^\infty \partial_Y^2 (\mathcal{L}_U^2 \mathcal{R} + \mathcal{C}[\partial_Y^2 \mathcal{L}_U V] + \mathcal{L}_U \mathcal{C}[\partial_Y^2 V]) hw_2 + \bar{C}s^{-P} + s^{-P} D_1. \end{aligned}$$

We now state the main intermediate results allowing us to prove the statement, namely a commutator estimate and a remainder estimate. We then prove each of the statements separately.

Concerning the commutators, we have the same type of estimates as in the proof of Proposition 2.9, which leads to

Lemma 5.6. *Assume that the assumptions of Proposition 2.13 are satisfied. There exist constants H_2, S_0 , depending on $a, m_i, \beta_i, M_1, M_2$ ($i = 1, 2$), such that if $s_0 \geq \max(S_0, C_0^8, J^4)$, for all $\delta > 0$,*

$$\begin{aligned} & \left| \int_0^\infty \int_0^\infty (\partial_Y^2 \mathcal{C}[\partial_Y^2 \mathcal{L}_U V] + \partial_Y^2 \mathcal{L}_U \mathcal{C}[\partial_Y^2 V]) hw_2 \right| \\ & \leq \delta (bE_2 + D_2) + \frac{H_2}{\delta} b^2 (b_s + b^2)^2 + \frac{H_2}{\delta} \left(s^{(7+a)\beta_1} D_1 E_2 + s^3 (b_s + b^2)^2 E_2 + s^{-2} D_1 \right). \end{aligned}$$

We now turn towards the remainder term. We have the following estimate:

Lemma 5.7. *Assume that the assumptions of Proposition 2.13 are satisfied. There exist constants H_2, S_0 , depending on $a, m_i, \beta_i, M_1, M_2$ ($i = 1, 2$), such that if $s_0 \geq \max(S_0, C_0^8, J^4)$, for all $\delta > 0$,*

$$\int_0^\infty \partial_Y^2 \mathcal{L}_U^2 \mathcal{R} \partial_Y^2 \mathcal{L}_U^2 V w_2 \leq \delta (bE_2 + D_2) + \frac{H_2}{\delta} b^2 (b_s + b^2)^2 + \frac{H_2}{\delta} s^{-2} D_1 + \frac{H_2}{\delta} s^{-7+(3-a)\beta_2}.$$

Gathering Lemmas 5.6 and 5.7 and choosing $\delta = \min(1/2, \bar{c}/2)$, we obtain

$$\begin{aligned} & \frac{dE_2}{ds} + 7bE_2 + \frac{\bar{c}}{2} D_2 \\ & \leq H_2 \left(b^2 (b_s + b^2)^2 + s^{(7+a)\beta_1} D_1 E_2 + s^3 (b_s + b^2)^2 E_2 + s^{-2} D_1 + s^{-7+(3-a)\beta_2} \right). \end{aligned}$$

We now multiply the above equation by s^5 and infer

$$\frac{d}{ds} (E_2(s) s^5) - H_2 \left(s^{(7+a)\beta_1} D_1 + s^3 (b_s + b^2)^2 \right) (E_2(s) s^5) \quad (5.21)$$

$$\leq H_2 \left(s^3 (b_s + b^2)^2 + s^3 D_1 + s^{-2+(3-a)\beta_2} \right). \quad (5.22)$$

Define

$$\phi_2(s) := H_2 \int_{s_0}^s \tau^{(7+a)\beta_1} D_1(\tau) + \tau^3(b_\tau + b^2)^2 d\tau.$$

According to Proposition 2.9, since $\beta_1 < 2/7 < 1/3$, we have $(7+a)\beta_1 < 6 - (11-a)\beta_1$ and $7\beta_1 < 13/4$, and therefore, if $s_0 \geq \max(S_0, J^4)$,

$$0 \leq \phi_2(s) \leq H_1 H_2 \left(1 + E_1(s_0) s_0^{13/4}\right).$$

Hence, multiplying (5.21) by $\exp(-\phi_2(s))$ and integrating over $[s_0, s]$, we obtain

$$\begin{aligned} & E_2(s) s^5 \exp(-\phi_2(s)) \\ & \leq E_2(s_0) s_0^5 + H_2 \int_{s_0}^s \left(\tau^3(b_\tau + b^2)^2 + \tau^3 D_1 + \tau^{-2+(3-a)\beta_2} \right) \exp(-\phi_2(\tau)) d\tau. \end{aligned}$$

Now, using assumption (5.4) together with Proposition 2.9 with $\alpha = 3$, we have, for $s_0 \geq \max(S_0, J^4)$,

$$\begin{aligned} \int_{s_0}^s \tau^3(b_\tau + b^2)^2 d\tau & \leq J s_0^{-1/4} \leq 1, \\ \int_{s_0}^s \tau^3 D_1 & \leq H_1(1 + C_0), \\ \int_{s_0}^s \tau^{-2+(3-a)\beta_2} d\tau & \leq \frac{1}{1 - (3-a)\beta_2} s_0^{-1+(3-a)\beta_2} \leq 1. \end{aligned}$$

We infer eventually

$$E_2(s) s^5 \leq \exp(H_2(1 + C_0)) \left[E_2(s_0) s_0^5 + H_2 H_1(1 + C_0) \right] \leq \exp(H_2(1 + C_0)) H_2(1 + C_0).$$

We now turn towards the proofs of the Lemmas.

The commutator terms: proof of Lemma 5.6

We start with the computation of the commutator terms. Looking at equation (5.20), the commutator integrals in the differential inequality for E_2 are

$$\int_0^\infty (\partial_Y^2 \mathcal{C}[\partial_Y^2 \mathcal{L}_U V] + \partial_Y^2 \mathcal{L}_U \mathcal{C}[\partial_Y^2 V]) \partial_Y^2 \mathcal{L}_U^2 V w_2. \quad (5.23)$$

We recall that the heuristics is that up to some corrector terms,

$$|(5.23)| \leq H_2 b \int_0^\infty \frac{1}{1+Y} (\partial_Y^2 \mathcal{L}_U^2 V)^2 w_2,$$

and the right-hand side of the above inequality is then handled by Lemma 4.14. However, there are a few complications, coming from the fact that the trace $\partial_Y \mathcal{L}_U^2 V|_{Y=0}$ is not zero. In the sequel, we will therefore focus on the difficulties and differences with respect to the treatment of the commutators in Proposition 2.9.

• We first consider the first term in the integral of (5.23). This term has the same type of structure as the term

$$\int_0^\infty \partial_Y^2 \mathcal{C}[\partial_Y^2 V] \partial_Y^2 \mathcal{L}_U V w_1,$$

which we treated in the previous section. However, there exists one substantial difference, due to the fact that $\partial_Y \mathcal{L}_U^2 V$ does not vanish at $Y = 0$, so that we cannot write Hardy inequalities for $\partial_Y \mathcal{L}_U^2 V$. To overcome this difficulty, we recall that $\partial_Y \mathcal{L}_U^2 V|_{Y=0} = -\frac{1}{2}(b_s + b^2)$, so that $\partial_Y^2 \mathcal{L}_U V \sim -\frac{1}{2}(b_s + b^2)L_U Y \sim -\frac{1}{4}(b_s + b^2)Y^2$ for $Y \ll 1$, and we write

$$\partial_Y^2 \mathcal{L}_U V = \left(\partial_Y^2 \mathcal{L}_U V + \frac{1}{2}(b_s + b^2)L_U Y \right) - \frac{1}{2}(b_s + b^2)L_U Y. \quad (5.24)$$

Now, we have $\partial_Y L_U^{-1} (\partial_Y^2 \mathcal{L}_U V + \frac{1}{2}(b_s + b^2)L_U Y)|_{Y=0} = 0$ by construction, so that we can apply Hardy inequalities to the first term. For instance

$$\int_0^\infty \frac{1}{Y^2} \left(\partial_Y L_U^{-1} \left(\partial_Y^2 \mathcal{L}_U V + \frac{1}{2}(b_s + b^2)L_U Y \right) \right)^2 w_2 \leq H_2 \int_0^\infty (\partial_Y^2 \mathcal{L}_U^2 V)^2 w_2.$$

Using the additional bound on $\partial_Y^3 U$ from Corollary 5.4, the computations are almost identical to the ones on page 57. The only difference lies in the treatment of one non-linear term, for which we do not apply exactly the same strategy (i.e. evaluate the term with the least number of derivatives in L^∞ , and the others in L^2) and for which we use the extra information coming from the bound on E_1 and D_1 . More precisely, the only term for which we do not use the same type of estimates as the ones on page 57 is

$$\int_0^\infty \partial_Y^3 \frac{\mathcal{D}}{U} \left(\int_0^Y (\mathcal{L}_U^2 V + \frac{1}{2}(b_s + b^2)Y) \right) \partial_Y^2 \mathcal{L}_U^2 V w_2.$$

For this term, the problem comes from the part of $\partial_Y^3 \frac{\mathcal{D}}{U}$ for which we do not have L^∞ bounds, namely $O_{1/3}(U^{-1})\partial_Y^3 \mathcal{L}_U V + O_{1/3}(Y^{-1}U^{-1})\partial_Y^2 \mathcal{L}_U V$. We first integrate by parts once; the most problematic term is then

$$- \int_0^\infty \frac{\partial_Y^2 \mathcal{L}_U V}{U} \left(\int_0^Y (\mathcal{L}_U^2 V + \frac{1}{2}(b_s + b^2)Y) \right) \partial_Y^3 \mathcal{L}_U^2 V w_2.$$

Here, although the middle integral term has less derivatives than $\partial_Y^2 \mathcal{L}_U V$, we choose to evaluate it in L^2 thanks to a Hardy inequality because cancellations occur between $\mathcal{L}_U^2 V$ and $\frac{1}{2}(b_s + b^2)Y$. More precisely, the above integral is bounded by

$$\int_0^\infty \left| \frac{\partial_Y^2 \mathcal{L}_U V}{U^{1/2}} Y^3 \right| \left| \frac{1}{Y^3} \int_0^Y (\mathcal{L}_U^2 V + \frac{1}{2}(b_s + b^2)Y) \right| \frac{|\partial_Y^3 \mathcal{L}_U^2 V|}{U^{1/2}} w_2.$$

Now,

$$\begin{aligned} \frac{\partial_Y^2 \mathcal{L}_U V}{U^{1/2}} Y^3 &= \int_0^Y \partial_Y \left(\frac{\partial_Y^2 \mathcal{L}_U V}{U^{1/2}} Y^3 \right) \\ &\leq H_2 D_1^{1/2} \left(\int_0^Y (Y^4 + Y^6) w_1^{-1} \right)^{1/2} = O_{\beta_1}(s^{(7+a)\beta_1/2} D_1^{1/2}). \end{aligned}$$

The part of the integral bearing on $Y \gtrsim s^{\beta_1}$ can be bounded by noticing that we have pointwise bounds on $\int_0^Y \mathcal{L}_U^2 V$ since

$$\left| \int_0^Y \mathcal{L}_U^2 V \right| = U \left| \int_0^Y \frac{\partial_Y^2 \mathcal{L}_U V}{U^2} \right| \leq 2U \int_1^Y \frac{|\partial_Y \mathcal{L}_U V| U_Y}{U^3} + \frac{|\partial_Y \mathcal{L}_U V|}{U}$$

For the term involving $(b_s + b^2)$ on the set $Y \gtrsim s^{\beta_1}$, we merely control $\partial_Y^2 \mathcal{L}_U V$ by E_1 .

Using Hardy inequalities and recalling that $\beta_2 < \beta_1$, $m_2 \gg m_1$, we infer that the problematic integral is bounded by

$$\begin{aligned} H_2 s^{7\beta_1/2} D_1^{1/2} E_2^{1/2} D_2^{1/2} + \delta D_2 + s^{-P} D_1 + |b_s + b^2|^2 s^{-P} E_1 \\ \leq 2\delta D_2 + \frac{H_2}{\delta} s^{(7+a)\beta_1} D_1 E_2 + s^{-P} D_1 + |b_s + b^2|^2 s^{-P} E_1. \end{aligned}$$

We will then choose β_1 so that $7\beta_1 < 6 - (11 - a)\beta_1$, which is possible since $\beta_1 < 1/3$. We conclude that for all $\delta > 0$, for $s_0 \geq \max(S_0, J^4, C_0^8, \delta^{-p})$ for some $p > 0$,

$$\begin{aligned} \left| \int_0^\infty \partial_Y^2 \mathcal{C} \left[\partial_Y^2 \mathcal{L}_U V + \frac{1}{2}(b_s + b^2) L_U Y \right] \partial_Y^2 \mathcal{L}_U^2 V w_2 \right| \\ \leq \delta(bE_2 + D_2) + \frac{H_2}{\delta} s^{(7+a)\beta_1} D_1 E_2 + s^{-P} D_1 + |b_s + b^2|^2 s^{-P} E_1 + s^{-P}. \end{aligned}$$

The next term coming from (5.24) is

$$\int_0^\infty \partial_Y^2 \mathcal{C} \left[\frac{1}{2}(b_s + b^2) L_U Y \right] \partial_Y^2 \mathcal{L}_U^2 V w_2.$$

It turns out that the main order term in $\partial_Y^2 \mathcal{C}[L_U Y]$ vanishes. More precisely, following the decomposition from Lemma 4.1, we write $\mathcal{D} = -bY/2 + \tilde{\mathcal{D}} + \mathcal{D}_{NL} =: \mathcal{D}_0 + \tilde{\mathcal{D}} + \mathcal{D}_{NL}$, and we decompose the operator \mathcal{C} into $\mathcal{C}_0 + \tilde{\mathcal{C}} + \mathcal{C}_{NL}$ accordingly. Concerning \mathcal{C}_0 , an explicit computation in Appendix A shows that $\partial_Y^2 \mathcal{C}_0[L_U Y] = O_{\beta_1}(b^2 Y)$, so that, for $s \geq s_0$ sufficiently large,

$$\begin{aligned} \left| \int_0^\infty \partial_Y^2 \mathcal{C}_0 \left[\frac{1}{2}(b_s + b^2) L_U Y \right] \partial_Y^2 \mathcal{L}_U^2 V w_2 \right| \\ \leq H_2 b^2 |b_s + b^2| s^{(3-a)\beta_2/2} E_2^{1/2} \leq \delta b E_2 + b^2 (b_s + b^2)^2. \end{aligned}$$

Concerning the terms involving the operator $\tilde{\mathcal{C}}$, we use the fact that $\partial_Y^k \tilde{\mathcal{D}} = O_{\beta_1}(b^3 Y^{7-k} + b^4 Y^{10-k})$ for $Y \gg 1$ and for $k = 0, 1, 2$. Therefore

$$\partial_Y^2 \tilde{\mathcal{C}}[L_U Y] = O_{\beta_1}(b^3 Y^4) - \partial_Y^3 \frac{\tilde{\mathcal{D}}}{U} \frac{Y^2}{2}.$$

Hence, integrating by parts the term involving $\partial_Y^3 \tilde{\mathcal{D}}$ and choosing s_0 large enough,

$$\begin{aligned} \left| \int_0^\infty \partial_Y^2 \tilde{\mathcal{C}} \left[\frac{1}{2}(b_s + b^2) L_U Y \right] \partial_Y^2 \mathcal{L}_U^2 V w_2 \right| \\ \leq b^3 |b_s + b^2| \int_0^\infty O_{\beta_1}(Y^4) |\partial_Y^2 \mathcal{L}_U^2 V| w_2 + b^3 |b_s + b^2| \int_0^\infty O_{\beta_1}(Y^5) |\partial_Y^3 \mathcal{L}_U^2 V| w_2 \\ \leq \delta b E_2 + \delta D_2 + \frac{H_2}{\delta} (b_s + b^2)^2 (s^{-5+9\beta_2} + s^{-6+13\beta_2}) \\ \leq \delta b E_2 + \delta D_2 + b^2 (b_s + b^2)^2. \end{aligned}$$

There remains to address the terms involving \mathcal{D}_{NL} . Notice that $\mathcal{D}_{NL}, \partial_Y \mathcal{D}_{NL}$ are bounded in L^∞ (see Corollary 5.4). As above, we integrate by parts the term involving $\partial_Y^3 \mathcal{D}_{NL}$. Eventually, we obtain, using Corollary 5.4

$$\begin{aligned} & \left| \int_0^\infty \partial_Y^2 \mathcal{C}_{NL} \left[\frac{1}{2}(b_s + b^2)L_U Y \right] \partial_Y^2 \mathcal{L}_U^2 V w_2 \right| \\ & \leq \delta b E_2 + \delta D_2 + \frac{H_2}{\delta} b^2 |b_s + b^2|^2 + s^{-P} D_1. \end{aligned}$$

Gathering all the estimates, we obtain, if $s_0 \geq \max(S_0, J^4, C_0^8)$,

$$\left| \int_0^\infty \partial_Y^2 \mathcal{C} [\partial_Y^2 \mathcal{L}_U V] \partial_Y^2 \mathcal{L}_U^2 V w_2 \right| \leq \delta (b E_2 + D_2) + \frac{H_2}{\delta} b^2 (b_s + b^2)^2 + \frac{H_2}{\delta} s^{7\beta_1} D_1 E_2 + s^{-P} D_1.$$

We then address the second term in (5.23), for which we use the same type of decomposition as above, writing

$$\partial_Y^2 V = \left(\partial_Y^2 V + \frac{1}{48}(b_s + b^2)L_U Y^4 \right) - \frac{1}{48}(b_s + b^2)L_U Y^4 =: \partial_Y^2 \tilde{V} - \frac{1}{48}(b_s + b^2)L_U Y^4.$$

Using the formula for $L_U^{-1}(Y^2)$ (4.2) together with the bounds on V stemming from E_1 , notice that

$$\partial_Y \mathcal{L}_U^2 \tilde{V}|_{Y=0} = \partial_Y \mathcal{L}_U^2 V|_{Y=0} + \frac{1}{4} (\partial_Y L_U^{-1} Y^2)|_{Y=0} = 0,$$

and that

$$\partial_Y^2 \mathcal{L}_U^2 \tilde{V} = \partial_Y^2 \mathcal{L}_U^2 V + \frac{1}{4}(b_s + b^2) \partial_Y^2 L_U^{-1} Y^2 = \partial_Y^2 \mathcal{L}_U^2 V + O_{\beta_1}((b_s + b^2)bY).$$

Concerning the term

$$\int_0^\infty \partial_Y^2 \mathcal{L}_U \mathcal{C} [\partial_Y^2 \tilde{V}] \partial_Y^2 \mathcal{L}_U^2 V w_2, \quad (5.25)$$

the computations are very similar to the ones above, using the formula for $\partial_Y^2 \mathcal{C}[\cdot]$ from Lemma 4.12 on the one hand, and the formula for $\partial_Y^2 L_U^{-1}$ from Lemma A.1 on the other hand. We leave the details of the estimates to the reader since they do not raise any additional difficulty. We end up with

$$(5.25) \leq \delta (D_2 + b E_2) + b^2 (b_s + b^2)^2 + \frac{H_2}{\delta} \left(s^{-3} D_1 + s^{(7+a)\beta_1} D_1 E_2 + s^3 (b_s + b^2)^2 E_2 + s^{-P} \right).$$

We then consider

$$\frac{1}{48}(b_s + b^2) \int_0^\infty \partial_Y^2 \mathcal{L}_U \mathcal{C} [L_U Y^4] \partial_Y^2 \mathcal{L}_U^2 V w_2. \quad (5.26)$$

A straightforward computation leads to

$$\partial_Y^2 \mathcal{L}_U \mathcal{C} [L_U Y^4] = O_{\beta_1} \left(\frac{b}{(1+Y)U} \right) + \sum_{i=0}^4 O_{\beta_1} ((1+Y)^{i-1}) \partial_Y^i \int_0^Y \mathcal{L}_U^2 V.$$

We have

$$\left| \int_0^\infty O_{\beta_1} \left(\frac{b(b_s + b^2)}{(1+Y)U} \right) \partial_Y^2 \mathcal{L}_U^2 V w_2 \right| \leq \delta D_2 + \frac{H_2}{\delta} b^2 (b_s + b^2)^2.$$

Expressing $\partial_Y \mathcal{L}_U^2 V$ and $\mathcal{L}_U^2 V$ in terms of $\mathcal{L}_U V$ and using the expressions of E_1, D_1 , it can be easily proved that the terms of the form $O_{\beta_1}((1+Y)^{i-1})\partial_Y^i \int_0^Y \mathcal{L}_U^2 V$ give rise to integrals that can be bounded by

$$\delta b E_2 + \delta D_2 + H_2 s^{-2} D_1 + s^{-P}.$$

Gathering all the terms, we obtain the estimate announced in Lemma 5.6.

The remainder terms: proof of Lemma 5.7

We now consider the remainder terms occurring in the right-hand side of the differential inequality for E_2 , namely

$$\int_0^\infty \partial_Y^2 \mathcal{L}_U^2 V \partial_Y^2 \mathcal{L}_U^2 \mathcal{R} w_2.$$

Following the decomposition of \mathcal{R} from Remark 2.7, we will write \mathcal{R} as $\sum_{i=1}^4 \mathcal{R}_i$, and study the contribution of each \mathcal{R}_i separately. We emphasize that the most important terms are \mathcal{R}_1 and \mathcal{R}_2 : indeed, they dictate the final convergence rate, whereas the terms involving \mathcal{R}_3 and \mathcal{R}_4 are small perturbations of the main dissipation terms $D_2 + bE_2$.

• Remainder stemming from \mathcal{R}_1 :

Using the decomposition (5.15) of the previous paragraph and using the fact that for $k \geq 8$ and $Y \gg 1$,

$$\partial_Y^2 L_U^{-1} \partial_Y^2 L_U^{-1} \left[Y^k \chi \left(\frac{Y}{s^{2/7}} \right) \right] = O_{\beta_1}(Y^{k-8}) + O_{\beta_1}(Y^{k-4}) \partial_Y^2 \mathcal{L}_U V + O_{\beta_1}(Y^{k-3}) \partial_Y \mathcal{L}_U V,$$

we find that

$$\begin{aligned} & \partial_Y \mathcal{L}_U^2(\mathcal{R}_1) \\ &= (b_s + b^2) b \partial_Y^2 L_U^{-1} \partial_Y^2 L_U^{-1} \left[\left(\frac{a_4}{2} Y^5 - \frac{61}{16} a_7 b Y^8 \right) \chi \left(\frac{Y}{s^{2/7}} \right) \right] \\ &+ O_{\beta_1}(b^4) + O_{\beta_1}(b^5 Y^4) + O_{\beta_1}(b^4 Y^5 + b^5 Y^9) \partial_Y^2 \mathcal{L}_U V + O_{\beta_1}(b^4 Y^6 + b^5 Y^{10}) \partial_Y \mathcal{L}_U V \\ &- \frac{1}{2} (b_s + b^2) \partial_Y L_U^{-1} \partial_Y^2 L_U^{-1} L_V Y + \partial_Y^2 L_U^{-1} \left(\tilde{P}_1(s, Y) (1 - \chi) \left(\frac{Y}{s^{2/7}} \right) \right). \end{aligned} \quad (5.27)$$

Define

$$\varphi(s, Y) := \partial_Y^2 L_U^{-1} \partial_Y^2 L_U^{-1} \left[Y^5 \chi \left(\frac{Y}{s^{2/7}} \right) \right].$$

Then

$$\begin{aligned} \varphi(s, Y) &= \partial_Y^3 U \int_0^Y \left(\frac{1}{U^2} \partial_Y^2 L_U^{-1} \left[Y^5 \chi \left(\frac{Y}{s^{2/7}} \right) \right] \right) + \frac{U_{YY}}{U^2} \partial_Y^2 L_U^{-1} \left[Y^5 \chi \left(\frac{Y}{s^{2/7}} \right) \right] \\ &- \frac{U_Y}{U^2} \partial_Y^3 L_U^{-1} \left[Y^5 \chi \left(\frac{Y}{s^{2/7}} \right) \right] + \frac{1}{U} \partial_Y^4 L_U^{-1} \left[Y^5 \chi \left(\frac{Y}{s^{2/7}} \right) \right]. \end{aligned}$$

Since we focus on the value of the above quantities for $Y \leq s^{\beta_1}$, we can replace $\chi(Y/s^{2/7})$ by 1; following by now usual arguments, the part of the integral bearing on $Y \geq s^{\beta_1}$ will

be smaller than s^{-P} for $P > 0$ arbitrarily large, provided $m_2 \gg m_1$ is chosen large enough. Using Lemma A.1, we have, for $Y \leq s^{\beta_1}$ and for $s \geq s_0$ large enough,

$$\begin{aligned} \partial_Y^2 L_U^{-1} \left[Y^5 \chi \left(\frac{Y}{s^{2/7}} \right) \right] &= \partial_Y^3 U \int_0^Y \frac{Y^5}{U^2} \\ &\quad + \frac{Y^5 - 5(1+Y)Y^4 + 20Y^3(Y + \frac{Y^2}{2}) + O(bY^7)}{U^2} \\ &= \frac{6Y^5 + 15Y^4 + O(bY^7)}{U^2} \geq \frac{5Y^5 + 14Y^4}{U^2} > 0 \end{aligned} \quad (5.28)$$

while

$$\begin{aligned} \partial_Y^3 L_U^{-1} \left[Y^5 \chi \left(\frac{Y}{s^{2/7}} \right) \right] &= \partial_Y^4 U \int_0^Y \frac{Y^5}{U^2} + 2 \left(\frac{\partial_Y^3 U}{U^2} - \frac{U_Y U_{YY}}{U^3} \right) Y^5 \\ &\quad + 10Y^4 \frac{U_Y^2}{U^3} - 40Y^3 \frac{U_Y}{U^2} + 120 \frac{Y^2}{U} \\ &= \frac{2Y^4(9Y^2 + 29Y + 45)}{U^3} + O(bY^2) + O(Y^2 U) \partial_Y^2 \mathcal{L}_U V. \end{aligned} \quad (5.29)$$

A similar formula holds for $\partial_Y^4 L_U^{-1}[Y^4]$. It follows that

$$\varphi(s, Y) = O_{\beta_1}((1+Y)^{-3}) + O_{\beta_1}(Y) \partial_Y^2 \mathcal{L}_U V + O_{\beta_1}(Y^2) \partial_Y^3 \mathcal{L}_U V + O_{\beta_1}(bY)$$

and

$$\begin{aligned} |\partial_Y^2 \mathcal{L}_U^2 \mathcal{R}_1| &= |b_s + b^2| b |\varphi(s, Y)| + O_{\beta_1}(b^4 + b^5 Y^4) \\ &\quad + O_{\beta_1}(b^4 Y^6 + b^5 Y^{10} + (b_s + b^2) Y^2) \partial_Y \mathcal{L}_U^2 V + \text{l.o.t.} \end{aligned}$$

As a consequence, since $\beta_2 < \beta_1$,

$$\begin{aligned} &\left| \int_0^\infty [(b_s + b^2) b \varphi(s, Y)] [\partial_Y^2 \mathcal{L}_U^2 V] w_2 \right| \\ &\leq H_2 b |b_s + b^2| D_2^{1/2} \left(\int_0^\infty \frac{U^2}{(1+Y)^6} w_2 \right)^{1/2} + H_2 b^2 |b_s + b^2| E_2^{1/2} \left(\int_0^\infty Y^2 w_2 \right)^{1/2} \\ &\quad + H_2 b |b_s + b^2| E_2^{1/2} \left(s^{6/7} D_1^{1/2} + s^{-P} \right). \end{aligned}$$

For any $\delta > 0$, if $s \geq s_0$ large enough (depending on δ), the right-hand side is bounded by

$$\begin{aligned} \delta (bE_2 + D_2) + \frac{H_2}{\delta} b^2 |b_s + b^2|^2 + \frac{H_2}{\delta} b |b_s + b^2|^2 s^{12/7} D_1 + s^{-P} \\ \leq \delta (bE_2 + D_2) + \frac{H_2}{\delta} b^2 |b_s + b^2|^2 + s^{-2} D_1 + s^{-P}. \end{aligned}$$

Gathering all the terms, it follows that

$$\int_0^\infty (\partial_Y^2 \mathcal{L}_U^2 \mathcal{R}_1) (\partial_Y^2 \mathcal{L}_U^2 V) Y^{-a} w_2 \leq \delta (bE_2 + D_2) + \frac{H_2}{\delta} b^2 |b_s + b^2|^2 + s^{-2} D_1(s) + s^{-P}$$

• **Remainder stemming from \mathcal{R}_2 :** As announced in Remark 2.7, the second remainder \mathcal{R}_2 will partly dictate the total size of the remainder. More precisely, we have, for $Y \gg 1$,

$$\begin{aligned}\mathcal{L}_U^2 \mathcal{R}_2 &= O_{\beta_1}(b^4 Y^3), \\ \partial_Y \mathcal{L}_U^2 \mathcal{R}_2 &= O_{\beta_1}(b^4 Y^2) + O_{\beta_1}(b^4 Y^5) \mathcal{L}_U^2 V, \\ \partial_Y^2 \mathcal{L}_U^2 \mathcal{R}_2 &= O_{\beta_1}(b^4 Y) + O_{\beta_1}(b^4 Y^5) \partial_Y \mathcal{L}_U^2 V + O_{\beta_1}(b^4 Y^4) \mathcal{L}_U^2 V.\end{aligned}$$

We infer that for all $\delta > 0$ and for $s \geq s_0$ large enough (depending on δ),

$$\left| \int_0^\infty \partial_Y^2 \mathcal{L}_U^2 \mathcal{R}_2 \partial_Y^2 \mathcal{L}_U^2 V w_2 \right| \leq \delta b E_2 + \frac{H_2}{\delta} s^{-7+(3-a)\beta_2} + \frac{H_2}{\delta} s^{-17/4} D_1.$$

• **Remainder stemming from \mathcal{R}_3 :**

An easy computation leads to

$$\begin{aligned}\mathcal{L}_U^2(L_U^{-1} L_V Y) &= O_{\beta_1}(1) \partial_Y \mathcal{L}_U^2 V + \text{l.o.t.} + U_Y \int_0^Y \frac{\partial_Y^2 \mathcal{L}_U L_U^{-1}(L_V Y)}{U^2}, \\ \partial_Y \mathcal{L}_U^2(L_U^{-1} L_V Y) &= O_{\beta_1}(1) \partial_Y^2 \mathcal{L}_U^2 V + O_{\beta_1}\left(\frac{1}{1+Y}\right) \partial_Y \mathcal{L}_U^2 V + \text{l.o.t.} \\ &\quad + U_{YY} \int_0^Y \frac{\partial_Y^2 \mathcal{L}_U L_U^{-1}(L_V Y)}{U^2}.\end{aligned}$$

In order to estimate $\partial_Y^2 \mathcal{L}_U^2(L_U^{-1} L_V Y)$, we use the same trick as in the commutator estimate and we replace V by its asymptotic expansion close to $Y = 0$. More precisely, we write

$$V = \left(V + \frac{1}{2}(b_s + b^2) \mathcal{L}_U^{-2} Y \right) - \frac{1}{2}(b_s + b^2) \mathcal{L}_U^{-2} Y =: V_0 - \frac{1}{2}(b_s + b^2) \mathcal{L}_U^{-2} Y,$$

with the convention $\partial_Y^{-1} = \int_0^Y$. Now, by definition $\partial_Y \mathcal{L}_U^2 V_0|_{Y=0} = 0$ and $\partial_Y^2 \mathcal{L}_U^2 V_0 = \partial_Y^2 \mathcal{L}_U^2 V$. Moreover, it can be easily checked that

$$\begin{aligned}& \partial_Y^2 \mathcal{L}_U^2 L_U^{-1} \left(Y \mathcal{L}_U^{-2} Y - \frac{Y^2}{2} \partial_Y \mathcal{L}_U^{-2} Y \right) \\ &= O_{\beta_1} \left(\frac{1}{Y(1+Y)^2} \right) + O_{1/3}(Y^5) \partial_Y^2 \mathcal{L}_U^2 V + O_{1/3}(Y^4) \partial_Y \mathcal{L}_U^2 V + \text{l.o.t.},\end{aligned}$$

while

$$\partial_Y^2 \mathcal{L}_U^2 L_U^{-1}(L_{V_0} Y) = O_{\beta_1} \sum_{i=1}^3 O_{1/3}(Y^{3+i}) \partial_Y^i \mathcal{L}_U^2 V + \text{l.o.t.}$$

It follows that

$$\begin{aligned}& \left| \int_0^\infty b \partial_Y^2 \mathcal{L}_U^2(L_U^{-1} L_V Y) \partial_Y^2 \mathcal{L}_U^2 V w_2 \right| \\ & \leq \delta D_2 + \frac{C}{\delta} b^2 \int_0^\infty U O_{\beta_1} \left(\frac{Y^4}{U^2} \right) (\partial_Y^2 \mathcal{L}_U^2 V)^2 w_2 + H_2 b \int_0^\infty \frac{1}{1+Y} (\partial_Y^2 \mathcal{L}_U^2 V)^2 w_2 \\ & \quad + H_2 b |b_s + b^2| \int_0^\infty O_{\beta_1} \left(\frac{1}{Y(1+Y)^2} \right) |\partial_Y^2 \mathcal{L}_U^2 V| w_2 \\ & \quad + H_2 b |b_s + b^2| \sum_{i=1}^3 \int_0^\infty O_{1/3}(Y^{3+i}) |\partial_Y^i \mathcal{L}_U^2 V| |\partial_Y^2 \mathcal{L}_U^2 V| w_2.\end{aligned}$$

Using the estimate on $\partial_Y \mathcal{L}_U^2 V$ from Corollary 5.4, we infer that the right-hand side is bounded by

$$\delta(D_2 + bE_2) + \frac{H_2}{\delta} b^2 (b_s + b^2)^2 + H_2 s^{-3} D_1.$$

The same method and estimates apply to $b^3 L_U^{-1}(L_V Y^7)$. At last, we consider

$$b^3 L_U^{-1} \left(\chi \left(\frac{Y}{s^{2/7}} \right) L_{-a_4 b Y^4 - a_7 b^2 Y^7 + a_{10} b^3 Y^{10} + a_{11} b^3 Y^{11} Y^7} \right) =: L_U^{-1}(\zeta(s, Y)).$$

Note that $\zeta(s, \cdot) \in \mathcal{C}^\infty(Y)$ and that

$$\partial_Y^k \zeta(s, Y) = O(b^4 Y^{11-k}) \quad \forall k \leq 11.$$

It follows that

$$\begin{aligned} \mathcal{L}_U^2 L_U^{-1} \zeta &= O(b^4 Y \ln Y) + O_{1/3}(s^{-10/3} Y^4 + s^{-4} Y^8) \partial_Y \mathcal{L}_U^2 V + \text{l.o.t.} \\ \partial_Y \mathcal{L}_U^2 L_U^{-1} \zeta &= O(b^4 \ln Y) + O_{1/3}(s^{-10/3} Y^4 + s^{-4} Y^8) \partial_Y^2 \mathcal{L}_U^2 V \\ &\quad + O_{1/3}(s^{-10/3} Y^3 + s^{-4} Y^7) \partial_Y \mathcal{L}_U^2 V + \text{l.o.t.} \\ \partial_{YY} \mathcal{L}_U^2 L_U^{-1} \zeta &= O\left(\frac{b^4}{1+Y}\right) + O_{1/3}(s^{-10/3} Y^2 + s^{-4} Y^6) \partial_Y^3 \mathcal{L}_U^2 V \\ &\quad + O_{1/3}(s^{-10/3} Y^3 + s^{-4} Y^7) \partial_Y \mathcal{L}_U^2 V + \text{l.o.t.} \end{aligned}$$

We obtain, since $\beta_2 < \beta_1 < 1/3$,

$$\begin{aligned} \left| \int_0^\infty (\mathcal{L}_U^2 L_U^{-1} \zeta \mathcal{L}_U^2 V \partial_{YY} w_2 + \partial_Y \mathcal{L}_U^2 L_U^{-1} \zeta \partial_Y \mathcal{L}_U^2 V w_2) \right| \\ \leq \delta b E_2 + \frac{H_2}{\delta} s^{-27/4} (\ln s)^2 + \frac{H_2}{\delta} s^{-7+(16+a)\beta_2} E_2 \end{aligned}$$

and, writing $\partial_Y \mathcal{L}_U^2 V = (\partial_Y \mathcal{L}_U^2 V + \frac{1}{2}(b_s + b^2)) - \frac{1}{2}(b_s + b^2)$,

$$\left| \int_0^\infty \partial_Y^2 \mathcal{L}_U^2 L_U^{-1} \zeta \partial_Y^2 \mathcal{L}_U^2 V w_2 \right| \leq \delta b E_2 + \delta D_2 + \frac{H_2}{\delta} s^{-7} + \frac{H_2}{\delta} s^{-7+(13-a)\beta_2} (b_s + b^2)^2 + s^{-M} D_1.$$

• **Remainder stemming from \mathcal{R}_4 :**

We recall that

$$\mathcal{R}_4 = P_1(s, Y)(1 - \chi_1) \left(\frac{Y}{s^{2/7}} \right) + L_U^{-1} \left(P_2(s, Y)(1 - \chi_2) \left(\frac{Y}{s^{2/7}} \right) \right),$$

and that for any $P > 0$, $k \geq 0$, $i = 1, 2$, choosing $\beta_1 < 2/7$,

$$\partial_Y^k (P_i(1 - \chi_i) \left(\frac{Y}{s^{2/7}} \right)) = O_{\beta_1}(s^{-P}).$$

Using the same type of computations as above, it follows that for any $M > 0$,

$$\begin{aligned} \mathcal{L}_U^2 \mathcal{R}_4 &= O_{\beta_1}(s^{-P}) + O_{\beta_1}(s^{-P}) \partial_Y \mathcal{L}_U^2 V + \text{l.o.t. in } L^2(\partial_{YY} w_2), \\ \partial_Y \mathcal{L}_U^2 L_U^{-1} \psi &= O_{\beta_1}(s^{-P}) + O_{\beta_1}(s^{-P}) \partial_Y^2 \mathcal{L}_U^2 V + O_{\beta_1}(s^{-P}) \partial_Y \mathcal{L}_U^2 V + \text{l.o.t. in } L^2(w_2) \\ \partial_{YY} \mathcal{L}_U^2 L_U^{-1} \psi &= O_{\beta_1}(s^{-P}) + O_{\beta_1}(s^{-P}) \partial_Y^3 \mathcal{L}_U^2 V + O_{\beta_1}(s^{-P}) \partial_Y \mathcal{L}_U^2 V + \text{l.o.t. in } L^2(w_2). \end{aligned}$$

Thus, choosing m_2 and m_2 large enough,

$$\left| \int_0^\infty (\mathcal{L}_U^2 \mathcal{R}_4 \mathcal{L}_U^2 V \partial_{YY} w_2 + \partial_Y \mathcal{L}_U^2 \mathcal{R}_4 \partial_Y \mathcal{L}_U^2 V w_2) \right| \leq \delta b E_2 + s^{-P} + s^{-P} E_2 + s^{-P} D_1,$$

$$\left| \int_0^\infty \partial_Y^2 \mathcal{L}_U^2 \mathcal{R}_4 \partial_Y^2 \mathcal{L}_U^2 V w_2 \right| \leq \delta b E_2 + \delta D_2 + s^{-P} + s^{-P} D_1.$$

Gathering all the terms, we obtain the estimates announced in Lemma 5.7.

Acknowledgements

A.-L. Dalibard is partially supported by the Agence Nationale de la Recherche, project Dyficolti, Grant ANR-13-BS01-0003-01. This project has received funding from the European Research Council (ERC) under the European Union's Horizon 2020 research and innovation program Grant agreement No 637653, project BLOC "Mathematical Study of Boundary Layers in Oceanic Motion". This work was initiated during A.-L. D.'s stay at the Courant Institute in 2013-2014, and she warmly thanks the Courant Institute for its hospitality. N. M is partially supported by the NSF grant DMS-1716466.

A Useful formulas

- We will often need to transform derivatives of V into quantities involving $\mathcal{L}_U V, \mathcal{L}_U^2 V$ and their derivatives. In order to do so, we start from the following observation

$$\partial_Y^2 V = L_U \mathcal{L}_U V = U \mathcal{L}_U V - U_Y \int_0^Y \mathcal{L}_U V, \quad (\text{A.1})$$

from which it follows, applying the same idea to $\partial_Y^2 \mathcal{L}_U V$,

$$\partial_Y^3 V = U \partial_Y \mathcal{L}_U V - U_{YY} \int_0^Y \mathcal{L}_U V,$$

$$\partial_Y^4 V = U^2 \mathcal{L}_U^2 V - U U_Y \int_0^Y \mathcal{L}_U^2 V + U_Y \partial_Y \mathcal{L}_U V - U_{YY} \mathcal{L}_U V - (\partial_Y^3 U^{\text{app}} + \partial_Y^3 V) \int_0^Y \mathcal{L}_U V. \quad (\text{A.2})$$

Notice that for derivatives of U higher than or equal to two, we decompose $\partial_Y^k U$ into $\partial_Y^k U^{\text{app}} + \partial_Y^k V$. This is related to the fact that we have pointwise estimates on $\partial_Y^2 U$, but not on higher derivatives. Now, in the formula giving $\partial_Y^4 V$, we can write $\partial_Y^3 V$ in terms of $\mathcal{L}_U V$. Obviously, we can iterate this procedure. As a consequence, for any $k \geq 2$, we can express $\partial_Y^k V$ in terms of $\mathcal{L}_U^l V$ for $l \leq k/2$.

- We will also need the explicit expression of $\partial_Y^k L_U^{-1}$, which is given in the following Lemma, whose proof is straightforward and left to the reader.

Lemma A.1. *For any function W which vanishes at a sufficiently high order near $Y = 0$,*

we have

$$\begin{aligned}\partial_Y L_U^{-1} W &= U_{YY} \int_0^Y \frac{W}{U^2} + \frac{\partial_Y W}{U}, \\ \partial_Y^2 L_U^{-1} W &= \partial_Y^3 U \int_0^Y \frac{W}{U^2} + \partial_Y^2 U \frac{W}{U^2} - \partial_Y U \frac{\partial_Y W}{U^2} + \frac{\partial_Y^2 W}{U} \\ &= \partial_Y^3 U \int_0^Y \frac{W}{U^2} + \partial_Y^2 U \frac{W}{U^2} + \partial_Y \frac{\partial_Y W}{U}\end{aligned}$$

and

$$\partial_Y^3 L_U^{-1} W = \partial_Y^4 U \int_0^Y \frac{W}{U^2} + 2 \left(\frac{U \partial_Y^3 U - U_{YY} U_Y}{U^3} \right) W + 2 \frac{U_Y^2}{U^3} \partial_Y W - 2 \frac{U_Y}{U^2} \partial_{YY} W + \frac{\partial_Y^3 W}{U}.$$

- Eventually, setting $\mathcal{D}_0 := -bY/2$ and

$$\mathcal{C}_0[W] := 2L_U^{-1} \left(\mathcal{D}_0 \frac{W}{U} \right) - \partial_Y \left(\frac{\mathcal{D}_0}{U} \int_0^Y L_U^{-1} W \right),$$

we need to compute $\partial_Y^2 \mathcal{C}_0[L_U Y]$. By definition, $L_U Y = YU - U_Y Y^2/2$, so that

$$\mathcal{C}_0[L_U Y] = -\frac{b}{2} \left\{ 2L_U^{-1}(Y^2) - L_U^{-1} \left(\frac{Y^3 U_Y}{U} \right) - \partial_Y \left(\frac{Y^3}{2U} \right) \right\}.$$

Now, integrating by parts,

$$L_U^{-1} \left(\frac{Y^3 U_Y}{U} \right) = \partial_Y \left[U \int_0^Y \frac{Y^3 U_Y}{U^2} \right] = -\partial_Y \left(\frac{Y^3}{2U} \right) + \frac{3}{2} L_U^{-1}(Y^2).$$

Therefore

$$\mathcal{C}_0[L_U Y] = -\frac{b}{4} L_U^{-1}(Y^2).$$

Using the formula above for $\partial_Y^2 L_U^{-1}$, we obtain

$$\partial_Y^2 \mathcal{C}_0[L_U Y] = -\frac{b}{4} \left\{ \partial_Y^3 U \int_0^Y \frac{Y^2}{U^2} + \frac{Y^2 U_{YY} - 2Y U_Y + 2U}{U^2} \right\}.$$

B Estimate on the modulation rate

Lemma B.1. *Let $\gamma \in (0, 5)$, and let $\varphi : [s_0, s_1] \rightarrow \mathbb{R}_+$ be such that*

$$\int_{s_0}^{s_1} s^\gamma \varphi(s) ds < +\infty.$$

Assume that there exists a constant $\epsilon > 0$ such that for all $s \in [s_0, s_1]$

$$\begin{aligned}|b_s + b^2| &\leq \sqrt{\varphi}, \\ \frac{1-\epsilon}{s} &\leq b(s) \leq \frac{1+\epsilon}{s}.\end{aligned}$$

Then for all $s \in [s_0, s_1]$,

$$\left| b(s) - \frac{1}{s} \right| \leq \frac{1+\epsilon}{1-\epsilon} \left| \frac{1}{s_0} - b(s_0) \right| \frac{s_0^2}{s^2} + \frac{1+\epsilon}{(1-\epsilon)^2 \sqrt{5-\gamma}} s^{\frac{1-\gamma}{2}} \left(\int_{s_0}^{s_1} t^\gamma \varphi(t) dt \right)^{1/2}.$$

In particular:

- If $\varphi(s) = Js^{-4-\eta}$, the inequality becomes, with $\gamma = 3 + \eta/2$,

$$\left| b(s) - \frac{1}{s} \right| \leq \frac{1+\epsilon}{1-\epsilon} \left| \frac{1}{s_0} - b(s_0) \right| \frac{s_0^2}{s^2} + \frac{J^{1/2}}{\sqrt{\eta}} \frac{1+\epsilon}{(1-\epsilon)^2} s^{-1-\frac{\eta}{2}}.$$

- More generally, if $\gamma > 3$, then there exists a constant $\eta > 0$ ($\eta = (\gamma - 3)/2$) such that for all $s \in [s_0, s_1]$,

$$\left| b(s) - \frac{1}{s} \right| \leq \frac{1+\epsilon}{1-\epsilon} \left| \frac{1}{s_0} - b(s_0) \right| \frac{s_0^2}{s^2} + \frac{(1+\epsilon)J^{1/2}}{(1-\epsilon)^2 \sqrt{2-2\eta}} s^{-1-\eta},$$

where $J = \int_{s_0}^{s_1} t^\gamma \varphi(t) dt$.

Proof. The assumption on b entails

$$\left| \frac{b_s}{b^2} + 1 \right| = \left| \frac{d}{ds} \left(s - \frac{1}{b} \right) \right| \leq \frac{\sqrt{\varphi}}{b^2} \leq \frac{1}{(1-\epsilon)^2} s^2 \sqrt{\varphi}.$$

Integrating the above inequality between s_0 and s and using a Cauchy-Schwarz inequality yields

$$\begin{aligned} \left| s - \frac{1}{b(s)} \right| &\leq \left| s_0 - \frac{1}{b(s_0)} \right| + \frac{1}{(1-\epsilon)^2} \left(\int_{s_0}^{s_1} t^\gamma \varphi(t) dt \right)^{1/2} \left(\int_{s_0}^s t^{4-\gamma} dt \right)^{1/2} \\ &\leq \left| s_0 - \frac{1}{b(s_0)} \right| + \frac{1}{(1-\epsilon)^2} \left(\int_{s_0}^{s_1} t^\gamma \varphi(t) dt \right)^{1/2} \frac{s^{\frac{5-\gamma}{2}}}{\sqrt{5-\gamma}}. \end{aligned}$$

Now, multiplying the above inequality by $b/s \leq (1+\epsilon)s^{-2}$, we obtain

$$\left| b(s) - \frac{1}{s} \right| \leq (1+\epsilon) \frac{s_0}{b(s_0)} \left| \frac{1}{s_0} - b(s_0) \right| \frac{1}{s^2} + \frac{1+\epsilon}{(1-\epsilon)^2} \left(\int_{s_0}^{s_1} t^\gamma \varphi(t) dt \right)^{1/2} \frac{s^{\frac{1-\gamma}{2}}}{\sqrt{5-\gamma}}.$$

Since $s_0/b(s_0) \leq (1-\epsilon)^{-1}s_0^2$, we obtain the inequality announced in the Lemma. \square

Lemma B.1 has in particular the following consequence:

Corollary B.2. *Assume that b satisfies the assumptions of Lemma B.1 for some $\gamma \in]3, 4[$. For $s \geq s_0$, define \tilde{b} by*

$$\tilde{b}_s + b\tilde{b} = 0, \quad \tilde{b}|_{s=s_0} = \frac{1}{s_0}.$$

Then there exists a universal constant \bar{C} such that if $s_0 \geq \bar{C}(J\epsilon^{-2})^{1/(\gamma-3)}$, for all $s \geq s_0$,

$$\frac{1-2\epsilon}{s} \leq \tilde{b}(s) \leq \frac{1+2\epsilon}{s}.$$

Proof. Since \tilde{b} satisfies a linear ODE, we have simply

$$\tilde{b}(s) = \frac{1}{s_0} \exp \left(- \int_{s_0}^s b(s') ds' \right).$$

According to Lemma B.1, for all $s \geq s_0$,

$$\ln \frac{s}{s_0} - C_{\epsilon, \gamma, s_0} \leq \int_{s_0}^s b(s') ds' \leq \ln \frac{s}{s_0} + C_{\epsilon, \gamma, s_0},$$

where

$$C_{\epsilon, \gamma, s_0} = \frac{1 + \epsilon}{(1 - \epsilon)^2} \epsilon + \frac{1 + \epsilon}{(1 - \epsilon)^2} J^{1/2} (5 - \gamma)^{-1/2} \frac{2s_0^{\frac{3-\gamma}{2}}}{\gamma - 3}.$$

Therefore

$$\frac{e^{-C_{\epsilon, \gamma, s_0}}}{s} \leq \tilde{b}(s) \leq \frac{e^{C_{\epsilon, \gamma, s_0}}}{s}.$$

Now, if $s_0 \geq \bar{C}(J\epsilon^{-2})^{1/(\gamma-3)}$, we have

$$e^{C_{\epsilon, \gamma, s_0}} \leq 1 + 2\epsilon, \quad e^{-C_{\epsilon, \gamma, s_0}} \geq 1 - 2\epsilon,$$

which completes the proof. \square

C Proof of Lemma 3.6

As much as possible, we treat sub-solutions and super-solutions simultaneously. For any $A_{\pm} > 0$, $k_{\pm} > 2$, we consider the function

$$W_{\pm}(s, \psi) := \frac{(6\psi)^{4/3}}{4} \pm A_{\pm} \psi^{k_{\pm}} \tilde{b}^{\frac{3k_{\pm}-2}{4}}.$$

We will choose $k_- = 7/3$ for sub-solutions and $k_+ = 10/3$ for super solutions.

We claim that W_{\pm} satisfy the following properties:

- Choosing $k_- = 7/3$, there exists \bar{C} such that if $C_- \geq \bar{C}$, then for all $A_- > 0$,

$$\partial_s W_- - 2bW_- + \frac{3b}{2} \psi \partial_{\psi} W_- - \sqrt{W_-} \partial_{\psi} W_- + 2 \leq 0 \quad (\text{C.1})$$

on the domain $\{\psi \geq C_- \tilde{b}^{-3/4}\} \cap \{W_- > 0\}$.

Similarly, choosing $k_+ = 10/3$, there exists $\bar{C} > 0$ such that if $C_- \geq \bar{C}$, for all $A_+ > 0$

$$\partial_s W_+ - 2bW_+ + \frac{3b}{2} \psi \partial_{\psi} W_+ - \sqrt{W_+} \partial_{\psi} W_+ + 2 \geq 0 \quad \forall \psi \geq C_- \tilde{b}^{-3/4}. \quad (\text{C.2})$$

- There exists a constants \bar{A} such that if $A_{\pm} \geq \bar{A}$ and if s_0 is large enough (depending on A_{\pm}, C_1, a and C_-),

$$W_-(s, C_- \tilde{b}^{-3/4}) \leq W(s, C_- \tilde{b}^{-3/4}) \leq W_+(s, C_- \tilde{b}^{-3/4}). \quad (\text{C.3})$$

- There exists a constant A_0 , depending on M_0 such that if $A_{\pm} \geq A_0$,

$$W_-(s_0, \psi) \leq W(s_0, \psi) \leq W_+(s_0, \psi) \quad \forall \psi \geq C_- s_0^{3/4}. \quad (\text{C.4})$$

Proof of inequalities (C.1) and (C.2)

We first compute the transport term. We have

$$\partial_s W_{\pm} - 2bW_{\pm} + \frac{3b}{2}\psi\partial_{\psi}W_{\pm} = \pm A_{\pm} \frac{3(k_{\pm} - 2)}{4} \tilde{b}^{\frac{3k_{\pm}-2}{4}} \psi^{k_{\pm}} \geq 0$$

provided $k_{\pm} > 2$. We now address the computation of the diffusion term, which we treat a bit differently for the sub- and for the supersolution. The heuristic is that if $\psi \geq Cs^{3/4}$ for some C large enough (i.e. $Y > C's^{1/4}$ for some large C'), transport dominates the diffusion term. We prove this fact by distinguishing between two different zones:

- When $\psi^{4/3} \gg \psi^{k_{\pm}} \tilde{b}^{\frac{3k_{\pm}-2}{4}}$, i.e. $\psi \ll \tilde{b}^{-\frac{3}{4} \frac{3k_{\pm}-2}{3k_{\pm}-4}}$, we can perform asymptotic expansions of $\sqrt{W_{\pm}}$ and $\partial_{\psi\psi}W_{\pm}$. We have

$$\begin{aligned} \sqrt{W_{\pm}} &= \frac{(6\psi)^{2/3}}{2} \left(1 \pm \frac{2A_{\pm}}{6^{4/3}} \psi^{k_{\pm}-\frac{4}{3}} \tilde{b}^{\frac{3k_{\pm}-2}{4}} + O(\psi^{2k_{\pm}-\frac{8}{3}} \tilde{b}^{\frac{3k_{\pm}-2}{2}}) \right), \\ \partial_{\psi\psi}W_{\pm} &= \frac{6^{4/3}\psi^{-2/3}}{9} \left(1 \pm \frac{9A_{\pm}k_{\pm}(k_{\pm}-1)}{6^{4/3}} \psi^{k_{\pm}-4/3} \tilde{b}^{\frac{3k_{\pm}-2}{4}} \right), \end{aligned}$$

so that

$$-\sqrt{W_{\pm}}\partial_{\psi\psi}W_{\pm} + 2 = \mp \frac{2A_{\pm}}{6^{4/3}} (9k_{\pm}^2 - 9k_{\pm} + 2) \psi^{k_{\pm}-\frac{4}{3}} \tilde{b}^{\frac{3k_{\pm}-2}{4}} + O\left(\psi^{2k_{\pm}-\frac{8}{3}} \tilde{b}^{\frac{3k_{\pm}-2}{2}}\right).$$

Therefore, if

$$\frac{3(k_{\pm} - 2)}{4} b > \frac{4}{6^{4/3}} (9k_{\pm}^2 - 9k_{\pm} + 2) \psi^{-4/3} \quad \forall \psi \geq C_- \tilde{b}^{-4/3},$$

then

$$\partial_s W_{\pm} - 2bW_{\pm} + \frac{3b}{2}\psi\partial_{\psi}W_{\pm} - \sqrt{W_{\pm}}\partial_{\psi\psi}W_{\pm} + 2 \geq 0.$$

Hence we define

$$C_{k_{\pm}} := 2 \left(\frac{16(9k_{\pm}^2 - 9k_{\pm} + 2)}{3 \times 6^{4/3}(k_{\pm} - 2)} \right)^{3/4},$$

and recalling Lemma 3.5, we obtain inequality (C.1) on $C_{k_{\pm}} \tilde{b}^{-3/4} \leq \psi \ll \tilde{b}^{-\frac{3}{4} \frac{3k_{\pm}-2}{3k_{\pm}-4}}$ provided ϵ is sufficiently small.

- We now consider larger values of ψ . This is where we treat separately the sub- and the supersolution. Concerning the subsolution, we can use the weaker estimate

$$\sqrt{W_-} \leq \frac{(6\psi)^{2/3}}{2}, \tag{C.5}$$

which leads to, taking $k_- = 7/3$,

$$\begin{aligned} \partial_s W_- - 2bW_- + \frac{3b}{2}\psi\partial_{\psi}W_- - \sqrt{W_-}\partial_{\psi\psi}W_- + 2 \\ \leq -A_- \frac{1}{4} b \tilde{b}^{5/4} \psi^{7/3} + \frac{6^{2/3}}{2} A_- \frac{28}{9} \psi \tilde{b}^{5/4} + 2. \end{aligned}$$

The right-hand side above is negative if ϵ is small enough and

$$\psi^{7/3} > \frac{16}{A_-} \tilde{b}^{-9/4}.$$

Noticing that $27/28 < 5/4$, we infer that if s_0 is large enough, $\tilde{b}^{-\frac{3k_-+2}{4}} \ll \tilde{b}^{-\frac{3}{4} \frac{3k_- - 2}{3k_- - 4}}$, and thus (C.1) is proved on the set $[C_- \tilde{b}^{-3/4}, \infty) \cap \{W_- > 0\}$ if s_0 is sufficiently large.

We now consider the supersolution, choosing $k_+ := 10/3$. We replace estimate (C.5) by

$$\sqrt{W_+} \leq \left(\frac{(6\psi)^{2/3}}{2} + \sqrt{A_+} \psi^{5/3} \tilde{b} \right),$$

so that

$$\begin{aligned} & \partial_s W_+ - 2bW_+ + \frac{3b}{2} \psi \partial_\psi W_+ - \sqrt{W_+} \partial_{\psi\psi} W_+ + 2 \\ & \geq A_+ b \tilde{b}^2 \psi^{10/3} - \bar{C} \left(1 + A_+^{3/2} \psi^3 \tilde{b}^3 \right), \end{aligned}$$

for some universal (and computable) constant \bar{C} . It is easily checked that the right-hand side is positive as soon as $\psi \gg s^{9/10}$. Furthermore, since $k_+ = 10/3$, $\frac{3}{4} \frac{3k_+ - 2}{3k_+ - 4} = 1$, and therefore $s^{9/10} \ll \tilde{b}^{-\frac{3}{4} \frac{3k_+ - 2}{3k_+ - 4}}$. Thus inequality (C.2) is proved for $\psi \geq C_{10/3} \tilde{b}^{-3/4}$ provided s_0 is sufficiently large.

Proof of inequality (C.3)

Inequality (C.3) is an immediate consequence of the asymptotic expansion (3.6). Indeed, we need to choose A_\pm such that

$$-A_- C_-^{7/3} \tilde{b}^{-1/2} \leq \tilde{b}^{-1/2} \left(\frac{(6C_-)^{2/3}}{2} - \frac{3}{5} a_4 \frac{b}{\tilde{b}} (6C_-)^2 \right) + O(s^{\frac{a+2}{8}}) \leq A_+ C_-^{10/3} \tilde{b}^{-1/2}.$$

It is clear that once C_- is fixed, we can pick A_\pm sufficiently large (depending only on k_\pm and C_-) so that the above inequality is satisfied provided ϵ is small (e.g. $\epsilon < 1/2$, recalling Lemma 3.5) and s_0 is sufficiently large, so that the $O(s^{\frac{a+2}{8}})$ term can be neglected.

Proof of inequality (C.4)

Since $-M_0 \inf(s_0^{-1} Y^2, 1) \leq U_{YY}(s_0) - 1 \leq 0$, we infer that

$$Y + \frac{Y^2}{2} - \frac{M_0}{12} s_0^{-1} Y^4 \leq U(s_0, Y) \leq Y + \frac{Y^2}{2} \quad \forall Y \geq 0.$$

Then for $1 \ll \psi \ll s_0^{3/2}$, performing the same computations as the ones leading to (3.6),

$$W(s_0, \psi) = \frac{(6\psi)^{4/3}}{4} \left(1 + 2(6\psi)^{-2/3} + O(s_0^{-1} \psi^{2/3} + \psi^{-1}) \right)$$

and therefore there exists a constant M , depending only on M_0 , such that

$$\frac{(6\psi)^{4/3}}{4} + \frac{(6\psi)^{2/3}}{2} - M s_0^{-1} \psi^2 - M \psi^{1/3} \leq W(s_0, \psi) \leq \frac{(6\psi)^{4/3}}{4} + \frac{(6\psi)^{2/3}}{2} + M s_0^{-1} \psi^2 + M \psi^{1/3}.$$

The estimate follows on the set $C_-s_0^{3/4} \leq \psi \ll s_0^{3/2}$, provided A_-, A_+ are large enough (depending on M_0). Furthermore, recall that $W_-(s_0, Cs_0^{\frac{3}{4}\frac{3k-2}{3k-4}}) = 0$ and $s_0^{\frac{3}{4}\frac{3k-2}{3k-4}} \ll s_0^{3/2}$ since $k > 2$ (if s_0 is large). Thus the inequality $W_-(s_0, \psi) \leq W(s_0, \psi)$ is valid on the domain of definition of ψ . On the other hand, for $\psi \geq cs_0^{6/5}$, $W_+(s_0, \psi) \geq \frac{A_+}{2}c^{10/3}s_0^2$. Therefore, since $W(s_0, \psi) \leq \lim_{Y \rightarrow \infty} U(s_0, Y)^2 \lesssim s_0^2$, we infer that $W(s_0, \psi) \leq W_+(s_0, \psi)$ for $\psi \geq cs_0^{6/5}$. Since $s_0^{6/5} \ll s_0^{3/2}$, we infer that $W(s_0, \psi) \leq W_+(s_0, \psi)$ on the domain of definition of W_+ .

Conclusion

Putting together inequalities (C.1), (C.2), (C.3) and (2.22) and applying the maximum principle on the domain $\{s \in [s_0, s_1], \psi \geq C_- \tilde{b}^{-3/4}\}$, we deduce that $W_-(s, \psi) \leq W(s, \psi) \leq W_+(s, \psi)$ within this parabolic domain.

D Proof of Lemma 3.7

As in paragraph 3.2, the real issue is to control $U_{YY} - 1$ in the zone $Y \gtrsim s^{1/4}$ (or equivalently, $\psi \gtrsim s^{3/4}$). To that end, we rely on the equation in von Mises variables, and we use the computations in the proof of Lemma 3.2. We set

$$F(s, \psi) := \sqrt{W} \partial_{\psi\psi} W - 2$$

and we recall that F satisfies (3.3). We now construct a function \underline{F} such that $|-M_2 b Y(s, \psi)^2 \leq \underline{F}(s, \psi) \leq 0$ for some constant M_2 and for $\psi = O(s)$, and such that

$$\begin{aligned} \partial_s \underline{F} - \frac{1}{2W} \underline{F}(\underline{F} + 2) + \frac{3b}{2} \psi \partial_{\psi} \underline{F} - \sqrt{W} \partial_{\psi\psi} \underline{F} &\leq 0 \text{ in } s > s_0, \psi > C \tilde{b}^{-3/4}, \\ \underline{F}(s, \psi) &\leq F(s_0, \psi) \text{ on } \{s_0\} \times (Cs_0^{3/4}, \infty) \cup \{\psi = C \tilde{b}^{-3/4}, s > s_0\} \cup \{\psi = \infty, s > s_0\}. \end{aligned}$$

Let us postpone for a moment the construction of \underline{F} and explain why the estimate of the Lemma follows. First, notice that $(\underline{F} - F)$ satisfies

$$\partial_s (\underline{F} - F) - \frac{1}{2W} (\underline{F} - F)(\underline{F} + F + 2) + \frac{3b}{2} \psi \partial_{\psi} (\underline{F} - F) - \sqrt{W} \partial_{\psi\psi} (\underline{F} - F) \leq 0 \text{ in } s > s_0, \psi > C \tilde{b}^{-3/4},$$

and $(\underline{F} - F)_+ = 0$ on the parabolic boundary of the domain $\{s \geq s_0, \psi \geq C \tilde{b}^{-3/4}\}$. We then multiply the above inequality by $(\underline{F} - F)_+$, integrate in ψ over $[C \tilde{b}^{-3/4}, +\infty)$ and use the same argument as in Lemma 3.2. It follows that $(\underline{F} - F)_+ \equiv 0$, and thus $F \geq \underline{F}$ for all $s \geq s_0, \psi \geq C \tilde{b}^{-3/4}$. In particular, for $Y \leq cs^{1/3}$,

$$U_{YY}(s, Y) - 1 \geq -\frac{M_2}{2} b Y^2,$$

and the estimate announced in the statement of the Lemma follows.

We now turn towards the construction of \underline{F} . According to Lemma 3.6, there exists $A_- > 0$ and $C_- > 0$ such that if $\psi \geq C_- \tilde{b}^{-3/4}$,

$$W(s, \psi) \geq \frac{(6\psi)^{4/3}}{4} - A_- \psi^{7/3} \tilde{b}^{5/4}$$

Let us construct \underline{F} by treating separately the intervals $(C_- \tilde{b}^{-3/4}, c\tilde{b}^{-5/4})$ and $(c\tilde{b}^{-5/4}, +\infty)$, for some small constant $c > 0$ to be determined.

- For $\psi \in (C_- \tilde{b}^{-3/4}, c\tilde{b}^{-5/4})$ we take $\underline{F}(s, \psi) = -\tilde{b}\alpha \underbrace{(\psi^{2/3} - \psi^{1/3})}_{=:g(\psi)}$, for some large constant α to be determined. Then

$$\begin{aligned} & \partial_s \underline{F} - \frac{1}{2W} \underline{F}(\underline{F} + 2) + \frac{3b}{2} \psi \partial_\psi \underline{F} - \sqrt{W} \partial_{\psi\psi} \underline{F} \\ &= -b\tilde{b} \frac{\alpha}{2} \psi^{1/3} - \frac{1}{2W} \underline{F}^2 + \tilde{b}\alpha \sqrt{W} \left[\frac{g}{W^{3/2}} + g'' \right]. \end{aligned}$$

Let us evaluate the term in brackets in the right-hand side. On the set $(C_- \tilde{b}^{-3/4}, c\tilde{b}^{-5/4})$, we have

$$\frac{1}{W^{3/2}} \leq \left(\frac{(6\psi)^{4/3}}{4} - A_- \psi^{7/3} \tilde{b}^{5/4} \right)^{-3/2} = \frac{2}{9\psi^2} \left(1 + \frac{A_-}{6^{1/3}} \psi \tilde{b}^{5/4} + O(\psi^2 s^{-5/2}) \right).$$

Therefore

$$\begin{aligned} & \frac{g}{W^{3/2}} + g'' \\ & \leq \frac{2}{9\psi^2} \left(1 + \frac{A_-}{6^{1/3}} \psi \tilde{b}^{5/4} + O(\psi^2 s^{-5/2}) \right) (\psi^{2/3} - \psi^{1/3}) - \frac{2}{9} \psi^{-4/3} + \frac{2}{9} \psi^{-5/3} \\ & \leq \frac{2A_-}{9 \cdot 6^{1/3}} \psi^{-1/3} \tilde{b}^{5/4} + O(\psi^{2/3} \tilde{b}^{5/2} + \psi^{-2/3} \tilde{b}^{5/4}). \end{aligned}$$

Using Lemma 3.6, we see that $W = O(\psi^{4/3})$ for $\psi \leq c\tilde{b}^{-1}$. Therefore, for any $\alpha > 0$, provided c is small enough and s_0 is large enough, we have

$$\left| \tilde{b}\alpha \sqrt{W} \left[\frac{g}{W^{3/2}} + g'' \right] \right| \leq b\tilde{b} \frac{\alpha}{4} \psi^{1/3},$$

whence

$$\partial_s \underline{F} - \frac{1}{2W} \underline{F}(\underline{F} + 2) + \frac{3b}{2} \psi \partial_\psi \underline{F} - \sqrt{W} \partial_{\psi\psi} \underline{F} \leq 0 \text{ on } s \in (s_0, s_1), \psi \in (C_- \tilde{b}^{-3/4}, c\tilde{b}^{-1}).$$

We also need to check that $\underline{F} \leq F$ on $\{s = s_0, \psi \in (C_- s_0^{3/4}, c s_0^{5/4})\} \cup \{\psi = C_- \tilde{b}^{-3/4}\}$. According to Lemma 3.4, we have, for $\psi = C_- \tilde{b}^{-3/4}$,

$$F(s, \psi) = -\frac{6^{2/3}}{2} b \psi^{2/3} + O(s^{\frac{\alpha-6}{8}}).$$

Therefore it is sufficient to take $\alpha \geq 6^{2/3}$ and s_0 large enough. On the other hand, on the set $\{s = s_0, \psi \in (C_- s_0^{3/4}, c s_0^{5/4})\}$, since we know that

$$Y + \frac{Y^2}{2} - M_0 s_0^{-1} \frac{Y^4}{12} \leq U(s_0, Y) \leq Y + \frac{Y^2}{2} \quad \forall Y \geq 0,$$

we have $Y = (6\psi)^{1/3} + O(1)$, and therefore assumption (3.9) implies

$$F(s_0, \psi) \geq -2M_0 6^{2/3} s_0^{-1} \psi^{2/3}.$$

Therefore, choosing $\alpha = \max(6^{2/3}, 12M_0)$, we have $\underline{F} \leq F$ on $\{s = s_0, \psi \in (C_-s_0^{3/4}, cs_0^{5/4})\} \cup \{\psi = C_- \tilde{b}^{-3/4}\}$. Note that since $Y \sim (6\psi)^{1/3}$ for $1 \ll \psi \lesssim s^{1/3}$, this choice of α amounts to taking $M_2 = \bar{M} \max(1, M_0)$.

• We now define \underline{F} for $\psi \geq \tilde{c}\tilde{b}^{-5/4}$. On that interval, we choose $\underline{F} = -f(s, \psi\tilde{b}^{5/4})$, for some function f to be determined. Since $\underline{F}, \partial_Y \underline{F}$ should be continuous at $\psi = \tilde{c}\tilde{b}^{-5/4}$, we require that

$$\begin{aligned} f(s, c) &= \alpha \left[c^{2/3} \tilde{b}^{1/6} - c^{1/3} \tilde{b}^{7/12} \right] =: g_1(s), \\ \partial_\zeta f(s, c) &= \frac{\alpha}{3} \left[2c^{-1/3} \tilde{b}^{1/6} - c^{-2/3} \tilde{b}^{7/12} \right] =: g_2(s). \end{aligned}$$

As a consequence, we choose

$$f(s, \zeta) := (g_1(s) + g_2(s)(\zeta - c)) \chi(\zeta) + H(\zeta),$$

where $H \in \mathcal{C}^2 \cap W^{2,\infty}(\mathbb{R})$ is strictly increasing on $[c, +\infty[$ and such that $H(c) = H'(c) = 0$, and $\chi \in \mathcal{C}_0^\infty(\mathbb{R})$ is a cut-off function. We make the following additional assumptions: there exists $c'' > c' > c$ such that $\chi \equiv 1$ on $[c, c']$, $\chi \equiv 0$ on $[c'', +\infty[$, and $H''(\zeta) \leq -1$ for $\zeta \in [c', c'']$, $H''(\zeta) \leq 0$ and $2 \leq H(\zeta) \leq 4$ for $\zeta \geq c''$. With this choice, and recalling that $W \geq \bar{C}c^{4/3}\tilde{b}^{-5/3}$ for $\psi \geq \tilde{c}\tilde{b}^{-5/4}$ for some universal constant \bar{C} , we have

$$\begin{aligned} & \partial_s \underline{F} - \frac{1}{2W} \underline{F}(\underline{F} + 2) + \frac{3b}{2} \psi \partial_\psi \underline{F} - \sqrt{W} \partial_{\psi\psi} \underline{F} \\ & \leq - \left[g_1'(s) + g_2'(s)(\zeta - c) + \frac{1}{4} b \zeta g_2(s) \right] \chi(\zeta) \\ & \quad + \bar{C} c^{-4/3} \tilde{b}^{5/3} ((g_1(s) + g_2(s)(\zeta - c)) \chi(\zeta) + H(\zeta)) \\ & \quad - \frac{1}{4} b \zeta H'(\zeta) + \sqrt{W} \tilde{b}^{5/2} \partial_\zeta^2 H \\ & \quad + (g_1(s) + g_2(s)(\zeta - c)) \left(-\frac{1}{4} b \zeta \chi'(\zeta) + \sqrt{W} \tilde{b}^{5/2} \chi''(\zeta) \right) + 2g_2(s) \sqrt{W} \tilde{b}^{5/2} \chi'(\zeta). \end{aligned}$$

We now prove that the right-hand side of the above inequality is non-positive by looking separately at the zones $(c'', +\infty)$, $[c', c'']$ and $[c, c']$:

- For $\zeta \geq c''$, we have $\chi(\zeta) = 0$, and therefore

$$\begin{aligned} & \partial_s \underline{F} - \frac{1}{2W} \underline{F}(\underline{F} + 2) + \frac{3b}{2} \psi \partial_\psi \underline{F} - \sqrt{W} \partial_{\psi\psi} \underline{F} \\ & = \frac{1}{2W} H(\zeta)(2 - H(\zeta)) - \frac{1}{4} b \zeta H'(\zeta) + \sqrt{W} \tilde{b}^{5/2} H''. \end{aligned}$$

The assumptions $H'(\zeta) \geq 0$, $H''(\zeta) \leq 0$, $H(\zeta) \geq 2$ on $(c'', +\infty)$ ensure that the right-hand side is non-positive on this interval.

- For $\zeta \in [c', c'']$, we have $H''(\zeta) \leq -1$, and without loss of generality, we also assume

that $H'(\zeta) \geq 1$ on this interval. It follows that

$$\begin{aligned} & \partial_s \underline{F} - \frac{1}{2W} \underline{F}(\underline{F} + 2) + \frac{3b}{2} \psi \partial_\psi \underline{F} - \sqrt{W} \partial_{\psi\psi} \underline{F} \\ & \leq \bar{C} \left(\alpha b \tilde{b}^{1/6} c^{-1/3} c' + c^{-4/3} \tilde{b}^{5/3} \sup_{[c', c'']} H \right) \\ & \quad - \frac{1}{4} c' b - \bar{C} c^{2/3} \tilde{b}^{5/3} \\ & \quad + \bar{C} \|\chi\|_{W^{2,\infty}} \alpha \tilde{b}^{11/6} c^{1/3} \max(c', 1). \end{aligned}$$

It is clear that the term $-1/4c'b$ dominates all others for s_0 sufficiently large.

- For $\zeta \in [c, c']$, the computation is slightly more complicated because we expect that $H''(\zeta) \geq 0$ in a vicinity of $\zeta = c$, and $H'(c) = 0$, so we cannot use the good sign of H' in a vicinity of $\zeta = c$. However, using the formulas for g_1, g_2 , we have

$$\begin{aligned} & g_1'(s) + g_2'(s)(\zeta - c) + \frac{1}{4} b \zeta g_2(s) \\ & = \frac{\alpha b}{18} \left[\tilde{b}^{1/6} c^{-1/3} (\zeta - c) + \tilde{b}^{7/12} c^{-2/3} (2\zeta + 7c) \right] \geq 0. \end{aligned}$$

Noticing that $\tilde{b}^{5/3} \ll \tilde{b}^{7/12}$ and $\sqrt{W} \tilde{b}^{5/2} = O(\tilde{b}^{5/3})$ on the interval $\zeta \in [c, c']$, we infer that all terms are easily dominated by the above quantity, so that

$$\partial_s \underline{F} - \frac{1}{2W} \underline{F}(\underline{F} + 2) + \frac{3b}{2} \psi \partial_\psi \underline{F} - \sqrt{W} \partial_{\psi\psi} \underline{F} \leq 0$$

in this region as well.

The assumptions on the initial data also ensure that $\underline{F}(s_0, \psi) \leq F(s_0, \psi)$ for $\psi \geq cs_0^{5/4}$. The result follows.

E Proof of Lemma 4.11

Recall that

$$\bar{C}_{a,\mu} := 4 \sup_{r>0} \varphi(r, a, \mu),$$

where

$$\varphi(r, a, \mu) := \left(\int_r^\infty \frac{Y^{-a}}{\left(\mu Y + \frac{Y^2}{2}\right)^2} dY \right) \left(\int_0^r Y^a \left(Y + \frac{Y^2}{2}\right) dY \right).$$

For all $\mu \in (0, 1)$, for all $a > 0$, $r > 0$

$$\begin{aligned} \varphi(r, a, \mu) & \leq \left(\int_r^\infty \frac{4}{Y^{4+a}} dY \right) \left(\int_0^r \left(Y^{1+a} + \frac{Y^{2+a}}{2} \right) dY \right) \\ & \leq \frac{2}{(3+a)^2} + \frac{4}{(3+a)(2+a)} \frac{1}{r} \leq \frac{2}{9} + \frac{2}{3r}. \end{aligned}$$

Let $K \geq 1$ such that

$$\frac{8}{9} + \frac{8}{3K} \leq \frac{9}{10}.$$

Then $4 \sup_{r \geq K} \varphi(r, a, \mu) \leq 9/10$, so that for all $\mu \in (0, 1)$, for all $a > 0$,

$$\bar{C}_{a, \mu} \leq \max \left(\frac{9}{10}, 4 \sup_{0 < r < K} \varphi(r, a, \mu) \right).$$

Now, for all $r \in [0, K]$, for $a, \mu > 0$,

$$\begin{aligned} \partial_a \varphi(r, a, \mu) &= - \left(\int_r^\infty \ln Y \frac{Y^{-a}}{\left(\mu Y + \frac{Y^2}{2}\right)^2} dY \right) \left(\int_0^r Y^a \left(Y + \frac{Y^2}{2}\right) dY \right) \\ &\quad + \left(\int_r^\infty \frac{Y^{-a}}{\left(\mu Y + \frac{Y^2}{2}\right)^2} dY \right) \left(\int_0^r \ln Y Y^a \left(Y + \frac{Y^2}{2}\right) dY \right). \end{aligned}$$

There exists a constant C_K such that for all $a \in (0, 1/2)$, for all $\mu \in (1/2, 1)$,

$$\int_K^\infty (1 + \ln Y) \frac{Y^{-a}}{\left(\mu Y + \frac{Y^2}{2}\right)^2} dY, \quad \int_0^K (1 + |\ln Y|) Y^a \left(Y + \frac{Y^2}{2}\right) dY \leq C_K.$$

It follows that for all $r \in [0, K]$, for all $a \in (0, 1/2)$, for all $\mu \in (1/2, 1)$,

$$\begin{aligned} |\partial_a \varphi(r, a, \mu)| &\leq C_K + C_K \sup_{0 \leq r \leq K} \left(\int_r^K \frac{|\ln Y|}{Y^{2+a}} dY \right) \left(\int_0^r Y^{1+a} dY \right) \\ &\quad + C_K \sup_{0 \leq r \leq K} \left(\int_r^K \frac{1}{Y^{2+a}} dY \right) \left(\int_0^r |\ln Y| Y^{1+a} dY \right). \end{aligned}$$

Notice that if $Y \in [0, K]$, then $|\ln Y| \leq \ln(2K) - \ln Y$. Then, performing explicit integrations by part in the integrals, we observe that there exists a constant C_K such that

$$\sup_{a \in [0, 1/2]} \sup_{\mu \in [1/2, 1]} \sup_{r \in [0, K]} |\partial_a \varphi(r, a, \mu)| \leq C_K.$$

We infer that for all $r \in [0, K]$, for all $a \in [0, 1/2]$, for all $\mu \in [1/2, 1]$,

$$\varphi(r, a, \mu) \leq \varphi(r, 0, \mu) + C_K a.$$

Let us now compute explicitly $\varphi(r, 0, \mu)$. We have

$$\frac{1}{\left(\mu Y + \frac{Y^2}{2}\right)^2} = \frac{1}{\mu} \left(\frac{1}{2\mu + Y} - \frac{1}{Y} \right) + \frac{1}{Y^2} + \frac{1}{(2\mu + Y)^2},$$

so that

$$\varphi(r, 0, \mu) = \frac{1}{\mu^2} \left(\frac{1}{\mu} \ln \left(\frac{r}{2\mu + r} \right) + \frac{1}{r} + \frac{1}{2\mu + r} \right) \left(\frac{r^2}{2} + \frac{r^3}{6} \right).$$

The function $(r, \mu) \mapsto \varphi(r, 0, \mu)$ is $W^{1,\infty}$ in $[0, K] \times [\frac{1}{2}, 1]$, and therefore $|\varphi(r, 0, \mu) - \varphi(r, 0, 1)| \leq C_K |\mu - 1|$.

A careful study of the function

$$r \mapsto \varphi(r, 0, 1) = \left(\ln \left(\frac{r}{2+r} \right) + \frac{1}{r} + \frac{1}{2+r} \right) \left(\frac{r^2}{2} + \frac{r^3}{6} \right)$$

shows that it is increasing and converges towards $2/9$ as $r \rightarrow \infty$. Eventually, we obtain

$$\begin{aligned} \bar{C}_{a,\mu} &\leq \max \left(\frac{9}{10}, 4 \sup_{0 \leq r \leq K} \varphi(r, 0, 1) + C_K a + C_K |\mu - 1| \right) \\ &\leq \max \left(\frac{9}{10}, \frac{8}{9} + C_K a + C_K |\mu - 1| \right). \end{aligned}$$

Therefore, choosing a sufficiently small and μ sufficiently close to 1, we obtain

$$\bar{C}_{a,\mu} \leq \frac{9}{10}.$$

F Proofs of Lemmas 2.14 and 2.15

Proof of the trace result (Lemma 2.14)

For any $Y \in [0, L]$, write

$$|g(Y) - g(0)| = \left| \int_0^Y \partial_Y g \right| \leq \frac{1}{\sqrt{1+a}} \left(\int_0^L (\partial_Y g)^2 Y^{-a} dY \right)^{1/2} Y^{\frac{1+a}{2}}.$$

It follows that

$$|g(0)|^2 \leq 2g(Y)^2 + 2 \left(\int_0^L (\partial_Y g)^2 Y^{-a} dY \right) Y^{1+a}.$$

Multiply the above equation by $(Y + Y^2)Y^{-a}$ and integrate over $[0, L]$. We obtain

$$L^{3-a} |g(0)|^2 \leq \bar{C} \left(\int_0^L |g(Y)|^2 (Y + Y^2) Y^{-a} dY + L^4 \left(\int_0^L (\partial_Y g)^2 Y^{-a} dY \right) \right),$$

which leads to the desired inequality. \square

Proof of the coercivity result (Lemma 2.15)

Let us consider the quantity

$$\int_0^\infty U (\partial_Y \mathcal{L}_U^2 V)^2 \tilde{w}_1, \tag{F.1}$$

where $\tilde{w}_1 = Y^{-a} (1 + s^{-\beta_1} Y)^{-m_1 - 2} = w_1 (1 + s^{-\beta_1} Y)^{-2}$. In order to prove the coercivity result, we go back to the diffusion term that is bounded from below by D_1 (plus some lower order terms), namely

$$- \int_0^\infty \partial_Y^2 \mathcal{L}_U^2 V \partial_Y^2 \mathcal{L}_U V w_1,$$

or rather, to the same integral where w_1 is replaced by \tilde{w}_1 . We set

$$\tilde{D}_1(s) := \int_0^\infty \frac{(\partial_Y^2 \mathcal{L}_U V)^2}{U^2} \tilde{w}_1 + \int_0^\infty \frac{(\partial_Y^3 \mathcal{L}_U V)^2}{U} \tilde{w}_1.$$

Note that we obviously have $\tilde{D}_1 \leq D_1$.

We recall (see the proof of Lemma 4.7 with $f = \partial_Y^2 \mathcal{L}_U V$) that for any $\delta > 0, P > 0$, provided m_1 and s_0 are sufficiently large,

$$\bar{c}\tilde{D}_1 - \delta b\tilde{E}_1 - s^{-P} - s^{-P}D_0 \leq - \int_0^\infty \partial_Y^2 \mathcal{L}_U^2 V \partial_Y^2 \mathcal{L}_U V \tilde{w}_1 \leq \bar{c}^{-1}\tilde{D}_1 + b\tilde{E}_1 + s^{-P} + s^{-P}D_0.$$

On the other hand, set $h := \mathcal{L}_U^2 V$. Then $\partial_Y^2 \mathcal{L}_U V = L_U h$, so that, using the identity $\partial_Y L_U = U \partial_Y - U_{YY} \int_0^Y$ and performing several integrations by parts,

$$\begin{aligned} & - \int_0^\infty \partial_Y^2 \mathcal{L}_U^2 V \partial_Y^2 \mathcal{L}_U V \tilde{w}_1 = \int_0^\infty \partial_Y h (\partial_Y L_U h) \tilde{w}_1 + \int_0^\infty \partial_Y h L_U h \partial_Y \tilde{w}_1 \\ & = \int_0^\infty U (\partial_Y h)^2 \tilde{w}_1 - \int_0^\infty \partial_Y h \left(\int_0^Y h \right) \tilde{w}_1 + \int_0^\infty (1 - U_{YY}) \partial_Y h \left(\int_0^Y h \right) \tilde{w}_1 \\ & \quad - \frac{1}{2} \int_0^\infty h^2 (U \partial_Y \tilde{w}_1)_Y - \int_0^\infty \partial_Y h \left(\int_0^Y h \right) U_Y \partial_Y \tilde{w}_1 \\ & = \int_0^\infty [U (\partial_Y h)^2 + h^2] \tilde{w}_1 \\ & \quad + \int_0^\infty (1 - U_{YY}) \partial_Y h \left(\int_0^Y h \right) \tilde{w}_1 - \frac{1}{2} \int_0^\infty \left(\int_0^Y h \right)^2 \partial_{YY} \tilde{w}_1 \\ & \quad + \int_0^\infty h^2 U_Y \partial_Y \tilde{w}_1 + \int_0^\infty h \left(\int_0^Y h \right) (U_{YY} - 1) \partial_Y \tilde{w}_1 \\ & \quad - \frac{1}{2} \int_0^\infty \left(\int_0^Y h \right)^2 ((1 + U_{YY}) \partial_{YY} \tilde{w}_1 + U_Y \partial_Y^3 \tilde{w}_1). \end{aligned}$$

The first term in the right-hand side is precisely (F.1). The two terms with $(U_{YY} - 1)$ in the integrand can be bounded in the same fashion as the analogous remainder terms in Lemma 4.7, and therefore satisfy, for any $\delta, P > 0$,

$$\begin{aligned} & \left| \int_0^\infty (1 - U_{YY}) \partial_Y h \left(\int_0^Y h \right) \tilde{w}_1 \right| + \left| \int_0^\infty h \left(\int_0^Y h \right) (U_{YY} - 1) \partial_Y \tilde{w}_1 \right| \\ & \leq \delta \left[D_1 + bE_1 + \int_0^\infty [U (\partial_Y h)^2 + h^2] \tilde{w}_1 \right] + s^{-P} + s^{-P}D_0. \end{aligned}$$

Using the bound $|U_{YY}| \leq M_2$ and noticing that $\partial_Y^3 \tilde{w}_1 < 0$, there remains to upper-bound

$$\int_0^\infty h^2 U_Y |\partial_Y \tilde{w}_1| + \int_0^\infty \left(\int_0^Y h \right)^2 \partial_{YY} \tilde{w}_1.$$

Notice that $|\partial_Y \tilde{w}_1| \leq C_{m_1, a} Y^{-1} \tilde{w}_1$, and $\partial_{YY} \tilde{w}_1 \leq C_{m_1, a} Y^{-2} \tilde{w}_1$. Hence, using a Hardy inequality together with the bounds on U_Y , it is sufficient to upper-bound

$$\int_0^\infty (1 + Y^{-1}) h^2 \tilde{w}_1.$$

Let us first consider the integral between 0 and 1. By a Hardy inequality, we have

$$\int_0^1 (1 + Y^{-1}) h^2 \tilde{w}_1 \leq 4 \int_0^1 h^2 \frac{dY}{Y^{1+a}} \leq C_a \int_0^1 Y^{1-a} (\partial_Y h)^2 dY.$$

Since

$$\partial_Y h = U_{YY} \int_0^Y \frac{\partial_Y^2 \mathcal{L}_U V}{U^2} + \frac{\partial_Y^3 \mathcal{L}_U V}{U},$$

we have, for $Y \in (0, 1)$,

$$|\partial_Y h|^2 \leq C_a \tilde{D}_1 Y^a + \frac{(\partial_Y^3 \mathcal{L}_U V)^2}{Y^2},$$

and therefore

$$\int_0^1 (1 + Y^{-1}) h^2 \tilde{w}_1 \leq C_a \tilde{D}_1.$$

There remains to control the integral for $Y \geq 1$. To that end, we write

$$h = L_U^{-1}(\partial_Y^2 \mathcal{L}_U V) = U_Y \int_0^Y \frac{\partial_Y^2 \mathcal{L}_U V}{U^2} + \frac{\partial_Y^2 \mathcal{L}_U V}{U}.$$

Once again, a simple Cauchy-Schwartz inequality yields

$$\left(\int_0^Y \frac{\partial_Y^2 \mathcal{L}_U V}{U^2} \right)^2 \leq \left(\int_0^\infty \frac{(\partial_Y^2 \mathcal{L}_U V)^2}{Y^2 U} w_1 \right) \left(\int_0^Y \frac{Y^2}{U^3 w_1} \right) \leq C_{m_1, a} \tilde{D}_1 (1 + Y^{-3} w_1^{-1}).$$

It follows that

$$\begin{aligned} \int_1^\infty h^2 \tilde{w}_1 &\leq 2 \int_1^\infty \frac{(\partial_Y^2 \mathcal{L}_U V)^2}{U^2} \tilde{w}_1 + C_{m_1, a} D_1 \int_1^\infty (1 + Y^{-3} w_1^{-1}) (1 + Y)^2 \tilde{w}_1 \\ &\leq 2 \tilde{D}_1 + C_{m_1, a} s^{(3-a)\beta_1} D_1 \leq C_{m_1, a} s^{(3-a)\beta_1} D_1. \end{aligned}$$

Eventually, we infer that for any $P > 0$, provided m_1 and s_0 are large enough, for any $s \geq s_0$,

$$\int_0^\infty [U(\partial_Y \mathcal{L}_U V)^2 + (\mathcal{L}_U V)^2] \tilde{w}_1 \leq C_{m_1, a} s^{(3-a)\beta_1} D_1 + bE_1 + s^{-P} + s^{-P} D_0. \quad (\text{F.2})$$

References

- [1] C. Bardos, *Existence et unicité de la solution de l'équation d'Euler en dimension deux*, J. Math. Anal. Appl. **40** (1972), 769–790.
- [2] H. Beirão da Veiga and F. Crispo, *Concerning the $W^{k,p}$ -inviscid limit for 3-D flows under a slip boundary condition*, J. Math. Fluid Mech. **13** (2011), no. 1, 117–135.
- [3] Thierry Clopeau, Andro Mikelić, and Raoul Robert, *On the vanishing viscosity limit for the 2D incompressible Navier-Stokes equations with the friction type boundary conditions*, Nonlinearity **11** (1998), no. 6, 1625–1636.
- [4] Weinan E, *Boundary layer theory and the zero-viscosity limit of the Navier-Stokes equation*, Acta Math. Sin. (Engl. Ser.) **16** (2000), no. 2, 207–218.

- [5] Weinan E and Bjorn Engquist, *Blowup of solutions of the unsteady Prandtl's equation*, Comm. Pure Appl. Math. **50** (1997), no. 12, 1287–1293.
- [6] David Gerard-Varet and Nader Masmoudi, *Well-posedness for the Prandtl system without analyticity or monotonicity*, Ann. Sci. Éc. Norm. Supér. (4) **48** (2015), no. 6, 1273–1325.
- [7] S. Goldstein, *On laminar boundary-layer flow near a position of separation*, Quart. J. Mech. Appl. Math. **1** (1948), 43–69.
- [8] Emmanuel Grenier, *On the nonlinear instability of Euler and Prandtl equations*, Comm. Pure Appl. Math. **53** (2000), no. 9, 1067–1091.
- [9] Yan Guo and Toan T. Nguyen, *Prandtl boundary layer expansions of steady Navier-Stokes flows over a moving plate*, Ann. PDE **3** (2017), no. 1.
- [10] Dragoş Iftimie and Franck Sueur, *Viscous boundary layers for the Navier-Stokes equations with the Navier slip conditions*, Arch. Ration. Mech. Anal. **199** (2011), no. 1, 145–175.
- [11] Sameer Iyer, *Global Steady Prandtl expansion over a moving boundary*, arXiv preprint arXiv:1609.05397 (2016).
- [12] ———, *Steady Prandtl boundary layer expansions over a rotating disk*, Arch. Ration. Mech. Anal. **224** (2017), no. 2.
- [13] ———, *Steady Prandtl layers over a moving boundary: Non-shear Euler flows*, arXiv preprint arXiv:1705.05936 (2017).
- [14] Tosio Kato, *Remarks on zero viscosity limit for nonstationary Navier-Stokes flows with boundary*, Seminar on nonlinear partial differential equations (Berkeley, Calif., 1983), Math. Sci. Res. Inst. Publ., vol. 2, Springer, New York, 1984, pp. 85–98.
- [15] James P. Kelliher, *Navier-Stokes equations with Navier boundary conditions for a bounded domain in the plane*, SIAM J. Math. Anal. **38** (2006), no. 1, 210–232.
- [16] James P. Kelliher, Roger Temam, and Xiaoming Wang, *Boundary layer associated with the Darcy-Brinkman-Boussinesq model for convection in porous media*, Phys. D **240** (2011), no. 7, 619–628.
- [17] Igor Kukavica, Nader Masmoudi, Vlad Vicol, and Tak Kwong Wong, *On the local well-posedness of the Prandtl and hydrostatic Euler equations with multiple monotonicity regions*, SIAM J. Math. Anal. **46** (2014), no. 6, 3865–3890.
- [18] Igor Kukavica and Vlad Vicol, *On the local existence of analytic solutions to the Prandtl boundary layer equations*, Commun. Math. Sci. **11** (2013), no. 1, 269–292.
- [19] Igor Kukavica, Vlad Vicol, and Fei Wang, *The van Dommelen and Shen singularity in the Prandtl equations*, Advances in Mathematics **307** (2017), 288–311.
- [20] L. D. Landau and E. M. Lifshitz, *Fluid mechanics*, Translated from the Russian by J. B. Sykes and W. H. Reid. Course of Theoretical Physics, Vol. 6, Pergamon Press, London-Paris-Frankfurt; Addison-Wesley Publishing Co., Inc., Reading, Mass., 1959.

- [21] Maria Carmela Lombardo, Marco Cannone, and Marco Sammartino, *Well-posedness of the boundary layer equations*, SIAM J. Math. Anal. **35** (2003), no. 4, 987–1004.
- [22] Nader Masmoudi, *The Euler limit of the Navier-Stokes equations, and rotating fluids with boundary*, Arch. Rational Mech. Anal. **142** (1998), no. 4, 375–394.
- [23] ———, *About the Hardy inequality*, An Invitation to Mathematics, Springer, 2011, pp. 165–180.
- [24] Nader Masmoudi and Frédéric Rousset, *Uniform regularity for the Navier-Stokes equation with Navier boundary condition*, Arch. Ration. Mech. Anal. **203** (2012), no. 2, 529–575.
- [25] ———, *Uniform regularity and vanishing viscosity limit for the free surface Navier-Stokes equations*, Arch. Ration. Mech. Anal. **223** (2017), no. 1, 301–417.
- [26] Nader Masmoudi and Tak Kwong Wong, *Local-in-time existence and uniqueness of solutions to the Prandtl equations by energy methods*, Comm. Pure Appl. Math. **68** (2015), no. 10, 1683–1741.
- [27] Frank Merle and Pierre Raphael, *On universality of blow-up profile for L^2 critical nonlinear Schrödinger equation*, Invent. Math. **156** (2004), no. 3, 565–672.
- [28] ———, *The blow-up dynamic and upper bound on the blow-up rate for critical nonlinear Schrödinger equation*, Ann. of Math. (2) **161** (2005), no. 1, 157–222.
- [29] O. A. Oleinik and V. N. Samokhin, *Mathematical models in boundary layer theory*, Applied Mathematics and Mathematical Computation, vol. 15, Chapman & Hall/CRC, Boca Raton, FL, 1999.
- [30] L. Prandtl, *Über flüssigkeitsbewegung bei sehr kleiner reibung*, Verhandl. 3rd Int. Math. Kongr. Heidelberg (1904) (1905), 484–491.
- [31] Marco Sammartino and Russel E. Caflisch, *Zero viscosity limit for analytic solutions, of the Navier-Stokes equation on a half-space. I. Existence for Euler and Prandtl equations*, Comm. Math. Phys. **192** (1998), no. 2, 433–461.
- [32] ———, *Zero viscosity limit for analytic solutions of the Navier-Stokes equation on a half-space. II. Construction of the Navier-Stokes solution*, Comm. Math. Phys. **192** (1998), no. 2, 463–491.
- [33] K. Stewartson, *On Goldstein’s theory of laminar separation*, Quart. J. Mech. Appl. Math. **11** (1958).
- [34] Catherine Sulem and Pierre-Louis Sulem, *The nonlinear Schrödinger equation*, Applied Mathematical Sciences, vol. 139, Springer-Verlag, New York, 1999, Self-focusing and wave collapse.
- [35] Ya-Guang Wang and Shi-Yong Zhu, *Separation of the two-dimensional unsteady prandtl boundary layers under adverse pressure gradient*, arXiv preprint arXiv:1801.10481 (2018).
- [36] Zhouping Xin, *Viscous boundary layers and their stability. I*, J. Partial Differential Equations **11** (1998), no. 2, 97–124.

- [37] V. E. Zakharov and A. B. Shabat, *Exact theory of two-dimensional self-focusing and one-dimensional self-modulation of waves in nonlinear media*, *Ž. Èksper. Teoret. Fiz.* **61** (1971), no. 1, 118–134.